arXiv:hep-ph/9912518v1 28 Dec 1999

Virtual Compton Scattering at High Energy

Zhang Chen

Advisor: Professor Alfred H. Mueller

Submitted in partial fulfillment of the
requirements for the degree
of Doctor of Philosophy
in the Graduate School of Arts and Sciences

COLUMBIA UNIVERSITY
1999

ABSTRACT

Virtual Compton Scattering at High Energy

ZHANG CHEN

In this dissertation we develop a theoretical framework in the context of perturbative QuantumChromoDynamics (pQCD) for studying non-forward scattering processes. In particular, we investigate a non-forward unequal mass virtual Compton scattering amplitude by performing the general operator product expansion (OPE) and the formal renormalization group (RG) analysis.

We discuss the general tensorial decomposition of the amplitude to obtain the invariant amplitudes in the non-forward kinematic region. We study the OPE to identify the relevant operators and their reduced matrix elements, as well as the corresponding Wilson coefficients. We find that the OPE now should be done in double moments with new moment variables. There are in the expansion new sets of leading twist operators which have overall derivatives. They mix under renormalization in a well-defined way. We compute the evolution kernels from which the anomalous dimensions for these operators can be extracted. We also obtain explicitly the lowest order Wilson coefficients.

In the high energy limit we find the explicit form of the dominantly contributing anomalous dimensions. We are then able to solve the resulting renormalization group equations (RGE) and give a prediction of the high energy behavior of the invariant amplitudes. We find that it is the same as is indicated by the conventional double leading logarithmic analysis.

Contents

List of Figures **iv**

Acknowledgments **vi**

Introduction **1**

1 Forward Scattering **11**

- 1.1 Amplitude and Kinematics 12
  - 1.1.1 Invariants of DIS 12
  - 1.1.2 Scattering Amplitude and Differential Cross-section 14
- 1.2 Structure Function 16
  - 1.2.1 Tensorial Decomposition 16
  - 1.2.2 The Structure Functions 18
- 1.3 Operator Product Expansion Analysis 20
  - 1.3.1 General Statement 21
  - 1.3.2 Renormalization Group Equations 22

1.3.3 OPE Description of DIS 24

1.4 Renormalization Group Analysis 37

1.4.1 Equations and Solutions 37

1.4.2 Relationship to DGLAP Evolution 39

1.4.3 The Anomalous Dimensions and Splitting Functions 41

1.5 High Energy Behavior 51

1.5.1 Anomalous Dimension Matrix 51

1.5.2 Dominant Moment Contribution 51

1.5.3 Double Leading Logarithmic Approximation 53

2 Decomposition of the Non-forward Amplitude 56

2.1 Kinematics 56

2.2 The Amplitude 59

2.3 Current Conservation 62

2.4 Decomposition and Invariant Amplitudes 63

3 Operator Product Expansion 70

3.1 Non-forward Operator Product Expansion 70

3.2 The Operators 73

3.3 Wilson Coefficients and Explicit OPE 76

4 Renormalization Group Analysis **82**

4.1 Renormalization Group Equation 82

4.2 Anomalous Dimensions 88

4.2.1 Quark-Quark Anomalous Dimension 88

4.2.2 Anomalous Dimension Kernels 97

4.2.3 Comments on Anomalous Dimensions 98

4.3 Lowest Order Wilson Coefficients 99

5 High Energy Limit **102**

5.1 Reduction of the Path Ordered Exponential 102

6 Conclusions and Open Problems **113**

Bibliography **115**

Appendices **121**

A Quark-Gluon Anomalous Dimension **121**

B Gluon-Gluon Anomalous Dimension **127**

C Gluon-Quark Anomalous Dimension **139**

Figures **144**

List of Figures

1 Diffractive Vector Meson Production 145

2 Deeply Inelastic Scattering 146

3 Forward Amplitude and Structure Function 147

4 Analyticity of the Forward Amplitude 148

5 Dispersion Integral Contour 149

6 Integration Region and Change of Variables 150

7 Forward Operator Product Expansion 151

8 The Lowest Order Quark Distribution Function 152

9 The Lowest Order Quark-Quark Diagram 153

10 Quark Self Energy Diagram 154

11 The Quark-Gluon Mixing Graph 155

12 Non-forward Scattering Amplitude 156

13 Symmetry of the Non-forward Scattering Amplitude 157

14 Handbag Diagrams in QCD-Parton Picture 158

15 Triangle Graphs for the Anomalous Dimensions 159

16 The Quark-Quark Diagram of the Anomalous Dimensions 160

17 The Poles and Contours of k_- Integration 161

18 The Tree Level Born Diagrams of the Non-forward Amplitude 162

19 The Gluon-Quark Diagram of the Anomalous Dimensions 163

20 The Quark-Gluon Diagram of the Anomalous Dimensions 164

21 The Gluon-Gluon Diagram of the Anomalous Dimensions 165

22 The Gluon Self-Energy Diagram 166

Acknowledgments

I am most thankful to Professor Alfred H. Mueller, my advisor, for his guidance throughout my studies at Columbia as a graduate student.

I wish to thank Professor Allan S. Blaer, for his advice and help in the development of my career, not only as a researcher, but also as an effective teacher.

I would like to thank Professor Norman Christ, Professor Frank Sciulli and Professor Michael Tuts for their support and help. Many thanks to Professor Xiangdong Ji at University of Maryland for stimulating physics discussions specially on Virtual Compton Scattering and to Anne Billups, Lalla Grimes and Roy Jerome for their administrative work.

I also wish to thank Professor T. D. Lee and all other faculty members of the Physics Department of Columbia University. Thanks to all members of the theory group during my study for the fun and stimulating atmosphere in which I am able to pursue my research. And special thanks all my friends for the happiness they bring to me.

Finally, with all my heart, I wish to thank my parents for their guidance, support and encouragement throughout my life.

Introduction

It is generally believed that *Quantum Chromodynamics* (QCD) is the fundamental theory for the strong interaction and it describes the properties of nuclear and subnuclear, or hadronic, matter from first principles. QCD is a quantum field theory that is invariant under a local non-Abelian gauge symmetry–the SU(3) color symmetry. The basic matter particles of QCD, the *quarks* (and antiquarks), interact with eight massless *gluons*, non-Abelian gauge bosons that also couple to themselves, to form confined bound states of hadrons. The SU(3) gauge symmetry uniquely determines the forms of interactions between quarks and gluons, and dictates that the theory is both color confined and asymptotically free. This results in the bifurcation of QCD into *Lattice* QCD at low energy and *Perturbative* QCD (pQCD) at high energy. For more detailed discussion on various aspects of Quantum Field Theory and QCD, see, for example, References [1, 2, 3, 4, 5, 6, 7, 8, 9].

At low energies, because the coupling of QCD is strong, the usual analytic method of perturbative expansion in quantum field theory breaks down. As a result many fundamental properties of QCD at low energies, such as quark confinement, the dynamical chiral symmetry breaking and ultimately the low energy hadron mass spectrum, remain unresolved, at least analytically. The lattice formulation of QCD [10] provides us with a new way of studying the theory non-perturbatively. By defining the quark and the gluon fields on a 4 dimensional Euclidean space-time lattice, the theory is regulated by an ultra-violet cut-off, the inverse lattice spacing. Physical quantities of interest defined through the path integral of the quantum fields can now be studied numerically, using Monte Carlo methods.

At high energies, on the other hand, perturbation theory is valid due to the smallness of the QCD coupling. Perturbative QCD has had great success in explaining and predicting results of high energy experiments, for example, from the ratio of cross sections of $e^+e^- \to \mu^+\mu^-$ to $e^+e^- \to hadrons$, to structure functions of the nucleon in deeply inelastic scattering (DIS) (e.g., see [11] and the references therein). In particular, pQCD has been successfully applied to the study of the structure of the nucleon, one of the most important frontiers in strong interaction physics. Here by structure we mean not only the traditional nuclear structure like spin structure and form factors but also, more relevant to the work in this thesis, the distribution functions of quarks and gluons, which we collectively call *partons*, inside the nucleon.

Physics processes usually involve both hard (high energy) and soft (low energy) physics. The key to a pQCD analysis is the proper separation ("factorization") between the hard part, where analytic calculation using perturbation theory is possible, and the soft part, where we have to rely on non-perturbative methods like numerical study and phenomenological modeling. By application of factorization theorems and resummation methods, and/or more formal methods like the operator product expansion (OPE) and renormalization group equation (RGE) [4, 5, 12, 13, 14], pQCD has been very successful in extracting the parton distribution functions of the nucleon and predicting their behaviors in a wide range of kinematic regions of high energy particle-particle scattering. However, there is still much work to be done, especially in extreme kinematic regions like very small and/or very large Bjorken-x (see equation 1.7), where we have to not only look for new physics but also take greater care in separating the hard scattering process where pQCD is applicable and the soft physics parts where perturbation theory breaks down.

Recently there is much interest in the analysis of non-forward scattering processes, for example, deeply virtual Compton scattering (DVCS) [15, 16] and hard diffractive electroproduction of vector mesons [17, 18, 19, 20] in DIS. This creates a new territory to explore the quark and gluon structure of the nucleon besides the traditional inclusive (forward parton distributions) and exclusive (form factors) processes. A new collection of parton distribution functions called the *skewed* parton distributions (SPDs) [1] appear in these exclusive, hard diffractive processes which contain new information on long distance physics due to the non-forwardness (see, for example, Ref. [21, 22, 23, 24]). At the same time, diffractive vector meson production also provides a new way of directly measuring the (forward) gluon density in the proton [17]. The work of this thesis is aimed at developing a theoretical frame work to properly study non-forward processes in the context of operator production expansion and renormalization group analysis [25]. The thrust of our research is obtaining the high energy behavior of these non-forward processes in general.

DVCS and Skewed Parton Distributions

The Compton process, referring to the elastic scattering of a photon off a charged object, historically provided one of the early evidences of the quantization and particle nature of the electromagnetic wave [26]. Its role in studying the structure of hadrons has been explored since the 50s with the derivation of Low's low-energy theorems [27] which assert, for instance, that at sufficiently low energy the spin-dependent part of the Compton amplitude is determined by the anomalous magnetic moment of a composite system. Going to higher-order terms in the low-energy expansion, one finds the electric and magnetic polarizabilities. In recent years experimental and theoretical works in measuring and understanding the polarizabilities of the nucleon and pion have flourished [28].

---

[1] SPD is the unified terminology used to replace the many terms like non-diagonal, non-forward, off-diagonal and off-forward from different parameterizations.

The virtual Compton amplitude for scattering a virtual photon off a hadron has become one of the basic tools in QCD to understand the short-distance behavior of the theory. The process of deeply virtual Compton scattering (DVCS) is, assuming the virtual photon is generated by inelastic lepton scattering, the Bjorken limit, i.e., the energy and momentum of the virtual photon going to infinity at the same rate, of virtual Compton scattering. The basic mechanism for DVCS is a quark absorbing the virtual photon, immediately radiating a real photon and falling back to the nucleon ground state. Ji [15, 22] was the first to introduce and study DVCS. He proposed that we can obtain from DVCS information on the so-called off-forward parton distributions (OFPD) which in this case contain new information on long distance physics.

The OFPDs have come up in different theoretical studies prior to Ji's work. In the late 1980s, Geyer and collaborators [29, 30] studied the relation between the Altarelli-Parisi evolution for parton distributions [13] and the Brodsky-Lepage evolution [31] for leading-twist meson wave functions. The "interpolating functions" introduced in Ref. [29] are essentially Ji's OFPDs. In the early 1990s, Jain and Ralston [32] studied hard processes involving hadron helicity flip in terms of an "off-diagonal transition amplitude" that involves off-forward matrix elements of bi-quark fields in the nucleon.

It was shown that the integral of this amplitude over the quark four-momentum yielded elastic form factors.

Ji [15, 21] introduced OFPDs in the study of the spin structure of the nucleon. His main observations were that the fractions of the spin carried by quarks and gluons can be determined from form factors of the QCD energy-momentum tensor, and that the latter can be extracted from the OFPDs. Furthermore, the DVCS process was proposed [15] as a practical way to measure the new distributions. Because the spin-2 twist-two quark and gluon operators are part of the QCD energy-momentum tensor and because the form factors of the energy-momentum tensor contain information about the quark and gluon contributions to the nucleon spin, DVCS provides a novel way to measure the fraction of the nucleon spin carried by the quark orbital angular momentum, a subject of great current interest [33]. The more relevant part of his work to ours is his study on the evolution and sum rules (in terms of form factors) of OFPDs. He also obtained certain estimates at low energy of the OFPDs and form factors [22] using the MIT bag model [34].

Radyushkin also studied the scaling limit of DVCS [16], and generalized the discussion to hard exclusive electroproduction processes [18, 23]. The non-perturbative information is incorporated in his double distributions $F(x, y; t)$ and non-forward distribution functions $F_\zeta(X; t)$. He discussed their spectral properties, the evolution equations they satisfy, their basic uses and some general aspects of factorization for hard exclusive processes.

A third parametrization for the non-perturbative information was proposed by Collins *et al* [19, 20]. In addition to the discussion of factorization theorem and evolution, they performed a numerical study of their non-diagonal parton distributions in leading logarithmic approximation. They found that the non-diagonal gluon distribution $x_2G(x_1, x_2, Q^2)$ can be well approximated at small x by the conventional (forward) gluon density $xG(x, Q^2)$.

It is clearly both interesting and important for us to study in depth these SPDs, because they will give us information on novel long distance physics due to the non-forwardness.

Diffractive Vector Meson Production

Conventionally, for example, in DIS, the gluon density $xG(x, Q^2)$ is measured indirectly. The structure function F_2 (the same as $\nu\mathrm{W}_2$ of, eg, equation 1.91) that comes out directly from the experimental data is (essentially) the sum of quark (and anti-quark) distribution functions while the gluonic content can only be obtained from a DGLAP type evolution analysis (see Chap. 1 and Ref. [11]). This is because gluons only have color charge and all probes in nature are color neutral due to color confinement. As a result any interaction with the gluonic content inside the nucleon has to have at least a quark loop involved and thus has to be of higher order.

On the other hand, at very small Bjorken-x, the gluonic sector dominates the high energy behavior of scattering cross sections (see Chap.1 for explanation). It is well-known that there is a sharp rise in the parton densities inside a nucleon at small x, which is driven by the gluons. While this might be an indication of the emergence of the so-called BFKL pomeron [35], higher order DGLAP evolution is able to account for most of the increase (see, for example, Ref. [11]). It is obvious that new ways of measuring the gluonic content in the nucleon, especially direct measurements, will be extremely useful.

It has been shown that the cross section for diffractive electroproduction of vector mesons can be predicted in pQCD [17, 36, 37] (see Figure 1). This process provides a novel probe of the dynamics of diffractive scattering in QCD. The prediction that the cross section is proportional to the square of the gluon density $xG(x, Q^2)$ in the hadron [17] means that we do have a process through which direct measurement of the gluon density is possible. Experimental data [38] appears to be in accord with the predictions, including an enhancement due to the rapid rise of the gluon density at small x.

The pQCD calculation leading to the above prediction in Ref. [17] was done under a double leading logarithmic approximation (DLLA)–resumming leading logarithms in both the longitudinal momentum (leading $\ln 1/x$) and the transverse momentum (leading $\ln Q^2/\Lambda^2$, where Λ is the QCD scale). For further studies we need to relax (at least one of) these two approximations and also incorporate consistently the BFKL contribution. One such possibility can be found in applying the high energy factorization developed by Catani *et al* [39], relaxing the leading logarithmic approximation (LLA) on the transverse momentum $\underline{k}$ and discussing the cross section in terms of the so-called un-integrated gluon distribution (see, for example, [40]). However, up to the time of the work in this thesis [25], there had not been a completely consistent treatment.

One of the basic problems lies in the fact that conventional forward factorization has been used on a non-forward process–because of the time-like four-momentum of the final state vector meson, diffractive vector meson production is necessarily a non-forward scattering process. It is therefore not obvious without theoretical investigation of non-forward factorization and/or OPE that it should be the forward gluon density that comes into the expression of the cross section.

The research of this thesis was indeed originally motivated by the diffractive vector meson production. The ultimate goal is a better understanding of the gluon distribution inside the nucleon, especially at small x. However, we realised that some formal theoretical development on OPE and renormalization group analysis in the non-forward cases has to be made before we can consistently attempt to relax the LLAs. In other words, we have to justify (or prove wrong) the use of forward factorization under DLLA before we can go further.

Unequal Mass Virtual Compton Scattering

This dissertation aims exactly at tackling the above problem. In this research, we formulate a general operator product expansion (OPE) description for a generic non-forward unequal mass virtual Compton scattering amplitude(see Figure 12). In order to avoid complications from dealing with meson light-cone wave functions (these have been shown by Collins *et al* [19] to be factorizable), we have replaced the vector meson vertex with another virtual photon coupling. At this point, the main goal is to study the influence of the non-forwardness on OPE, rather than the phenomenology of the vector meson production itself. DVCS and DIS are extreme kinematic cases of this generic unequal mass double virtual Compton scattering (see section 2.1).

General proofs of the DVCS factorization have been given by Radyushkin in his approach based on α-representation [16, 18, 23] and by Ji in an alternative method [41]. It has been established that factorization for DVCS is on the same footing as other well-known examples like DIS. Collins *et al* [19] have also proved the factorization theorem for general diffractive vector meson production.

Some early studies of unequal mass Compton processes can be found in Refs.

[30, 42]. The factorization property of the general two virtual photon process can be summarized beautifully in terms of Wilson's OPE. This expansion now requires operators with total derivatives [25, 30, 42] to describe the non-forward nature of the process. While it is well-known that these derivative operators contribute to the wave functions of mesons [31, 43], and that the scale evolution of total derivative operators can best be studied using conformally-symmetric operators [44], our work will concentrate on the renormalization group evolutions of these operators (and their corresponding Wilson coefficients) in the high energy (and small x) limit (see later chapters of this thesis).

Because of the non zero momentum transfer $r = p - p'$ from the initial state proton to the final state proton (Fig. 12), the scaling behavior must be different from that in a forward case because of the new kinematic degree of freedom (the non-forwardness). Indeed we will show that compared with forward scattering one has two scaling variables, or equivalently, two moment variables ω and ν (see equation 3.13). The amplitude should now be expanded in terms of double moments with respect to these two moment variables. There are associated with these double moments new sets of operators that have overall derivatives in front (see equation 3.11), which mix among themselves under renormalization via an anomalous dimension matrix. The reduced non-forward matrix elements of these operators are thus double distribution functions. However, they do not seem to have a simple probability interpretation. While formally we can go from DVCS to DIS using the same formalism and in principle we can proceed beyond DLL to next to leading order, we will focus our attention on the high energy behavior of these distribution functions. We solve their renormalization group equations at high energy, and show that in the high energy limit the gluonic double distribution function actually reduces to the conventional (forward) gluon density and the high energy behavior is the same as obtained by a conventional double leading logarithmic (DLL) analysis. This also means that we have justified the DLLA calculations [17].

The outline of the thesis is as follows. In Chapter 1 we review the standard operator product expansion and renormalization group equation analysis of forward scattering amplitudes, using DIS as an example. In Chapter 2 we define the process and the kinematics of the non-forward scattering amplitude $\mathrm{T}_{\mu\nu}$. A general tensorial decomposition of $\mathrm{T}_{\mu\nu}$ into invariant amplitudes is given. In Chapter 3 we perform a general operator product expansion of $\mathrm{T}_{\mu\nu}$, define new moment variables and give their relationship to the more conventional Bjorken type scaling variables. In Chapter 4 we write down the renormalization group equations, present the equivalence of evolution kernels for the double moments, and calculate explicitly the lowest order Wilson Coefficients. In Chapter 5 we go to the high energy limit to solve the renormalization group equations and make connection with the double leading logarithmic analysis. Chapter 6 gives conclusions and outlook for future work.

Chapter 1

Forward Scattering

In this chapter, we give a review of the traditional operator product expansion (OPE) and renormalization group (RG) analysis for a deeply inelastic scattering (DIS) of lepton and proton (e-p), which serves to lay out the framework and conventions that will be used in later discussions of non-forward scatterings. This will be the forward case that we refer and compare to in those chapters.

We start with the parton model description of DIS to present the problem and establish the kinematics. We proceed to decompose the amplitude into invariant amplitudes and define the structure functions. We then perform an OPE of the invariant amplitude and discuss the RG equation and its solution of the operators and Wilson coefficients involved. To set the conventions, we also include a review of the Light Cone(LC) gauge convention and explicitly calculate the quark-quark splitting function in the LC gauge.

Most results in this chapter concerning quantum field theory are more or less standard material and can be found in, for example, [2], [4] or [5], while more specific details of

DIS process can be found in for example, [11]. However, we will be presenting them in a different way [45]. We will generally not make explicit references to the standard literature on well-known results.

1.1 Amplitude and Kinematics

The process we consider here is an electron-proton scattering at very high energy, such as those that occur at DESY. Figure 2 depicts one such process, where the initial proton has four momentum p and spin σ while that of the initial electron is k and λ, respectively. After the scattering, the electron has four momentum k' and spin λ' while the final state of the proton is collectively labeled n.

The exchanged virtual particle, with momentum $q = k - k'$, can be either a photon, a Z or a W boson (in figure 2 it is a photon). For the purpose of our discussion, we will consider only one photon exchange, which is a good approximation at the leading level. In the kinematic region we will be working, that is, with $Q^2 = -q^2 \geq 1\ GeV^2$ and energy of the electron at tens of GeV (for example, DESY has a electron beam energy of about 30 GeV), we can neglect the mass of the electron in the following discussion, which means $k^2 = k'^2 = 0$.

1.1.1 Invariants of DIS

We first work in the rest frame of the proton to define the kinematic variables used to describe such a process. We have

$$\begin{cases} p = (M, 0, 0, 0) \\ k = (\,\epsilon,\ \ 0, 0, \epsilon) \end{cases} . \tag{1.1}$$

If the scattering angle of the electron is θ, then

$$k = (\,\epsilon', \epsilon' \sin\theta, 0, \epsilon' \cos\theta)\,, \tag{1.2}$$

thus the exchanged four momentum square is

$$Q^2 = -q^2 = -(k-k')^2 = 2k\cdot k' - k^2 - k'^2 = 2\epsilon\epsilon'(1-\cos\theta)\,. \tag{1.3}$$

The first invariant of a DIS process, Q^2, is then given by

$$Q^2 = 4\epsilon\epsilon' \sin^2\frac{\theta}{2}\,, \tag{1.4}$$

which indicates the hardness of the scattering. We can see that large Q^2 needs large ϵ, ϵ', and θ not approaching zero.

A second invariant ν is defined by

$$M\nu \equiv p\cdot q = p\cdot(k-k') = M(\epsilon - \epsilon')\,, \tag{1.5}$$

which gives

$$\nu = \epsilon - \epsilon'\,. \tag{1.6}$$

We can see that ν is the energy loss of the electron in the proton rest frame.

A third invariant, the Bjorken-x variable, is more commonly used instead of ν and is defined as

$$x_{Bj} \equiv \frac{1}{\omega} \equiv \frac{Q^2}{2M\nu} \equiv \frac{Q^2}{2p\cdot q}\,. \tag{1.7}$$

We note $0 \leq x_{Bj} \leq 1$.

Q^2 and x_{Bj} are the usual variables used to describe DIS, and sometimes we use equivalently Q^2 and ν. Although they are defined above in the photon rest frame, they are by definition invariant when we go to any other frame.

1.1.2 Scattering Amplitude and Differential Cross-section

The scattering amplitude of the DIS process in figure 2 can be written as

$$T^{(n)}_{\sigma\lambda'\lambda} = -ie\tilde{U}(k')\gamma_\mu U(k)\frac{-ig_{\mu\nu}}{q^2}\langle n^{(-)}|j_\nu(0)|p\sigma\rangle ie\,, \tag{1.8}$$

with the minus sign on the n state meaning outgoing states and j_ν the electromagnetic current given by

$$j_\nu = \sum_f e_f \tilde{q}_f \gamma_\nu q_f\,. \tag{1.9}$$

The differential cross-section is then given by

$$d\sigma = \frac{1}{4}\sum_{\lambda'\lambda}\sum_{\sigma n}|T^{(n)}_{\sigma\lambda'\lambda}|^2(2\pi)^4\delta^4(p+q-p_n)\frac{d^3k'}{2\epsilon 2\epsilon'}\,, \tag{1.10}$$

where the factor $\frac{1}{4}$ comes from averaging over initial proton and electron spin and the phase space and kinematic factor are obtained from a plane wave normalization.

To simplify the expression of $d\sigma$ we define the electron factor $l_{\mu\nu}$ and the proton factor $L_{\mu\nu}$ such that

$$\frac{d^3\sigma}{dk'^3} = \frac{1}{2\epsilon 2\epsilon'}\frac{e^4}{q^4}L^{\mu\nu}l_{\mu\nu}\,. \tag{1.11}$$

We have, for the electron part,

$$\begin{aligned} l_{\mu\nu} &\equiv \frac{1}{2}\sum_{\lambda'\lambda}[\tilde{U}_{\lambda'}(k')\gamma_\mu U_\lambda(k)]^*[\tilde{U}_{\lambda'}(k')\gamma_\nu U_\lambda(k)] \\ &= \frac{1}{2}\sum_{\lambda'\lambda}\tilde{U}_\lambda(k)\gamma_\mu U_{\lambda'}(k')\tilde{U}_{\lambda'}(k')\gamma_\nu U_\lambda(k)\,. \end{aligned} \tag{1.12}$$

Because we know that

$$\sum_\lambda U_{\lambda,\alpha}(k)\tilde{U}_{\lambda,\beta}(k) = (\not{k} + m_e)_{\alpha\beta}\,, \tag{1.13}$$

we arrive at

$$\begin{aligned} l_{\mu\nu} &= \frac{1}{2}Tr[\gamma_\mu(\not{k}' + m_e)\gamma_\nu(\not{k} + m_e)] \\ &= \frac{1}{2}Tr[\gamma_\mu \not{k}'\gamma_\nu \not{k}] \, , \end{aligned} \tag{1.14}$$

where we have taken the limit $m_e = 0$. Thus we obtain

$$l_{\mu\nu} = 2(k'_\mu k_\nu + k_\mu k'_\nu - g_{\mu\nu} k \cdot k') + O(m_e^2) \, , \tag{1.15}$$

which is determined completely from the kinematics of the process.

On the other hand, for the proton part, we have

$$\begin{aligned} L^{\mu\nu} &= \frac{1}{2}\sum_\sigma \sum_n \{\langle n^{(-)}|j^\mu(0)|p\sigma\rangle^* \langle n^{(-)}|j^\nu(0)|p\sigma\rangle (2\pi)^4 \delta^4(p+q-p_n)\} \\ &= \frac{1}{2}\sum_{\sigma,n} \langle p\sigma|j^\mu(0)|n^{(-)}\rangle \langle n^{(-)}|j^\nu(0)|p\sigma\rangle (2\pi)^4 \delta^4(p+q-p_n) \, , \end{aligned} \tag{1.16}$$

where we have used the fact that j_μ is Hermitian.

Because the presence of the δ-function, the sum of n state is not a complete set summation, but rather an on-shell summation. We rewrite the δ-function and obtain

$$\begin{aligned} L^{\mu\nu} &= \frac{1}{2}\sum_{\sigma,n} \int d^4x e^{i(p+q-p_n)\cdot x} \langle p\sigma|j^\mu(0)|n^{(-)}\rangle \langle n^{(-)}|j^\nu(0)|p\sigma\rangle \\ &= \frac{1}{2}\sum_{\sigma,n} \int d^4x e^{iq\cdot x} \langle p\sigma|j^\mu(x)|n^{(-)}\rangle \langle n^{(-)}|j^\nu(0)|p\sigma\rangle \, , \end{aligned} \tag{1.17}$$

where we have used the translation relationship

$$j_\mu(x) = e^{i\hat{p}\cdot x} j_\mu(0) e^{-i\hat{p}\cdot x} \, . \tag{1.18}$$

This translation operation is exactly analogous to the spacial translation by the three momentum operator $\hat{\vec{p}}$ or the time translation by the Hamiltonian $\hat{H}$ in quantum mechanics (see, for example, [46]). In the above, $\hat{p}$ is the 4-momentum operator where

$$\begin{cases} \hat{p} \cdot x = \hat{H} t - \hat{\vec{p}} \cdot \vec{x} \\ \hat{p} | \, Vacuum \, \rangle = 0 \\ \hat{p}_\mu |n\rangle = p_{n\mu} |n\rangle \, . \end{cases} \tag{1.19}$$

The n state summation is now of a complete set, thus by the completeness relationship we arrive at the expression of the proton factor as

$$L^{\mu\nu} = \frac{1}{2} \sum_\sigma \int d^4 x e^{iq \cdot x} \langle p\sigma | j^\mu(x) j^\nu(0) | p\sigma \rangle \, . \tag{1.20}$$

1.2 Structure Function

The structure functions of the proton are defined from the invariant amplitudes of the proton factor, which in turn are obtained by a general tensorial decomposition.

1.2.1 Tensorial Decomposition

$L_{\mu\nu}$ as a rank-2 tensor depends on the kinematic variables p and q only. Thus the general tensorial decomposition of $L_{\mu\nu}$ can be written as

$$L_{\mu\nu}(p,q) = A g_{\mu\nu} + B p_\mu p_\nu + C(p_\mu q_\nu + p_\nu q_\mu) + D q_\mu q_\nu + E(p_\mu q_\nu - p_\nu q_\mu) + F \varepsilon_{\mu\nu\rho\sigma} q^\rho p^\sigma \, , \tag{1.21}$$

where $A, B, ...$ are invariant amplitudes depending only on the invariants of the process, that is, $A = A(p \cdot q, q^2)$ and so on. We have deliberately grouped the terms into symmetric and anti-symmetric parts to make the symmetry properties of $L\mu\nu$ more manifest. The last term does not occur for an electron scattering, however, we do have to consider it if it is a neutrino scattering.

The conservation of the electromagnetic current gives (cf equation 2.26)

$$q^\mu L_{\mu\nu} = q^\nu L_{\mu\nu} = 0\,. \tag{1.22}$$

Thus we have

$$q^\mu L_{\mu\nu} = Aq_\nu + Bp\cdot qp_\nu + C(p\cdot qq_\nu + q^2p_\nu) + Dq^2q_\nu + E(p\cdot qq_\nu - q^2p_\nu) = 0\,, \tag{1.23}$$

being true for any p and q. We can vary p and q while keeping $p\cdot q$ and q^2 constant and this means that the coefficients of the individual p_ν and q_ν have to vanish. We have

$$\begin{cases} A + Cp\cdot q + Ep\cdot q + Dq^2 = 0 \\ Bp\cdot q + Cq^2 - Eq^2 = 0 \end{cases}. \tag{1.24}$$

Similarly, from

$$q^\nu L_{\mu\nu} = Aq_\mu + Bp\cdot qp_\mu + C(p\cdot qq_\mu + q^2p_\mu) + Dq^2q_\mu + E(q^2p_\mu - p\cdot qq_\mu) = 0 \tag{1.25}$$

we have

$$\begin{cases} A + Cp\cdot q - Ep\cdot q + Dq^2 = 0 \\ Bp\cdot q + Cq^2 + Eq^2 = 0 \end{cases}. \tag{1.26}$$

Combine equations 1.24 and 1.26 we arrive at

$$\begin{cases} E = 0 \\ A + Cp\cdot q + Dq^2 = 0 \\ Bp\cdot q + Cq^2 = 0 \end{cases}, \tag{1.27}$$

which eventually leads to

$$\begin{cases} C = -\frac{p\cdot q}{q^2}B \\ D = -\frac{1}{q^2}(A - \frac{(p\cdot q)^2}{q^2}B) \\ E = 0 \end{cases} . \tag{1.28}$$

Therefore, only two of the invariant amplitudes survive after considering current conservation and we have

$$L_{\mu\nu} = Ag_{\mu\nu} + Bp_\mu p_\nu - B\frac{p\cdot q}{q^2}(p_\mu q_\nu + p_\nu q_\mu) - \frac{1}{q^2}(A - \frac{(p\cdot q)^2}{q^2}B)q_\mu q_\nu \,, \tag{1.29}$$

or eventually, after collecting terms,

$$L_{\mu\nu} = A(g_{\mu\nu} - \frac{q_\mu q_\nu}{q^2}) + B\left(p_\mu p_\nu - \frac{p\cdot q}{q^2}(p_\mu q_\nu + p_\nu q_\mu) + \frac{(p\cdot q)^2}{(q^2)^2}q_\mu q_\nu\right) . \tag{1.30}$$

1.2.2 The Structure Functions

The traditional structure structure function $\mathrm{W}_{\mu\nu}$ of the proton is indeed just $L_{\mu\nu}$ with kinematic factors. Explicitly (see 1.20),

$$\frac{1}{4\pi^2}\frac{M}{E_p}\mathrm{W}_{\mu\nu}(p,q) \equiv L_{\mu\nu} = \frac{1}{2}\sum_\sigma \int d^4x e^{iq\cdot x}\langle p\sigma|j^\mu(x)j^\nu(0)|p\sigma\rangle \,. \tag{1.31}$$

And after the tensorial decomposition, the two proton structure functions W_1 and W_2, which are only functions of the kinematic invariants of the problem, are defined by

$$\begin{aligned} \mathrm{W}_{\mu\nu}(p,q) &= \mathrm{W}_1(Q^2,x)\left(-g_{\mu\nu} + \frac{q_\mu q_\nu}{q^2}\right) \\ &+ \frac{\mathrm{W}_2(Q^2,x)}{M^2}\left(p_\mu p_\nu - \frac{p\cdot q}{q^2}(p_\mu q_\nu + p_\nu q_\mu) + \frac{(p\cdot q)^2}{(q^2)^2}q_\mu q_\nu\right) . \end{aligned} \tag{1.32}$$

Now in terms of the structure functions, the differential cross-section 1.11 becomes

$$\frac{d^3\sigma}{dk'^3} = \frac{1}{4\pi^2}\frac{M}{E_p}\frac{1}{2\epsilon 2\epsilon'}\frac{e^4}{Q^4}\mathrm{W}^{\mu\nu}l_{\mu\nu}\,. \tag{1.33}$$

In calculating $\mathrm{W}^{\mu\nu}l_{\mu\nu}$ we note that $q_\mu l^{\mu\nu} = 0 + O(m_e^2)$, so we can drop all terms having explicit factor of q_μ or q_ν in $\mathrm{W}_{\mu\nu}$. We get

$$\begin{aligned}
l_{\mu\nu}\mathrm{W}^{\mu\nu} &= 2(k'_\mu k_\nu + k_\mu k'_\nu - g_{\mu\nu}k\cdot k')\,(-\mathrm{W}_1 g^{\mu\nu} + \frac{\mathrm{W}_2}{M^2}p^\mu p^\nu) \\
&= 4\mathrm{W}_1 k\cdot k' + 2\frac{\mathrm{W}_2}{M^2}(2p\cdot kp\cdot k' - M^2 k\cdot k') \\
&= 4\epsilon\epsilon'(2\mathrm{W}_1\sin^2\frac{\theta}{2} + \mathrm{W}_2\cos^2\frac{\theta}{2})\,. \qquad (1.34)
\end{aligned}$$

Thus the differential cross-section is given by

$$\frac{d^3\sigma}{dk'^3} = 4\frac{\alpha_{em}^2}{Q^4}\frac{M}{E_p}(2\mathrm{W}_1\sin^2\frac{\theta}{2} + \mathrm{W}_2\cos^2\frac{\theta}{2})\,, \tag{1.35}$$

where $\alpha_{em} \equiv \frac{e^2}{4\pi}$ is the fine structure constant.

It is conventional to give $d\sigma/dQ^2 d\nu$ rather than $d\sigma/d^3k'$, so we need the Jacobian of the variable transformation. We have

$$\begin{aligned}
d^3k' &= 2\pi d\cos\theta(\epsilon')^2 d\epsilon' \\
&= 2\pi(\epsilon')^2 \begin{vmatrix} \frac{\partial\cos\theta}{\partial Q^2} & \frac{\partial\cos\theta}{\partial\nu} \\ \frac{\partial\epsilon'}{\partial Q^2} & \frac{\partial\epsilon'}{\partial\nu} \end{vmatrix} d\nu dQ^2 = \pi(\epsilon')^2 d\nu dQ^2 \begin{vmatrix} \frac{\partial Q^2}{\partial\cos\theta} & \frac{\partial Q^2}{\partial\epsilon'} \\ \frac{\partial\nu}{\partial\cos\theta} & \frac{\partial\nu}{\partial\epsilon'} \end{vmatrix}^{-1} \\
&= \pi\frac{\epsilon'}{\epsilon}d\nu dQ^2\,. \qquad (1.36)
\end{aligned}$$

Therefore, the expression of the differential cross-section eventually becomes

$$\frac{d\sigma}{d\nu dQ^2} = 4\pi\frac{\alpha_{em}^2}{Q^4}\frac{\epsilon'}{\epsilon}\frac{M}{E_p}(2\mathrm{W}_1\sin^2\frac{\theta}{2} + \mathrm{W}_2\cos^2\frac{\theta}{2})\,. \tag{1.37}$$

In the laboratory frame where θ is the scattering angle, it is possible to fix x and Q^2 (or equivalently Q^2 and ν) and vary θ, thus experimentally W_1 and W_2 can be determined separately. Since there are more small angle experimental data, W_2 is better measured.

Theoretically, what we usually do is to relate the cross-section, in terms of the structure functions, to the imaginary part of a forward scattering amplitude [12] via an optical theorem.

To be more specific, define the forward scattering amplitude $\mathrm{T}_{\mu\nu}$ as

$$\mathrm{T}_{\mu\nu} = i4\pi^2 \frac{E_p}{M} \frac{1}{2} \sum_{\sigma} \int d^4x e^{iq\cdot x} \langle p\sigma | T j^{\mu}(x) j^{\nu}(0) | p\sigma \rangle \, , \tag{1.38}$$

we have (see 1.31)

$$\mathrm{W}_{\mu\nu} = 2Im \; \mathrm{T}_{\mu\nu} \, . \tag{1.39}$$

The forward amplitude is shown in figure 3(a) while the structure function is shown in figure 3(b), both up to kinematic factors. The only difference in the graphs is the cut in the middle in figure 3(b), which means that all the intermediate lines the cut goes through are put on the mass shell. Mathematically this is equivalent to replacing the Feynman propagators of all the intermediate particles with its corresponding on-shell δ-function and the proper factors (see section 1.4.3).

1.3 Operator Product Expansion Analysis

We will use a straight forward operator product expansion (OPE) analysis to discuss the deeply inelastic scattering (DIS) of previous sections. While similar and more complete discussions can be found in [5], we will present ours in a slightly different notation, adopted from [45].

1.3.1 General Statement

Let $\hat{O}_1(x)$ and $\hat{O}_2(x)$ be local operators built out of fundamental fields like $q, \tilde{q}$ (quark and anti-quark fields) or A_μ (gauge fields) and finite number of (covariant) derivatives at one space-time point, for example,

$$j^f_\mu(x) = \tilde{q}^f(x)\gamma_\mu q(x) \qquad ; \hat{O}_\mu(x) = \tilde{q}D_\mu(x)q(x) \; ; \ldots \,. \tag{1.40}$$

We have, in the short distance limit,

$$\hat{O}_1(x)\hat{O}_2(0) \overset{x\rightarrow 0}{\longrightarrow} \sum_{r=1}^{N} \hat{O}^{(r)}_{\mu_1\mu_2\ldots\mu_{n_r}}(0)\; E^{(r)}_{\mu_1\mu_2\ldots\mu_{n_r}}(x) + \mathit{Remainder}\,. \tag{1.41}$$

$\hat{O}^{(r)}$s are local operators which can include the identity operator $\hat{I}$ and $E^{(r)}(x)$ are the so-called Wilson coefficients, c-number functions that are usually singular in the limit of $x \rightarrow 0$.

By a dimensional analysis we can obtain the asymptotic behavior of the Wilson coefficients $E^{(r)}(x)$ at short distance. The convention we use is such that mass and energy has dimension +1, i.e., $[E] = dim(E) = [M] = dim(M) = +1$, while coordinate has dimension of -1, ie, $[x] = -1$. Thus, we have under our convention, $[q] = -\frac{3}{2}$, $[A_\mu] = -1$, and so on.

Now let d_r be the naive dimension of $\hat{O}^{(r)}$, d_1 and d_2 be those of $\hat{O}_1$ and $\hat{O}_2$, respectively, $E^{(r)}(x)$ has an x behavior/dependence, for small x, given by $(\sqrt{x^2})^{-\lambda}$, where $\lambda = d_r - d_1 - d_2$ since we must have $d_1 + d_2 = d_r - \lambda$. That is, we have

$$E^{(r)}_{\mu_1\mu_2\ldots\mu_{n_r}}(x) \overset{x\rightarrow 0}{\sim} (\sqrt{x^2})^{d_1+d_2-d_r}\,. \tag{1.42}$$

When we increase N, we can only generate new operators by adding new fields or more derivatives into the operators, these new operators must have more negative dimensions, that is, d_r decreases with increasing N, therefore, we may choose N large enough such that the remainder becomes as small as desired, namely, *remainder* $\sim (x^2)^R$ for any desired positive value R in the small x limit.

1.3.2 Renormalization Group Equations

We work in the zero quark mass limit. The operators under renormalization behaves like

$$\hat{O}_i(x,\mu^2,\alpha_\mu) = Z_i^{-1}(\frac{\mu^2}{\mu_0^2},\alpha_{\mu 0})\, \hat{O}_i(x,\mu_0^2,\alpha_{\mu 0})\,, \tag{1.43}$$

where we have explicitly shown the dependence of the operators on the renormalization scale. This is the so-called multiplicative renormalization and Z_i more generally can be a matrix when there is mixing among the operators.

Taking the logarithmic derivative of the scale μ^2 on both sides of 1.43 we have

$$\begin{aligned} \mu^2\frac{d}{d\mu^2}\hat{O}_i(x,\mu^2,\alpha_\mu) &= [\mu^2\frac{d}{d\mu^2}Z_i^{-1}(\frac{\mu^2}{\mu_0^2},\alpha_{\mu 0})]\, \hat{O}_i(x,\mu_0^2,\alpha_{\mu 0}) \\ &= -\frac{1}{Z_i}\mu^2\frac{dZ_i}{d\mu^2}\; Z_i^{-1}\, \hat{O}_i(x,\mu_0^2,\alpha_{\mu 0}) \\ &\equiv \gamma_i(\alpha_\mu)\, \hat{O}_i(x,\mu^2,\alpha_\mu)\;, \end{aligned} \tag{1.44}$$

where in the last step we have defined the so-called anomalous dimension γ_i of the operator $\hat{O}_i$ as

$$\gamma_i(\alpha_\mu) = -\frac{1}{Z_i}\mu^2\frac{dZ_i}{d\mu^2}\,. \tag{1.45}$$

Note that γ_i is a function of α_μ only. This is because from 1.45 γ_i can only be a function of μ^2 and α_μ and is dimensionless. Because we have taken all the quark masses to be zero, there is no scale to set μ, thus γ_i is a function of the coupling only. On the other hand, if $m_q \neq 0$, then $\gamma_i \to \gamma_i(\alpha_\mu, \frac{\mu^2}{m_q^2})$, complications will arise.

The operators, at the same time, obey a renormalization group equation of the form

$$\mu^2 \frac{d}{d\mu^2} \hat{O}_i(x, \mu^2, \alpha_\mu) = \gamma_i(\alpha_\mu)\, \hat{O}_i(x, \mu^2, \alpha_\mu)\ , \tag{1.46}$$

or more generally, when there is operator mixing,

$$\mu^2 \frac{d}{d\mu^2} \hat{O}_i(x, \mu^2, \alpha_\mu) = \sum_j \gamma_{ij}(\alpha_\mu)\, \hat{O}_j(x, \mu^2, \alpha_\mu)\ . \tag{1.47}$$

To establish the renormalization group equations of the Wilson coefficients, let us look at a simple case where only one term is kept in the operator product expansion, that is,

$$\hat{O}_1(x, \mu^2, \alpha_\mu) \hat{O}_2(0, \mu^2, \alpha_\mu) \xrightarrow{x \to 0} \hat{O}^{(3)}(0, \mu^2, \alpha_\mu)\ E^{(3)}(x, \mu^2, \alpha_\mu) + \mathit{Remainder}\ . \tag{1.48}$$

Taking the logarithmic derivative of μ^2 on both sides and omitting the contribution from the remainder, we have, suppressing the μ^2 and α_μ dependence,

$$(\gamma_1 + \gamma_2) \hat{O}_1(x) \hat{O}_2(0) \xrightarrow{x \to 0} \gamma_3 \hat{O}^{(3)}(0) E^{(3)} + \hat{O}^{(3)}(0) \mu^2 \frac{d}{d\mu^2} E^{(3)}\ . \tag{1.49}$$

Combined with the operator product expansion itself, we have

$$\mu^2 \frac{d}{d\mu^2} E^{(3)}(x) \equiv (\mu^2 \frac{\partial}{\partial \mu^2} + \beta \frac{\partial}{\partial \alpha_\mu}) E^{(3)}(x) = (\gamma_1 + \gamma_2 - \gamma_3) E^{(3)}(x)\ , \tag{1.50}$$

where $\beta = \beta(\alpha_\mu) = \mu^2 \frac{d\alpha_\mu}{d\mu^2}$ is the QCD beta-function.

Therefore, the renormalization group equation for the Wilson coefficient functions is

$$\left(\mu^2 \frac{\partial}{\partial \mu^2} + \beta \frac{\partial}{\partial \alpha_\mu} + \gamma_3(\alpha_\mu) - \gamma_1(\alpha_\mu) - \gamma_2(\alpha_\mu) \right)\ E^{(3)}(x, \mu^2, \alpha_\mu) = 0\ . \tag{1.51}$$

1.3.3 OPE Description of DIS

We now put the operator product expansion formalism to use on the deeply inelastic scattering process in 1.1. We work on the forward amplitude $\mathrm{T}_{\mu\nu}$ defined in 1.38 and the structure functions (and thus the cross-sections) can be obtained through 1.39. As usual we work in the so-called Bjorken limit where we have

$$\begin{cases} -q^2 = Q^2 & \text{large} \\ m\nu = p \cdot q & \text{large} \\ x_{Bj} = \frac{1}{\omega} = \frac{Q^2}{2p\cdot q} & \text{fixed .} \end{cases} \tag{1.52}$$

Forward Operator Product Expansion

We expand $Tj_\mu(x)j_\nu(0)$ as $x_\mu \to 0$. The tensor structure of $\mathrm{T}_{\mu\nu}$ can be proven as [45]

$$\begin{aligned} Tj_\mu(x)j_\nu(0) = \hat{A}_{\mu\nu}E^{(0)}(x^2)\hat{I} + \hat{A}_{\mu\nu}\sum_{i,n} F_n^{(i)}(x^2)\hat{O}^{(i,n)}_{\mu_1\cdots\mu_n}(0)x^{\mu_1}x^{\mu_2}...x^{\mu_n} \\ + \hat{B}_{\mu\nu\alpha\beta}\sum_{i,n} E_n^{(i)}(x^2)\hat{O}^{(i,n)}_{\alpha\beta;\mu_1\cdots\mu_n}(0)x^{\mu_1}x^{\mu_2}...x^{\mu_n} + Rem. \ , \end{aligned} \tag{1.53}$$

where $\hat{A}_{\mu\nu}$ and $\hat{B}_{\mu\nu\alpha\beta}$ are conserved tensor structure operators and (see also section 3.1)

$$\begin{aligned} \hat{A}_{\mu\nu} &= g_{\mu\nu}\Box - \partial_\mu\partial_\nu \\ \hat{B}_{\mu\nu\alpha\beta} &= g_{\mu\alpha}g_{\nu\beta}\Box + g_{\mu\nu}\partial_\alpha\partial_\beta - g_{\mu\alpha}\partial_\nu\partial_\beta - g_{\nu\beta}\partial_\mu\partial_\alpha \ . \end{aligned} \tag{1.54}$$

We suppose symmetric combinations in μ, ν (since from section 1.2.1, $\mathrm{T}_{\mu\nu}$, after all, is symmetric) and also we suppose that all indices in $\hat{O}$s are symmetrized (even α, β with μ_is). We note that the $\hat{O}$s are the same operator sets for both terms and that the label n is actually the angular momentum quantum number, or spin, of the corresponding operator. The term *twist* is defined as the difference between the naive dimension of $\hat{O}$ and its spin, ie, *twist* of $\hat{O}_n = [\hat{O}_n] - n$.

We need to evaluate the above OPE between two symmetric external proton states (see 1.38), however, in practice in the kinematic region where high energy particle experiments are conducted, especially in the small x_{Bj} region where most of our interest lies, we can put in some transverse momentum to make the external states asymmetric and physics will not see the difference at this level (note it is the main goal of this thesis to discuss what will happen with asymmetric external states when we go to higher level of accuracy). That is,

$$\langle p|Tj_\mu(x)j_\nu(0)|p\rangle = \lim_{r\rightarrow 0}\langle p-r|Tj_\mu(x)j_\nu(0)|p\rangle \, . \tag{1.55}$$

Therefore the identity operator $\hat{I}$ does not contribute to $\mathrm{T}_{\mu\nu}$ in the OPE. We need to work to the next terms in 1.53 and we have

$$\begin{aligned}\langle p|Tj_\mu(x)j_\nu(0)|p\rangle \quad \xrightarrow{x\rightarrow 0} \quad & (g_{\mu\nu}\Box - \partial_\mu\partial_\nu)\sum_{i,n} F_n^{(i)}(x^2)x^{\mu_1}x^{\mu_2}...x^{\mu_n}\langle p|\hat{O}^{(i,n)}_{\mu_1...\mu_n}(0)|p\rangle \\ & + (g_{\mu\alpha}g_{\nu\beta}\Box + g_{\mu\nu}\partial_\alpha\partial_\beta - g_{\mu\alpha}\partial_\nu\partial_\beta - g_{\nu\beta}\partial_\mu\partial_\alpha) \\ & \quad \sum_{i,n} E_n^{(i)}(x^2)x^{\mu_1}x^{\mu_2}...x^{\mu_n}\langle p|\hat{O}^{(i,n)}_{\alpha\beta;\mu_1...\mu_n}(0)|p\rangle \, . \end{aligned} \tag{1.56}$$

For the matrix elements $\langle p|\hat{O}^{(i,n)}_{\mu_1...\mu_n}(0)|p\rangle$, the indices can be made only from either $g_{\mu\nu}$ or p_μ because x_μ is gone, however, the contribution of the two types of indices are different. After the Fourier transformation, we know from dimensional analysis that

$$x_{\mu_i} \Rightarrow \frac{q_{\mu_i}}{q^2} \, . \tag{1.57}$$

Thus we have, for the $g_{\mu\nu}$ terms,

$$g_{\mu_i\mu_j}\frac{q_{\mu_i}}{q^2}\frac{q_{\mu_j}}{q^2} \sim \frac{1}{q^2} \, , \tag{1.58}$$

while at the same time,

$$p_{\mu_i}p_{\mu_j}\frac{q_{\mu_i}}{q^2}\frac{q_{\mu_j}}{q^2} = (\frac{p\cdot q}{q^2})^2 \sim \mathrm{O}(1)\,. \tag{1.59}$$

Therefore the $g_{\mu\nu}$ contributions are small compared with those of $p_\mu p_\nu$. So the only term we keep is the contribution from a totally symmetric (in μ_i) combination $p_{\mu_1}p_{\mu_2}...p_{\mu_n}$. It is clear that this approximation is good to $O(\frac{1}{q^2})$. That is, we write

$$\langle p|\hat{O}^{(i,n)}_{\mu_1...\mu_n}(0)|p\rangle = p_{\mu_1}p_{\mu_2}...p_{\mu_n}\,\langle p||\hat{O}^{(i)}_n||p\rangle + O(\frac{1}{q^2})\,, \tag{1.60}$$

where $\langle p||\hat{O}^{(i)}_n||p\rangle$ is the so-called reduced matrix element of the operator $\hat{O}^{(i,n)}_{\mu_1...\mu_n}$. Because the external states are on-shell protons, $\langle p||\hat{O}^{(i)}_n||p\rangle$ does not depend on kinematic variables and thus is a number depending only on i and n, the flavor and spin indices, respectively.

Note indeed this approximation is the so-called leading twist approximation because non-leading twist operators will have in their contributions extra factors of $\frac{1}{q^2}$ following similar discussion as the above.

We therefore obtain, after taking leading twist approximation,

$$\begin{aligned}\langle p|Tj_\mu(x)j_\nu(0)|p\rangle \xrightarrow{x\to 0} \sum_{i,n}\Big\{&(g_{\mu\nu}\Box - \partial_\mu\partial_\nu)(p\cdot x)^n F^{(i)}_n(x^2)\\ &+ [g_{\mu\nu}(p\cdot\partial)^2 + p_{\mu\nu}p_\nu\Box - (p_\mu\partial_\nu + p_\nu\partial_\mu)p\cdot\partial](p\cdot x)^{n-2}E^{(i)}_{n-2}(x^2)\Big\}\,.\end{aligned} \tag{1.61}$$

The forward amplitude $\mathrm{T}_{\mu\nu}$ is essentially the Fourier transform with momentum q of the matrix elements (1.38). And under the Fourier transform, it is clear that $x_\mu \Rightarrow -i\frac{\partial}{\partial q_\mu}$. Thus we can rewrite the part concerning the Fourier transform of the Wilson coefficients $E^{(i)}_n(x^2)$ in terms of logarithmic derivatives of q^2 as

$$\int d^4x\, e^{iq\cdot x}(p\cdot x)^n E^{(i)}_n(x^2) \;=\; (-ip_\mu\frac{\partial}{\partial q_\mu})^n \int d^4x\, e^{iq\cdot x}E^{(i)}_n(x^2)$$

$$\begin{aligned} &= ((-i2p\cdot q)\frac{\partial}{\partial q^2})^n\ \tilde{e}_n^{(i)}(q^2) \\ &= \left(\frac{-i2p\cdot q}{q^2}\right)^n \left(q^2\frac{\partial}{\partial q^2}\right)^n \tilde{e}_n^{(i)}(q^2)\,, \qquad (1.62)\end{aligned}$$

where $\tilde{e}_n^{(i)}$ is the Fourier transform of $E_n^{(i)}$; and we have a similar set of equations for $F_n^{(i)}$. (Note in our notation, $\left(q^2\frac{\partial}{\partial q^2}\right)^n \tilde{e}_n^{(i)}(q^2)$ is exactly the coefficient $\tilde{C}^{Ja}(Q)$ in [5] with the replacement of $n \to J$ (spin index) and $i \to a$(flavor index).)

We then define the Wilson coefficients in the momentum space $\tilde{E}_n^{(i)}$ and $\tilde{F}_n^{(i)}$ as

$$\begin{aligned}\int d^4x\, e^{iq\cdot x}(p\cdot x)^n E_n^{(i)}(x^2) &= \frac{-2i}{(Q^2)^2}\left(\frac{2p\cdot q}{Q^2}\right)^n \tilde{E}_n^{(i)}(Q^2) \equiv \frac{-2i}{Q^4}\omega^n \tilde{E}_n^{(i)}(Q^2) \\ \int d^4x\, e^{iq\cdot x}(p\cdot x)^n F_n^{(i)}(x^2) &= \frac{i}{Q^2}\left(\frac{2p\cdot q}{Q^2}\right)^n \tilde{F}_n^{(i)}(Q^2) \equiv \frac{i}{Q^2}\omega^n \tilde{F}_n^{(i)}(Q^2)\,, \qquad (1.63)\end{aligned}$$

where we have used equation 1.7. Substitute these definitions into the Fourier transform of the matrix elements and note that using integration by parts we can show that the derivatives in the conserved tensor operators (1.54) become factors of momentum q in the fashion $\partial_\mu \to iq_\mu$, we arrive at

$$\begin{aligned}\int d^4x\, e^{iq\cdot x}\langle p|Tj_\mu(x)j_\nu(0)|p\rangle = -\sum_{i,n} i\langle p||\hat{O}_n^{(i)}||p\rangle &\left\{-\left(g_{\mu\nu} - \frac{q_\mu q_\nu}{q^2}\right)\omega^n \tilde{F}_n^{(i)}(Q^2)\right. \\ &\left. + \frac{2}{Q^2}\left(p_\mu p_\nu - \frac{p_\mu q_\nu + p_\nu q_\mu}{q^2}p\cdot q + g_{\mu\nu}\frac{(p\cdot q)^2}{q^2}\right)\omega^{n-2}\tilde{E}_{n-2}^{(i)}(Q^2)\right\}\,. \qquad (1.64)\end{aligned}$$

Rewriting the last term in the parenthesis in front of $\tilde{E}_{n-2}^{(i)}$ as

$$g_{\mu\nu}\frac{(p\cdot q)^2}{q^2} = (g_{\mu\nu} - \frac{q_\mu q_\nu}{q^2} + \frac{q_\mu q_\nu}{q^2})\frac{(p\cdot q)^2}{q^2}\,, \qquad (1.65)$$

we obtain

$$\begin{aligned}&\int d^4x\, e^{iq\cdot x}\langle p|Tj_\mu(x)j_\nu(0)|p\rangle \\ &= -\sum_{i,n} i\langle p||\hat{O}_n^{(i)}||p\rangle\left\{-\left(g_{\mu\nu} - \frac{q_\mu q_\nu}{q^2}\right)\omega^n\left(\tilde{F}_n^{(i)}(Q^2) - \frac{2}{Q^2\omega^2}\frac{(p\cdot q)^2}{q^2}\tilde{E}_{n-2}^{(i)}(Q^2)\right)\right. \\ &\qquad \left. + \left(p_\mu p_\nu - \frac{p_\mu q_\nu + p_\nu q_\mu}{q^2}p\cdot q + \frac{q_\mu q_\nu}{q^2}\frac{(p\cdot q)^2}{q^2}\right)\frac{2}{Q^2}\omega^{n-2}\tilde{E}_{n-2}^{(i)}(Q^2)\right\}\,. \qquad (1.66)\end{aligned}$$

From equation 1.7 we have

$$-\frac{2}{Q^2\omega^2}\frac{(p\cdot q)^2}{q^2}=\frac{1}{2}\,, \tag{1.67}$$

thus

$$\begin{aligned}&\int d^4x\, e^{iq\cdot x}\langle p|Tj_\mu(x)j_\nu(0)|p\rangle\\&=-\sum_{i,n} i\langle p||\hat{O}_n^{(i)}||p\rangle\left\{-\left(g_{\mu\nu}-\frac{q_\mu q_\nu}{q^2}\right)\omega^n(\tilde{F}_n^{(i)}(Q^2)+\frac{1}{2}\tilde{E}_{n-2}^{(i)}(Q^2))\right.\\&\qquad\left.+\frac{1}{M\nu}\left(p_\mu p_\nu-\frac{p\cdot q}{q^2}(p_\mu q_\nu+p_\nu q_\mu)+\frac{(p\cdot q)^2}{(q^2)^2}q_\mu q_\nu\right)\omega^{n-1}\tilde{E}_{n-2}^{(i)}(Q^2)\right\}\end{aligned} \tag{1.68}$$

Recall the definition of $\mathrm{T}_{\mu\nu}$, equation 1.38, and the tensorial decomposition of the structure function, equation 1.32, we obtain the tensorial decomposition of the forward amplitude as

$$\begin{aligned}\mathrm{T}_{\mu\nu} &= i4\pi^2\frac{E_p}{M}\int d^4xe^{iq\cdot x}\langle p|Tj^\mu(x)j^\nu(0)|p\rangle\\&=\left(-g_{\mu\nu}+\frac{q_\mu q_\nu}{q^2}\right)\mathrm{T}_1+\frac{1}{M^2}\left(p_\mu p_\nu-\frac{p\cdot q}{q^2}(p_\mu q_\nu+p_\nu q_\mu)+\frac{(p\cdot q)^2}{(q^2)^2}q_\mu q_\nu\right)\mathrm{T}_2\,,\end{aligned} \tag{1.69}$$

where the forward invariant amplitude $\mathrm{T}_{1,2}$ are related to the structure functions $\mathrm{W}_{1,2}$ by

$$\mathrm{W}_1=2Im\,T_1\,;\qquad \mathrm{W}_2=2Im\,T_2\,. \tag{1.70}$$

Comparing with the expression of the operator product expansion of $\mathrm{T}_{\mu\nu}$, equation 1.68, we obtain the formula for the invariant amplitudes as

$$\begin{aligned}\mathrm{T}_1 &= \frac{4\pi^2E_p}{M}\sum_n\omega^n\sum_i\langle p||\hat{O}_n^{(i)}||p\rangle(\tilde{F}_n^{(i)}(Q^2)+\frac{1}{2}\tilde{E}_{n-2}^{(i)}(Q^2))\\ \nu\mathrm{T}_2 &= 4\pi^2E_p\sum_n\omega^{n-1}\sum_i\langle p||\hat{O}_n^{(i)}||p\rangle\,\tilde{E}_{n-2}^{(i)}(Q^2)\,.\end{aligned} \tag{1.71}$$

Dispersion Relation and Optical Theorem

Equation 1.71 is essentially a power series expansion of the invariant amplitudes. To further the computation we need the analyticity properties of T_1 and T_2

Explicitly write out the time ordered product in $\mathrm{T}_{\mu\nu}$ (suppressing the spin average) by inserting a summation over arbitrary intermediate states $|r\rangle$ we get

$$\begin{aligned}
\mathrm{T}_{\mu\nu} &= i4\pi^2 \frac{E_p}{M} \int d^4x e^{iq\cdot x} \langle p|Tj^{\mu}(x)j^{\nu}(0)|p\rangle \\
&= 4\pi^2 \frac{E_p}{M} i \sum_r \int d^4x\, e^{iq\cdot x} \{\, \theta(x_0)\langle p|j_\mu(x)|r\rangle\langle r|j_\nu(0)|p\rangle \\
&\qquad + \theta(-x_0)\langle p|j_\nu(0)|r\rangle\langle r|j_\mu(x)|p\rangle \,\} \,. \qquad (1.72)
\end{aligned}$$

By applying the translation operators (see equation 1.18) we have

$$\begin{aligned}
\mathrm{T}_{\mu\nu} &= i4\pi^2 \frac{E_p}{M} i \sum_r \int d^4x\, e^{iq\cdot x} \{\, \theta(x_0) e^{i(p-p_r)\cdot x} \langle p|j_\mu(0)|r\rangle\langle r|j_\nu(0)|p\rangle \\
&\qquad + \theta(-x_0) e^{-i(p-p_r)\cdot x} \langle p|j_\nu(0)|r\rangle\langle r|j_\mu(0)|p\rangle \,\} \\
&= i4\pi^2 \frac{E_p}{M} (2\pi)^3 \sum_r \int dx_0 \{\delta^3(\vec{q}+\vec{p}-\vec{p}_r) e^{i(q_0+p_0-p_{r,0})\cdot x_0} \theta(x_0)\langle p|j_\mu(0)|r\rangle\langle r|j_\nu(0)|p\rangle \\
&\qquad + \delta^3(\vec{q}-\vec{p}+\vec{p}_r) e^{i(q_0-p_0+p_{r,0})\cdot x_0} \theta(-x_0)\langle p|j_\nu(0)|r\rangle\langle r|j_\mu(0)|p\rangle \,\} \\
&= -4\pi^2 \frac{E_p}{M} (2\pi)^3 \sum_r \left(\frac{\delta^3(\vec{q}+\vec{p}-\vec{p}_r)\langle p|j_\mu(0)|r\rangle\langle r|j_\nu(0)|p\rangle}{q_0+p_0-p_{r,0}+i\epsilon} \right. \\
&\qquad \left. - \frac{\delta^3(\vec{q}-\vec{p}+\vec{p}_r)\langle p|j_\nu(0)|r\rangle\langle r|j_\mu(0)|p\rangle}{q_0-p_0+p_{r,0}-i\epsilon} \right) \,, \qquad (1.73)
\end{aligned}$$

where in the second step we have performed the spacial integration that yields the 3-d momentum δ-function. In the last step when we perform the time integration we have put in the proper damping factors $\pm\epsilon$ in the exponent to assure the convergence at the boundaries in an adiabatic approximation.

To visualize the analytic properties of $\mathrm{T}_{\mu\nu}$, again we go to the rest frame of the proton where $p \cdot q = Mq_0 = M\nu$. The independent variables $\mathrm{T}_{\mu\nu}$ depends on are p

and q while for the invariant amplitudes the independent variables are actually Q^2 and ω, with now (see equation 1.7)

$$\omega = \frac{2p \cdot q}{Q^2} = \frac{2M}{Q^2} q_0 \,. \tag{1.74}$$

Thus, for fixed Q^2, singularities in ω are the same as singularities in q_0. Note this result is covariant–it is just for the purpose of visualization that we need to choose a frame.

From equation 1.73 we have singularities in q_0 at (i) $q_0 + p_0 = p_{r,0} \equiv E_r$ and (ii) $q_0 - p_0 = -E_r$. We shall discuss them one by one.

(i) The first set of poles in q_0 occur when

$$(q_0 + p_0)^2 = E_r^2 = M_r^2 + \vec{p}_r^{\,2} = M_r^2 + (\vec{q} + \vec{p})^2 \,, \tag{1.75}$$

where in the last step we have used the spacial δ-function. M_r is the invariant mass of the intermediate state $|r\rangle$. We therefore have

$$M_r^2 = (q + p)^2 = -Q^2 + 2p \cdot q + M^2 \,, \tag{1.76}$$

and the corresponding poles in ω are

$$\omega_r = 1 + \frac{M_r^2 - M^2}{Q^2} \,. \tag{1.77}$$

(ii) The second set of poles in q_0 occur when, after similar discussion as above,

$$M_r^2 = (q - p)^2 = q^2 - 2p \cdot q + M^2 \,, \tag{1.78}$$

and the ω poles are at

$$\omega_r = -1 - \frac{M_r^2 - M^2}{Q^2} \,. \tag{1.79}$$

The intermediate states $|r\rangle$ can be $|p\rangle$, $|p,\pi\rangle$, $|p,\pi,\pi\rangle$, etc. Electro-magnetic current and Baryon number conservation require that M, mass of the proton ($|p\rangle$ state), be the smallest invariant mass of all possible states involving any baryon. This means we always have $M_r \geq M$. Therefore we can clearly visualize the analytic property of $\mathrm{T}_{\mu\nu}$ on the complex ω-plane as shown in figure 4:

- $\mathrm{T}_{\mu\nu}$ has a pole at $\omega = 1$ and another one at $\omega = -1$. They correspond to an elastic scattering where $|r\rangle = |p\rangle$ and thus $M_r = M$.
- $\mathrm{T}_{\mu\nu}$ also has two branch cuts. One on the positive real ω-axis and starts from the point $\omega = 1 + \frac{(M+m_\pi)^2 - M^2}{Q^2}$, which corresponds to the lowest excited state $|p,\pi\rangle$ with a pion generated almost at rest and as such $M_r = M + m_\pi$. It extends out to infinity, or rather, as long as the collision center of mass energy is big enough to generate the states. The other one is simply a mirror reflection of this cut about the imaginary ω-axis.

It is therefore obvious that $\mathrm{T}_{\mu\nu}$ has an analytic circle of unit radius on the ω-plane and thus its Taylor expansion in ω exists as long as we are inside the unit circle.

However, the region of $0 \leq \omega \leq 1$ is not physical. The physics, namely, deeply inelastic scattering in the Bjorken limit, is happening in the kinematic region where $0 \leq x_{Bj} \leq 1$ (see section 1.1) and corresponds to $\omega \geq 1$. Thus the operator product expansion of equation 1.71, although called the OPE of the forward amplitude, can not be applied directly as it is to the physical scattering, nor the computation of structure functions and cross sections.

It turns out [12] that by deriving a dispersion relation for $\nu\mathrm{T}_2$, we can indeed correctly apply the operator product expansion to the physical situation and relate directly to the structure function $\nu\mathrm{W}_2$.

We start by writing a dispersion integral of $\nu \mathrm{T}_2$,

$$\nu \mathrm{T}_2(\omega, Q^2) = \frac{1}{2\pi i} \int_c \frac{d\omega'}{\omega' - \omega} \nu \mathrm{T}_2(\omega', Q^2) \tag{1.80}$$

where c is any coutour enclosing the origin on ω-plane and lying complete inside the unit circle (see figure 4). The analyticity of $\mathrm{T}_{\mu\nu}$ within the unit circle will guarantee its convergence.

We can continuously and analytically distort the contour c to c' as shown in figure 5, where c' crosses no cut, encloses no pole and has its boundary segments pushed to infinity. At infinity although actually $\nu \mathrm{T}_2$ approaches a constant [45], there are cancellations between $\pm\infty$ resulting in extra convergent factors and as such we can drop the integral region of the two semi-circles at infinity and obtain

$$\begin{aligned}
\nu \mathrm{T}_2(\omega, Q^2) &= \frac{1}{2\pi i} \int_{c \to c'} \frac{d\omega'}{\omega' - \omega} \nu \mathrm{T}_2(\omega', Q^2) \\
&= \frac{1}{2\pi i} \int_{1^-}^{\infty} \frac{d\omega'}{\omega' - \omega} (\nu \mathrm{T}_2(\omega' + i\epsilon, Q^2) - \nu \mathrm{T}_2(\omega' - i\epsilon, Q^2)) \\
&\quad + \frac{1}{2\pi i} \int_{-1^+}^{-\infty} \frac{d\omega'}{\omega' - \omega} (\nu \mathrm{T}_2(\omega' - i\epsilon, Q^2) - \nu \mathrm{T}_2(\omega' + i\epsilon, Q^2)) \,,
\end{aligned} \tag{1.81}$$

where ϵ is a infinitesimal positive number.

To simplify the above expression, we need the symmetry property of $\mathrm{T}_{\mu\nu}$ in ω, or equivalently, in q_0. $\mathrm{T}_{\mu\nu}$ as a time-ordered product can be represented systematically by a set of Feynman diagrams. Equation 1.73 is for real q_0 values. If we take q_0 to be complex and forget $i\epsilon$, we have an extended $\mathrm{T}^c_{\mu\nu}(q^c_0)$ on a complex plane. The usual time-ordered product is obviously obtained when we take $q_0 \to |q_0| + i\epsilon$ or $q_0 \to -|q_0| - i\epsilon$. This extended $\mathrm{T}^c_{\mu\nu}(q^c_0)$ clearly obeys

$$\mathrm{T}_{\mu\nu}(q_0) = \mathrm{T}_{\mu\nu}(-q_0) \,, \tag{1.82}$$

which can be obtained directly from equation 1.73. When we go to real values, we actually have

$$\mathrm{T}_2(\omega + i\epsilon, Q^2) = \mathrm{T}_2(-\omega - i\epsilon, Q^2)\,, \tag{1.83}$$

which means that for $\nu\mathrm{T}_2$ we have

$$\nu\mathrm{T}_2(\omega \pm i\epsilon, Q^2) = -\nu\mathrm{T}_2(-\omega \mp i\epsilon, Q^2)\,. \tag{1.84}$$

Therefore, by setting $\omega' \to -\omega'$ in the second integral of equation 1.81 and use the above we obtain

$$\begin{aligned}
\nu\mathrm{T}_2(\omega, Q^2) &= \frac{1}{2\pi i}\int_{1^-}^{\infty} \frac{d\omega'}{\omega' - \omega}(\nu\mathrm{T}_2(\omega' + i\epsilon, Q^2) - \nu\mathrm{T}_2(\omega' - i\epsilon, Q^2)) \\
&\quad + \frac{1}{2\pi i}\int_{1^-}^{\infty} \frac{d\omega'}{\omega' + \omega}(\nu\mathrm{T}_2(-\omega' - i\epsilon, Q^2) - \nu\mathrm{T}_2(-\omega' + i\epsilon, Q^2)) \\
&= \frac{1}{2\pi i}\int_{1^-}^{\infty} \frac{d\omega'}{\omega' - \omega}(\nu\mathrm{T}_2(\omega' + i\epsilon, Q^2) - \nu\mathrm{T}_2(\omega' - i\epsilon, Q^2)) \\
&\quad + \frac{1}{2\pi i}\int_{1^-}^{\infty} \frac{d\omega'}{\omega' + \omega}(-\nu\mathrm{T}_2(\omega' + i\epsilon, Q^2) + \nu\mathrm{T}_2(\omega' - i\epsilon, Q^2)) \\
&= \frac{1}{2\pi i}\int_{1^-}^{\infty} d\omega' \left(\frac{1}{\omega' - \omega} - \frac{1}{\omega' + \omega}\right)\left[\nu\mathrm{T}_2(\omega' + i\epsilon, Q^2) - \nu\mathrm{T}_2(\omega' - i\epsilon, Q^2)\right]. \\
&= \frac{1}{2\pi i}\int_{1^-}^{\infty} d\omega' \frac{2\omega}{\omega'^2 - \omega^2}\left[\nu\mathrm{T}_2(\omega' + i\epsilon, Q^2) - \nu\mathrm{T}_2^*(\omega' + i\epsilon, Q^2)\right] \\
&= \frac{\omega}{\pi i}\int_{1^-}^{\infty} \frac{d\omega'}{\omega'^2 - \omega^2} 2i\, Im\, \nu\mathrm{T}_2(\omega' + i\ eps, Q^2)\,,
\end{aligned} \tag{1.85}$$

where in the last step we have used the Hermiticity of the electro-magnetic current. Recalling equation 1.39, the optical theorem, we obtain the dispersion relation of $\nu\mathrm{T}_2$ as

$$\nu\mathrm{T}_2(\omega, Q^2) = \frac{\omega}{\pi}\int_{1^-}^{\infty} \frac{d\omega'}{\omega'^2 - \omega^2}\nu\mathrm{W}_2(\omega', Q^2)\,. \tag{1.86}$$

Note that in the above, ω should still be seen as inside the convergence circle (the unit circle) of $\nu\mathrm{T}_2$ while only ω' has the physical meaning of the inverse of Bjorken-x.

Therefore we can expand the denominator of the dispersion integrand and get

$$\begin{aligned} \nu\mathrm{T}_2(\omega, Q^2) &= \frac{\omega}{\pi}\int_{1^-}^{\infty}\frac{d\omega'}{\omega'^2}\nu\mathrm{W}_2(\omega', Q^2)\frac{1}{1-\frac{\omega^2}{\omega'^2}} \\ &= \frac{\omega}{\pi}\int_{1^-}^{\infty}\frac{d\omega'}{\omega'^2}\nu\mathrm{W}_2(\omega', Q^2)\sum_{n=0}^{\infty}\left(\frac{\omega^2}{\omega'^2}\right)^n \\ &= \sum_{n=0}^{\infty}\frac{\omega^{2n+1}}{\pi}\int_{1^-}^{\infty}\frac{d\omega'}{(\omega')^{2n+2}}\nu\mathrm{W}_2(\omega', Q^2)\,. \end{aligned} \tag{1.87}$$

Change the integration variable from ω' to $x_{Bj} = \omega'^{-1} \equiv x$ we have

$$\begin{aligned} dx &= -\frac{d\omega'}{\omega'^2} \\ \Rightarrow \frac{d\omega'}{(\omega')^{2n+2}} &= -x^{2n}dx\,, \end{aligned} \tag{1.88}$$

and thus

$$\begin{aligned} \nu\mathrm{T}_2(\omega, Q^2) &\overset{\omega\epsilon O(0)}{=} \sum_{n=0}^{\infty}\omega^{2n+1}\frac{1}{\pi}\int_0^1 dx\; x^{2n}\nu\mathrm{W}_2(x, Q^2) \\ &= \sum_{n\,even}^{\infty}\omega^{n+1}\frac{1}{\pi}\int_0^1 dx\; x^{n}\nu\mathrm{W}_2(x, Q^2)\,. \end{aligned} \tag{1.89}$$

Again we would like to stress that ω and x are completely different quantities. ω is essentially a mathematical qunatity introduced to Taylor expand the (invariant) amplitude(s) around the origin in an operator product expansion. It is the so-called *moment variable* and has its value limited between 0 and 1. On the other hand, x is the Bjorken-x, x_{Bj}, a kinematic variable of the actual physical scattering (see 1.7) with its value also limited between 0 and 1. The convolution of the structure function $\nu\mathrm{W}_2$ with the $n+1$-th power of x is called taking the n-th moment of the structure function. We can see that because of the symmetry properties of $\nu\mathrm{T}_2$, only odd moments of the structure function enters the expansion of the invariant amplitude.

Comparing equations 1.89 and 1.71 we can relate the operator product expansion of the forward amplitude with the moments of the (physical) structure function and we get

$$4\pi^2 E_p \sum_i \langle p||\hat{O}_n^{(i)}||p\rangle \; \tilde{E}_n^{(i)}(Q^2) = \frac{1}{\pi}\int_0^1 dx \; x^n \nu \mathrm{W}_2(x, Q^2) \; , \qquad n \; even \, . \tag{1.90}$$

That is, although the structure function $\nu\mathrm{W}_2$ itself can not be related directly to an expansion of local operators and their corresponding Wilson coefficients, its moments in x_{Bj} can indeed be expressed as a product of local operators and Wilson coefficients by application of operator product expansion and dispersion relations. The result, after a trivial rewriting, is

$$\int_0^1 dx \; x^n \nu \mathrm{W}_2(x, Q^2) = \frac{(2\pi)^3 E_p}{2} \sum_i \langle p||\hat{O}_n^{(i)}||p\rangle \; \tilde{E}_n^{(i)}(Q^2) \; , \qquad n \; even \; . \tag{1.91}$$

The Operators

Before we discuss the momentum evolution and renormalization group properties of the operators and their corresponding Wilson coefficients in equation 1.91, let us identify the operators themselves.

By recalling equation 1.60 we know that the dominant operators should have the smallest negative dimension since those with larger negative ones will be accompanied by extra suppressing factors of $\frac{1}{Q^2}$ at high energy. In QCD, the leading twist operators are the quark operators $\hat{O}^f_{\mu_1 \ldots \mu_n}$ and the gluon operators $\hat{O}^G_{\mu_1 \ldots \mu_n}$ as the following

$$\left\{ \begin{array}{l} \hat{O}^f_{\mu_1 \ldots \mu_n} = 2 \; \tilde{q}_f \gamma_{\mu_1} D_{\mu_2} \ldots D_{\mu_n} q_f \\ \hat{O}^G_{\mu_1 \ldots \mu_n} = -2 F^{\alpha}_{\mu_1} \mathcal{D}_{\mu_2} \ldots \mathcal{D}_{\mu_{n-1}} F_{\alpha \mu_n} \; . \end{array} \right. \tag{1.92}$$

In the quark operator, D_μ is the covariant derivative and $D_\mu = \partial_\mu - igA_\mu$ with g the QCD coupling $A_\mu = \sum_i A^i_\mu \lambda^i/2$ where λ^i is the fundamental representation of the $SU(3)$ color group and A^i_μ the gauge field. f is the flavor label and we can form singlets and octets out of the flavor indices. The Dirac indices are all symmetrized. The dimension of the quark operator is $[O_n^f] = -n-2$.

The dimension of the gluon operator is also $[O_n^G] = -n-2$. The Dirac indices are also symmetrized. In detail, O_n^G is

$$\hat{O}^G_{\mu_1 \ldots \mu_n} = -2 \sum_{i_1,\ldots,i_{n-1}} F^{i_1,\alpha_{\mu_1}} (\mathcal{D}_{\mu_2})^{i_1 i_2} (\mathcal{D}_{\mu_3})^{i_2 i_3} \ldots (\mathcal{D}_{\mu_{n-1}})^{i_{n-2} i_{n-1}} F^{i_{n-1}}_{\alpha\mu_n} \,, \tag{1.93}$$

where the i's are color indices and $\mathcal{D}_\mu^{ij} = \partial_\mu + \sum_l g f_{ilj} A_\mu^l$ with f_{ilj} the adjoint representation of the $SU(3)$ color group.

We will not give an explicit proof of equation 1.92, but rather present some motivations and explanations.

$\hat{O}^f_{\mu_1 \ldots \mu_n}$ Because the Dirac indices are symmetrized, only one gamma matrix can be used since two of them will give $g_{\mu\nu}$ terms. The other indices have to be covariant derivatives not only to preserve gauge symmetry but also to have the most efficient way to get indices except for gamma matrices. For example, in the case of $n = 2$, we can have $\tilde{q}_f \gamma_{\mu_1} q_f \tilde{q}_f \gamma_{\mu_2} q_f$, which has dimension of -6, or $\tilde{q}_f \gamma_{\mu_1} D_{\mu_2} q_f$, which has a dimension of -4. The Wilson coefficient (E function) of the former will have an extra power of $\frac{1}{Q^2}$ and thus the latter dominates.

$\hat{O}^G_{\mu_1 \ldots \mu_n}$ The gluon field $F_{\mu\nu}$ has two indices and dimension of $[F] = -2$, however, they are anti-symmetric. To symmetrize the μ_i indices we have to contract one of the two indices in each $F_{\mu\nu}$. Thus they are not as efficient as the (gluonic) covariant derivative $\mathcal{D}_\mu$'s.

1.4 Renormalization Group Analysis

Recall that the operators need renormalization and they obey renormalization group equations 1.46 and 1.47. Their corresponding Wilson coefficients obey a related equation 1.51 so that their products, which are the n moments of the physical structure function, do not depend on the renormalization scale.

1.4.1 Equations and Solutions

We define the Parton distribution functions $xP^f(x, Q^2)$s by

$$\sum_f e_f^2 \int_0^1 dx\, x^n x P^f(x, Q^2) \equiv \int_0^1 dx\, x^n \nu \mathrm{W}_2(x, Q^2) = \frac{(2\pi)^3 E_p}{2} \sum_i \langle p || \hat{O}_n^{(i)} || p \rangle \; \tilde{E}_n^{(i)}(Q^2) \; . \tag{1.94}$$

At this point xP^fs seem more like mathematical objects rather than physical ones. However, in the Parton Model of DIS, xP^f are indeed the (momentum fraction) distribution functions of quarks of flavor f inside the proton [11].

In leading order of the QCD coupling $\alpha(Q^2)$ (when Q^2 is large), only quark operators come in because photons only couple to quarks directly. Thus in equation 1.94 the $\hat{O}$'s are quark operators $\hat{O}^f$ from equation 1.92. We have chosen the normalization of $\hat{O}^f$ such that

$$(2\pi)^3 E_p \langle p | \hat{O}^f_{\mu_1 \dots \mu_n} | p \rangle = 2 p_{\mu_1 \dots \mu_n} \tag{1.95}$$

for quarks of flavor f in free field theory. On the other hand, in QCD $\hat{O}^f$ requires renormalization and so a scale μ has to be introduced. There will in general be complicated Q^2 and μ^2 dependence in the reduced matrix elements and Wilson coefficients.

To be more specific, if $\mu^2/Q^2 << 1$ then $\tilde{E}_n^f$ would have terms involving μ^2 as

$$\tilde{E}_n^f(Q^2,\mu^2) = \tilde{E}((\alpha(\mu^2)\log Q^2/\mu^2)^n) = \mathcal{C}_f + O((\alpha(\mu^2)\log Q^2/\mu^2)^n)\,. \tag{1.96}$$

If we simply choose $\mu^2 = Q^2$, then all the Q^2 dependence is in the reduced matrix elements obtained by equation 1.60 and we have

$$\tilde{E}_n^f(Q^2) = e_f^2 + O(\alpha(Q^2)) \tag{1.97}$$

under the normalization convention we use. Therefore we have, in leading order, at the renormalization scale $\mu^2 = Q^2$,

$$\sum_f e_f^2 \int_0^1 dx\, x^n x P^f(x,Q^2) = \frac{(2\pi)^3 E_p}{2} \sum_f \langle p||\hat{O}_n^f||p\rangle e_f^2\,, \tag{1.98}$$

which means that up to kinematic factors the moments of the parton distribution functions are the reduced matrix elements in operator product expansion, or explicitly,

$$\int_0^1 dx\, x^n x P^f(x,Q^2) = \frac{(2\pi)^3 E_p}{2} \langle p||\hat{O}_n^f||p\rangle_{Q^2}\,. \tag{1.99}$$

The subscript Q^2 means that the reduced matrix elements are renormalized at the scale $\mu^2 = Q^2$.

Recall that the operators obey the renormalization group equation (see equations 1.46 and 1.47), at an arbitrary scale μ,

$$(\mu^2 \frac{\partial}{\partial \mu^2} + \beta \frac{\partial}{\partial \alpha_\mu})\, \hat{O}_n^f(\mu^2,\alpha_\mu) = \gamma_n^f(\alpha_\mu)\, \hat{O}_n^f(\mu^2,\alpha_\mu)\,, \tag{1.100}$$

where γ_n^f is the anomalous dimension of the operator.

The matrix elements of the operators will obey the same equation. Furthermore, because the differences between the matrix elements and the reduced matrix elements are only kinematic factors (see equation 1.60), the reduced matrix elements and thus the n-moments of the parton distribution functions also obey the same renormalization equation, namely, we have

$$(\mu^2\frac{\partial}{\partial\mu^2}+\beta\frac{\partial}{\partial\alpha_\mu})\int_0^1 dx\,x^n xP^f(x,Q^2)=\gamma_n^f(\alpha_\mu)\int_0^1 dx\,x^n xP^f(x,Q^2)\,. \tag{1.101}$$

The solution to the renormalization group equation 1.100 is (formally when we have operator mixing and an anomalous dimension matrix)

$$\hat{O}_n^f(\mu^2,\alpha_\mu)=e^{-\int_{\alpha_\mu}^{\alpha_{\mu 0}}d\alpha'\frac{\gamma_n^f(\alpha')}{\beta(\alpha')}}\hat{O}_n^f(\mu_0^2,\alpha_{\mu 0})\,, \tag{1.102}$$

and the moments of parton distribution functions will obey exactly the same evolution in momentum scale.

The anomalous dimensions are calculable in perturbation theory. A lowest order calculation gives the dominant large μ^2 dependence.

1.4.2 Relationship to DGLAP Evolution

The DGLAP equations (Dokthitze, Gribov, Lipatov, Altarelli, Parisi) (for example, see [13]) state that the momentum scale Q^2 evolution of the parton distribution functions $P^f(x,Q^2)$ obeys

$$Q^2\frac{d}{dQ^2}\,xP^f(x,Q^2)=\frac{\alpha(Q^2)}{2\pi}\int_x^1\frac{dx'}{x'}\,\gamma^f(\frac{x}{x'})\,x'P^f(x',Q^2)\,, \tag{1.103}$$

where $\gamma^f(x)$ is the so-called Altarelli-Parisi splitting function of the corresponding parton distribution(s). $\gamma^f(x)$ is defined to be zero outside the range $(0,1)$. Again in general $\gamma^f(x)$ can be a matrix due to mixing among the parton distribution functions.

The DGLAP evolution equations and the renormalization group equations 1.101 are actually equivalent, with the proper identification of the splitting function with the anomalous dimension. We will not explicitly prove this (for that see, for example, [2, 4]), but rather show that given the DGLAP equation 1.103 we can obtain equation 1.101 and illustrate the explicit relationship between the splitting function and the anomalous dimension (in the leading order).

Taking the $n+1$th moment (here we adopt the convention that nth moment is take by convoluting with x^{n-1}) of both sides of equation 1.103 we obtain

$$Q^2\frac{d}{dQ^2}\int_0^1 dx\, x^n x P^f(x,Q^2) = \frac{\alpha(Q^2)}{2\pi}\int_0^1 dx\, x^n \int_x^1 \frac{dx'}{x'}\gamma^f(\frac{x}{x'})x' P^f(x',Q^2)\,. \tag{1.104}$$

Change the integration variables in the right-hand-side (rhs) from x and x' to x' and x/x' by $dx = x'\, d(x/x')$ and $dx\, dx'/x' = d(x/x')\, dx'$, so that

$$\int_0^1 dx \int_x^1 \frac{dx'}{x'} = \int_0^1 dx' \int_0^1 d\left(\frac{x}{x'}\right)\,, \tag{1.105}$$

where we have also changed the integration limits appropriately (see figure 6 for illustration). Rewrite x^n as $(x/x')^n(x')^n$ we have

$$\begin{aligned} rhs &= \frac{\alpha(Q^2)}{2\pi}\int_0^1 dx' \int_0^1 d(\frac{x}{x'})x' P^f(x',Q^2)(x')^n(\frac{x}{x'})^n\gamma^f(\frac{x}{x'}) \\ &= \frac{\alpha(Q^2)}{2\pi}\int_0^1 dx' x'^n x' P^f(x',Q^2)\int_0^1 d(\frac{x}{x'})(\frac{x}{x'})^n\gamma^f(\frac{x}{x'}) \\ &\equiv \frac{\alpha(Q^2)}{2\pi}\int_0^1 dx\, x^n\, x P^f(x,Q^2)\gamma^f_n\,, \end{aligned} \tag{1.106}$$

where we have defined the n-moment of the splitting function as

$$\gamma^f_n \equiv \int_0^1 dx\, x^n \gamma^f(x)\,. \tag{1.107}$$

By the proper identification between the $n(+1)$th moment of the splitting function and the anomalous dimension in leading order as

$$\gamma^f_n(\alpha(Q^2)) = \frac{\alpha(Q^2)}{2\pi}\gamma^f_n\,, \tag{1.108}$$

and noting that at the normalization scale $\mu^2 = Q^2$ we have $\alpha_\mu = \alpha(Q^2)$ and thus

$$Q^2 \frac{d}{dQ^2} \equiv Q^2 \frac{\partial}{\partial Q^2} + \beta \frac{\partial}{\partial \alpha} \tag{1.109}$$

we have indeed recovered equation 1.101.

1.4.3 The Anomalous Dimensions and Splitting Functions

In terms of diagrams, the operator product expansion of the DIS forward amplitude into the product of reduced matrix elements and Wilson coefficients is shown in figure 7. If we put a cut through both sides, it becomes the OPE of the structure function into Wilson coefficient and parton distribution function. Only the lowest order contribution is shown where the connecting part on the right hand side is a quark operator.

The parton distribution on the bottom evolves in the momentum scale via an anomalous dimension until it reaches the scale of the hard scattering Q^2. We will continue the computation and discussion in the light cone (LC) gauge.

Review of the Light Cone Gauge

We take the convention that the 0 component is the time component of a four vector, and the $1, 2, 3$ components are the (spacial) x, y, z components, respectively.

For any four vector v, we define the light cone components

$$v_\pm \equiv \frac{1}{\sqrt{2}}(v_0 \pm v_3), \quad \underline{v} \equiv \begin{pmatrix} v_1 \\ v_2 \end{pmatrix}. \tag{1.110}$$

This means

$$v^2 \equiv v \cdot v = 2v_+v_- - \underline{v}^2, \tag{1.111}$$

while for any two 4-vectors v_1 and v_2,

$$v_1 \cdot v_2 = v_{1+}v_{2-} + v_{1-}v_{2+} - \underline{v}_1 \cdot \underline{v}_2. \tag{1.112}$$

We can then define the LC metric $g_{\alpha\beta}$ by

$$\begin{aligned} g_{+-} = g_{-+} = 1, \quad g_{++} = g_{--} = 0, \quad g_{\perp\perp} = -1, \\ g_{+\perp} = g_{\perp+} = g_{-\perp} = g_{\perp-} = 0. \end{aligned}$$

such that for $\alpha, \beta = +, -, \perp$

$$v_1 \cdot v_2 = \sum_{\alpha,\beta} g_{\alpha\beta} v_{1\alpha} v_{2\beta}. \tag{1.113}$$

The Dirac matrices γ_α with $\alpha = 0, 1, 2, 3$ can be viewed as a four vector and we can similarly define

$$\gamma_\pm \equiv \frac{1}{\sqrt{2}}(\gamma_0 \pm \gamma_3), \quad \underline{\gamma} \equiv \begin{pmatrix} \gamma_1 \\ \gamma_2 \end{pmatrix}. \tag{1.114}$$

It can be easily verified that the light cone components obey the anti-commutation relationship

$$\{\gamma_\alpha, \gamma_\beta\} = 2g_{\alpha\beta}\,. \tag{1.115}$$

In particular, we have

$$\begin{aligned} \gamma_+^2 = \gamma_-^2 = 0 \\ \gamma_+\gamma_- + \gamma_-\gamma_+ = 2 \\ \gamma_\pm\gamma_{1,2} + \gamma_{1,2}\gamma_\pm = 0 \end{aligned} \tag{1.116}$$

and therefore,

$$\begin{aligned}\gamma_+\gamma_-\gamma_+ &= 2\gamma_+ \\ \gamma_+ \not{v}\gamma_+ &= 2v_+\gamma_+\end{aligned} \tag{1.117}$$

Introducing a vector n_μ such that for any 4-vector v

$$n \cdot v = v_+ \,. \tag{1.118}$$

This means

$$n_- = 1\,, \qquad n_+ = n_\perp = 0\,, \qquad \&\ n^2 = n_\mu n^\mu = 0\,. \tag{1.119}$$

n_μ is the so called LC null vector.

The light cone (LC) gauge is defined by requiring for the gauge field A,

$$n \cdot A = A_+ = 0\,. \tag{1.120}$$

The gluon propagator in the light cone gauge is

$$\mathcal{D}_{\mu\nu}(k) = \frac{-i}{k^2 + i\epsilon}\left(g_{\mu\nu} - \frac{n_\mu k_\nu + k_\mu n_\nu}{n \cdot k}\right) \equiv \frac{-i}{k^2 + i\epsilon} D_{\mu\nu}(k)\,, \tag{1.121}$$

where $D_{\mu\nu}(k)$ is the so-called light cone gluon projector. We have

$$n_\mu \mathcal{D}^{\mu\nu} = n_\nu \mathcal{D}^{\mu\nu} = 0\,. \tag{1.122}$$

The light cone gauge is also a *ghostless* gauge.

The Quark-Quark Anomalous Dimension

The diagram of the quark distribution function $xP^f(x)$ is shown in figure 8, where we take the lowest order quark-quark vertex function to be ([45])

$$\Gamma_+^{(0)} = \gamma_+ \left(\frac{k_+}{p_+}\right)^{n-1} . \tag{1.123}$$

Let us look at the first radiative corrections. The quark-quark diagram is shown in figure 9, where we have an extra on-shell gluon line.

Using standard Feynman rules we can write the value of part of the diagram within the dotted circle as

$$\Gamma_+^{(1)} = C \int \frac{d^4k}{(2\pi)^4} \gamma_\alpha \frac{i}{\not{k}} \gamma_+ \left(\frac{k_+}{p_+}\right)^{n-1} (-2\pi)\delta^+[(k_1-k)^2] \frac{i}{\not{k}} \gamma_\beta D^{\alpha\beta}(k_1-k) . \tag{1.124}$$

In the above, C is the color factor and

$$C = (ig)^2 \sum_{i,b} (T^i_{ab} T^i_{ba'}) , \tag{1.125}$$

where $T^i_{ab} = \left(\frac{\lambda^i}{2}\right)_{ab}$ is the generator of the $SU(3)$ color group. Since we have [6]

$$\sum_i (T^i)^2 = C_F I , \tag{1.126}$$

where C_F is the Casmir operator of the fundamental representation of the gauge group and for $SU(N)$

$$C_F = \frac{N^2-1}{2N} . \tag{1.127}$$

Defining the strong coupling constant $\alpha_s \equiv \alpha$ by $g^2 = 4\pi\alpha$, we have

$$C = (ig)^2 C_F \delta_{a'a} = -4\pi\alpha C_F , \tag{1.128}$$

where in the last step we have set $a = a'$.

In arriving at equation 1.124 we have also used the transition from the usual gluon propagator 1.121 to the on-shell gluon by

$$\frac{-i}{k^2 + i\epsilon} \to (-2\pi)\delta^+[k^2]\,, \tag{1.129}$$

which is consistent with the optical theorem 1.39.

Therefore we obtain

$$\Gamma_+^{(1)} = \frac{\alpha C_F}{2\pi^2} \int \frac{d^4k\delta^+[(k_1-k)^2]}{(k^2+i\epsilon)^2} \left(\frac{k_+}{p_+}\right)^{n-1} \gamma_\alpha \not{k} \gamma_+ \not{k} \gamma_\beta D^{\alpha\beta}(k_1-k)\,. \tag{1.130}$$

Note in the light cone gauge, there is only one solution to $\delta(k^2)$ because of the linearization of the light cone variable. We write the integral over d^4k in light cone variables

$$d^4k = dk_0 dk_1 dk_2 dk_3 \equiv dk_+ dk_- d^2\underline{k}\,, \tag{1.131}$$

and integrate over k_- with the δ-function to get

$$dk_-\delta[(k_1-k)^2] = dk_-\delta[2(k_1-k)_+(k_1-k)_- - (\underline{k}_1-\underline{k})^2] = \frac{1}{2(k_1-k)_+}\,. \tag{1.132}$$

Define three momentum ratio variables

$$x = \frac{k_+}{p_+}\,, \qquad x_1 = \frac{k_{1+}}{p_+}\,, \qquad \omega = \frac{k_+}{k_{1+}} = \frac{x}{x_1}\,, \tag{1.133}$$

we evaluate k^2 with $(k_1-k)^2 = 0$ and get

$$\begin{aligned} k^2 &= 2k_+k_- - \underline{k}^2 \\ &= 2k_+(k_{1-} - \frac{(\underline{k}_1-\underline{k})^2}{2(k_1-k)_+}) - \underline{k}^2 \\ &= \frac{k_+}{k_{1+}}(k_1^2 + \underline{k}_1^2) - \underline{k}^2 - \frac{k_+}{(k_1-k)_+}(\underline{k}_1-\underline{k})^2\,, \end{aligned} \tag{1.134}$$

where in the last step we have used

$$2k_+k_{1-} = \frac{k_+}{k_{1+}}2k_{1+}k_{1-} = \frac{k_+}{k_{1+}}(k_1^2 + \underline{k}_1^2)\,. \tag{1.135}$$

Expressing things in terms of ω we have

$$\begin{aligned} k^2 &= \omega k_1^2 + \omega \underline{k}_1^2 - \underline{k}^2 - \frac{\omega}{1-\omega}(\underline{k}_1 - \underline{k})^2 \\ &= -\frac{1}{1-\omega}[-\omega(1-\omega)k_1^2 + \omega(\underline{k}_1 - \underline{k})^2 - \omega(1-\omega)\underline{k}_1^2 + (1-\omega)\underline{k}^2 \\ &= -\frac{1}{1-\omega}[(\underline{k} - \omega\underline{k}_1)^2 - \omega(1-\omega)k_1^2]\,. \end{aligned} \tag{1.136}$$

Our goal is not to calculate Γ_+ completely, but only to calculate the ultraviolet divergent part of it. In evaluating the numerator of 1.130 we keep only the terms of the highest $\underline{k}$ power, which turns out to be quadratic. Note because of the δ-function, $(k_1 - k)_-$ also has a $\underline{k}^2$ contribution. The denominator is simple after we take the leading contribution, that is,

$$k^2 = -\frac{\underline{k}^2}{1-\omega}\,. \tag{1.137}$$

As for the numerator, we now need

$$\begin{aligned} \gamma^\alpha \not{k} \gamma_+ \not{k} \gamma^\beta D_{\alpha\beta} &= \gamma^\alpha(\{\not{k}, \gamma_+\} - \gamma_+ \not{k})\not{k}\gamma^\beta D_{\alpha\beta} \\ &= 2k_+\gamma^\alpha \not{k}\gamma^\beta D_{\alpha\beta} - k^2\gamma^\alpha\gamma_+\gamma^\beta D_{\alpha\beta}\,. \end{aligned} \tag{1.138}$$

The second term of the above is

$$\begin{aligned} -k^2\gamma^\alpha\gamma_+\gamma^\beta\left(g_{\alpha\beta} - \frac{n_\alpha(k_1-k)_\beta + (k_1-k)_\alpha n_\beta}{n\cdot(k_1-k)}\right) &= -k^2\gamma^\alpha\gamma_+\gamma_\alpha + 0 \\ &= 2k^2\gamma_+ = -2\gamma_+\frac{\underline{k}^2}{1-\omega}\,, \end{aligned} \tag{1.139}$$

where in the first step we used the facts that $n \cdot \gamma = \gamma_+$ and $\gamma_+^2 = 0$ while in the last step we used the anti-commuting relation 1.115. The first term of the numerator, on the other hand, can be expanded to be

$$2k_+\{\gamma_\alpha \not{k}\gamma^\alpha - \frac{1}{(k_1-k)_+}[\gamma_+ \not{k}(\not{k}_1-\not{k}) + (\not{k}_1-\not{k})\not{k}\gamma_+]\}\,. \tag{1.140}$$

We have from the δ-function

$$k_- \sim \frac{\underline{k}^2}{2(k_1-k)_+}\,, \tag{1.141}$$

and thus

$$\gamma_\alpha \not{k}\gamma^\alpha = -2\not{k} = -2\gamma_+ k_- = \gamma_+ \frac{\underline{k}^2}{(k_1-k)_+}\,. \tag{1.142}$$

At the same time,

$$\begin{aligned} \gamma_+ \not{k}(\not{k}_1-\not{k}) &= -\gamma_+ \not{k}\not{k} = -\gamma_+ k^2 \\ &= \gamma_+ \frac{\underline{k}^2}{1-\omega}\,, \end{aligned} \tag{1.143}$$

while similarly

$$(\not{k}_1-\not{k})\not{k}\gamma_+ = \gamma_+ \frac{\underline{k}^2}{1-\omega}\,. \tag{1.144}$$

Therefore the first term of the numerator is actually equal to

$$2\gamma\underline{k}^2 \frac{\omega}{1-\omega}\{1 - \frac{2}{1-\omega}\} = -2\gamma\underline{k}^2 \frac{\omega}{1-\omega}\frac{1+\omega}{1-\omega}\,. \tag{1.145}$$

Combined with the second term, we find that the divergent contribution to the numerator is

$$-2\gamma_+ \frac{\underline{k}^2}{(1-\omega)^2}[\omega(1+\omega) + (1-\omega)] = -2\gamma_+\underline{k}^2 \frac{1+\omega^2}{(1-\omega)^2}\,. \tag{1.146}$$

Rewrite the $\underline{k}$ integral as

$$\int d^2\underline{k} = 2\pi \int \underline{k} d\underline{k} = \pi \int d\underline{k}^2\,, \tag{1.147}$$

we arrive at the expression of the divergent part of $\Gamma_+^{(1)}$ as

$$\begin{aligned}\Gamma_+^{(1)} &= \frac{\alpha C_F}{2\pi}\int\frac{dk_+}{2(k_1-k)_+}\left(\frac{k_+}{p_+}\right)^{n-1}\int\frac{d\underline{k}^2}{\underline{k}^2}\gamma_+\left(\frac{k_+}{k_{1+}}\right)^{n-1}\left(\frac{k_{1+}}{p_+}\right)^{n-1}2(1+\omega^2)\\ &= \gamma_+\left(\frac{k_{1+}}{p_+}\right)^{n-1}\frac{\alpha C_F}{2\pi}\int\frac{d\underline{k}^2}{\underline{k}^2}\int d\omega\,\omega^{n-1}\frac{1+\omega^2}{1-\omega}\,. \qquad (1.148)\end{aligned}$$

Our original graph, one that is figure 3(b) plus an on-shell gluon running from the incoming quark to the outgoing quark, is not divergent in $\underline{k}^2$. The divergence we encountered arises because we let Q^2 become very large for fixed $\underline{k}^2$. This means that the cutoff for the logarithmic $\underline{k}^2$ integral is at Q^2. That is, explicitly,

$$\Gamma_+^{(1)} = \gamma_+\left(\frac{k_{1+}}{p_+}\right)^{n-1}\frac{\alpha C_F}{2\pi}\int^{Q^2}\frac{d\underline{k}^2}{\underline{k}^2}\int_0^1 d\omega\,\omega^{n-1}\frac{1+\omega^2}{1-\omega}\,, \qquad (1.149)$$

where we have also explicitly written out the limit on the ω integration.

Taking the logarithmic derivative with respect to Q^2, and recall equation 1.123, we obviously have

$$Q^2\frac{d}{dQ^2}\Gamma_+^{(1)}(x,Q^2) = \frac{\alpha C_F}{2\pi}\int_0^1 d\omega\,\omega^{n-1}\frac{1+\omega^2}{1-\omega}\Gamma_+^{(0)}(x,Q^2)\,. \qquad (1.150)$$

Remember the graphic definition of the parton distribution function and $\Gamma_+^{(0),(1)}$ as the lowest and first order factors, we realize that we indeed have

$$Q^2\frac{d}{dQ^2}\int_0^1 x^{n-1}xP^f(x,Q^2) = \frac{\alpha C_F}{2\pi}\int_0^1 x^{n-1}xP^f(x,Q^2)\int_0^1 d\omega\,\omega^{n-1}\frac{1+\omega^2}{1-\omega}\,. \qquad (1.151)$$

By exactly reversing the argument in section 1.4.2 (with x' now being x_1), it is straight forward to show that we indeed have the DGLAP evolution equation for the quark distribution function

$$Q^2\frac{d}{dQ^2}xP^f(x,Q^2) = \frac{\alpha C_F}{2\pi}\int_x^1\frac{dx_1}{x_1}\gamma(\frac{x}{x_1})xP^f(x_1,Q^2)\,, \qquad (1.152)$$

where the Altarelli-Parisi splitting function is

$$\gamma(y) = \frac{1+y^2}{1-y}\,. \tag{1.153}$$

The integral in equation 1.152 is actually divergent at the end point when $x_1 \to x$. This is an infrared divergence due to emission of soft gluons with $(k_1 - k)_+ \to 0$. This divergence should not be present in physical processes and indeed it is removed once we add the quark self-energy graphs shown in figure 10.

After virtual corrections are included, the full quark-quark splitting function that is free of singularities is

$$\gamma(x) = C_F[\frac{1+x^2}{(1-x)_+} + \frac{3}{2}\delta(1-x)]\,, \tag{1.154}$$

where the special function $(1-x)_+^{-1}$ is defined by

$$\int_0^1 \frac{f(x)dx}{(1-x)_+} = \int_0^1 dx \frac{f(x)-f(1)}{(1-x)} \tag{1.155}$$

for any function $f(x)$ that has reasonable behavior.

The Mixing of Evolution

There are additional order α terms in the Q^2-evolution equation due to operator mixing, for example, see figure 11. We thus have the necessity of considering both quark and gluon distributions.

To organize, define the flavor singlet quark distribution as

$$\Sigma(x,Q^2) = x\sum_f [P^f(x,Q^2) + P^{\bar{f}}(x,Q^2)]\,, \tag{1.156}$$

while the flavor octet distribution is

$$\Delta^{ff'}(x,Q^2) = x[P^f(x,Q^2) - P^{f'}(x,Q^2)]\,, \tag{1.157}$$

and a similarly defined function $\Delta^{\bar{f}\bar{f}'}$ for anti-quarks.

The octet distributions obey a generic DGLAP evolution equation

$$Q^2\frac{d}{dQ^2}\Delta^{ff'}(x,Q^2) = \frac{\alpha(Q^2)}{2\pi}\int_x^1 \frac{dx_1}{x_1}\gamma_{qq'}\left(\frac{x}{x_1}\right)\Delta^{ff'}(x_1,Q^2) \;+\; O(\alpha^2)\,. \tag{1.158}$$

However, the singlet distribution will mix under renormalization with the gluon distribution $G(x,Q^2)$. The DGLAP equation has a matrix form and explicitly

$$Q^2\frac{d}{dQ^2}\begin{pmatrix} \Sigma(x,Q^2) \\ G(x,Q^2) \end{pmatrix} = \frac{\alpha(Q^2)}{2\pi}\int_x^1 \frac{dx_1}{x_1}\begin{pmatrix} \gamma_{qq'}(x/x_1) & \gamma_{qg}(x/x_1) \\ \gamma_{gq}(x/x_1) & \gamma_{gg}(x/x_1) \end{pmatrix}\begin{pmatrix} \Sigma(x_1,Q^2) \\ G(x_1,Q^2) \end{pmatrix}\,. \tag{1.159}$$

Let C_F again be the Casmir operator in the fundamental representation with $C_F = 4/3$ for $SU(3)$ color group, and let C_A be the Casmir operator in the adjoint representation with $C_A = N_c = 3$ for the color group, we have the explicit expression of the four splitting functions as (eg, see [2])

$$\gamma_{qq}(x) = C_F\left[\frac{1+x^2}{(1-x)_+} + \frac{3}{2}\delta(1-x)\right] \tag{1.160}$$

$$\gamma_{gq}(x) = C_F\,\frac{1+(1-x)^2}{x} \tag{1.161}$$

$$\gamma_{qg}(x) = \frac{1}{2}\left[x^2 + (1-x)^2\right] \tag{1.162}$$

$$\gamma_{gg}(x) = 2C_A\left[\frac{x}{(1-x)_+} + \frac{1-x}{x} + x(1-x)\right] + \frac{11C_A - 2n_f}{6}\,\delta(1-x) \tag{1.163}$$

1.5 High Energy Behavior

It is more convenient to discuss the high energy behavior of the parton distribution functions when we go to the n-moment space by taking the n-th moment of the DGLAP evolution equation, or equivalently directly using the renormalization group equation from operator product expansion.

1.5.1 Anomalous Dimension Matrix

To the leading order in α we obtain, after taking the moments of the splitting functions, the anomalous dimension matrix for the singlet-gluon mixing to be

$$\gamma_n(\alpha(Q^2)) = \frac{\alpha(Q^2)}{2\pi} \begin{pmatrix} \gamma_n^{qq} & \gamma_n^{qg} \\ \gamma_n^{gq} & \gamma_n^{gg} \end{pmatrix} \tag{1.164}$$

where γ_n is the n-th moment of the corresponding $\gamma(x)$ in equation 1.160 and explicitly

$$\begin{aligned} \gamma_n^{qq} &= -\frac{2}{3}\left[1 + 4\sum_2^n \frac{1}{j} - \frac{2}{n(n+1)}\right] && (1.165) \\ \gamma_n^{qg} &= 2\frac{n^2+n+2}{n(n+1)(n+2)} && (1.166) \\ \gamma_n^{gq} &= \frac{8}{3}\frac{n^2+n+2}{n(n^2-1)} && (1.167) \\ \gamma_n^{gg} &= -3\left[\frac{1}{3} + \frac{2}{9}n_f + 4\sum_2^n \frac{1}{j} - \frac{4}{n(n-1)} - \frac{4}{(n+1)(n+2)}\right]. && (1.168) \end{aligned}$$

1.5.2 Dominant Moment Contribution

The anomalous dimension matrix elements are usually singular at some values of the moment index n (see equation 1.165). In the high energy limit, the dominant contribution at small x is from the right most pole of the anomalous dimension. We will illustrate this by looking again at the DIS structure function $\nu\mathrm{W}_2$.

Adopting the common terminology let us now call $\nu\mathrm{W}_2$ structure function F_2. The n-th moment of F_2 is defined as

$$F_2^{(n)}(Q^2) = \int_0^1 dx\, x^{n-1} F_2(x, Q^2)\,. \tag{1.169}$$

As we already established $F_2^{(n)}$ can be factorized as the product of parton distribution functions and perturbatively calculable Wilson coefficients with well defined Q^2 evolution. To predict cross-section, however, we do need directly $F_2(x, Q^2)$ itself. We can reconstruct F_2 from its moments by a Mellin transformation

$$F_2(x, Q^2) = \int_c \frac{dn}{2\pi i} e^{n \log \frac{1}{x}} F_2^{(n)}(Q^2)\,, \tag{1.170}$$

where c is a contour lies in the complex n plane parallel to the imaginary axis and to the right of all singularities.

Let us first check to see whether this is a self-consistent definition. Substitute 1.169 into equation 1.170 we obtain for the right hand side (rhs)

$$\begin{aligned}
rhs &= \int_c \frac{dn}{2\pi i} \int_0^1 dx' x'^{n-1} F_2(x', Q^2) e^{n \log \frac{1}{x}} \\
&= \int_c \frac{dn}{2\pi i} \int_0^1 \frac{dx'}{x'} e^{n \log \frac{1}{x} - n \log \frac{1}{x'}} F_2(x', Q^2) \\
&= \int_0^1 \frac{dx'}{x'} F_2(x', Q^2) \int_c \frac{dn}{2\pi i} e^{n[\log \frac{1}{x} - \log \frac{1}{x'}]} \\
&= \int_0^1 d \log x'\, F_2(x', Q^2)\, \delta\left(\log \frac{1}{x} - \log \frac{1}{x'}\right) \\
&= F_2(x, Q^2) = lhs\,,
\end{aligned} \tag{1.171}$$

where in the fourth step we have used the mathematical identity

$$\int_{L-i\infty}^{L+i\infty} \frac{dn}{2\pi i} e^{(n-c)z} = \delta(z)\,. \tag{1.172}$$

We can always distort the contour c to the left (but not crossing any poles) so that it'll pick out the residues of the n poles. Each pole n_p will eventually translate to a power factor of the Bjorken-x as $(x)^{-n_p}$. Therefore, at high energy and small-x the dominant contribution comes from the right most n pole of the anomalous dimension on the complex n-plane.

1.5.3 Double Leading Logarithmic Approximation

To illustrate the point that leading (right most) pole in moment space of the anomalous dimension dominates in the high energy limit, we will explicitly compute the high energy behavior of the structure function F_2 of deeply inelastic scattering.

According to previous analysis, the moments of the structure function obey the momentum evolution equation (see, eg, 1.102)

$$F_2^{(n)}(Q^2) = F_2^{(n)}(Q_0^2) e^{\int_{Q_0^2}^{Q^2} \frac{d\lambda^2}{\lambda^2} \gamma_n(\alpha(\lambda^2))} . \tag{1.173}$$

Thus the structure function itself is given by

$$F_2(x, Q^2) = \int_c \frac{dn}{2\pi i} F_2^{(n)}(Q_0^2) e^{n \log \frac{1}{x} + \int_{Q_0^2}^{Q^2} \frac{d\lambda^2}{\lambda^2} \gamma_n(\alpha(\lambda^2))} . \tag{1.174}$$

From equation 1.165 it is obvious that the gluon-gluon anomalous dimension has the right most pole among others at $n = 1$. Using the lowest order running coupling we can write the leading contributing term of the anomalous dimension

$$\gamma_n(\alpha(\lambda^2)) = \frac{C_A}{\pi(n-1)} \frac{1}{b_0 \log \lambda^2/\Lambda^2} , \tag{1.175}$$

where b_0 is the first coefficient of the QCD β-function.

By substitution we have

$$\begin{aligned} F_2(x,Q^2) &= \int_c \frac{dn}{2\pi i} F_2^{(n)}(Q_0^2) e^{n\log\frac{1}{x}+\frac{C_A}{\pi b_0}\log\frac{\log Q^2/\Lambda^2}{\log Q_0^2/\Lambda^2}\frac{1}{n-1}} \\ &\equiv \int_c \frac{dn}{2\pi i} F_2^{(n)}(Q_0^2) e^{an+\frac{b}{n-1}}\,, \end{aligned} \tag{1.176}$$

where we have defined at high energy,

$$\begin{aligned} a &\equiv \log\frac{1}{x} >> 1 \\ b &\equiv \frac{C_A}{\pi b_0}\log\frac{\log Q^2/\Lambda^2}{\log Q_0^2/\Lambda^2} \end{aligned} \tag{1.177}$$

The high energy limit corresponds to $n \to 1$ and $a >> 1$ so we can make a saddle point approximation for the exponent function $f(n) = an + b\frac{1}{n-1}$. We have

$$f'(n) = a - \frac{b}{(n-1)^2} = 0 \Rightarrow n_0 = 1 + \sqrt{\frac{b}{a}}\,. \tag{1.178}$$

Thus at the saddle point n_0,

$$\begin{aligned} f(n_0) &= a(\sqrt{\frac{b}{a}}+1) + \frac{b}{\sqrt{\frac{b}{a}}} = a + 2\sqrt{ab} \\ f''(n_0) &= \frac{2b}{(\sqrt{b/a})^3} = \frac{2a^{3/2}}{b^{1/2}} > 0\,, \end{aligned} \tag{1.179}$$

which means c is a fine contour. Let $i\nu = n-1$ we obtain finally,

$$\begin{aligned} F_2(x,Q^2) &= F_2^{(n)}(Q_0^2)\int_{-\infty}^{\infty}\frac{d\nu}{2\pi} e^{(a+2\sqrt{ab})} e^{-\frac{2a^{3/2}}{b^{1/2}}\nu^2} \\ &= F_2^{(n)}(Q_0^2) e^{n\log\frac{1}{x}} e^{2\sqrt{\log\frac{1}{x}\frac{C_A}{\pi b_0}\log\frac{\log Q^2/\Lambda^2}{\log Q_0^2/\Lambda^2}}} \sqrt{\frac{\pi b^{1/2}}{a^{3/2}}} \\ &= F_2^{(n)}(Q_0^2)\left(\frac{1}{x}\right)^1 e^{2\sqrt{\log\frac{1}{x}\frac{C_A}{\pi b_0}\log\frac{\log Q^2/\Lambda^2}{\log Q_0^2/\Lambda^2}}} \sqrt{\frac{\pi\sqrt{\frac{C_A}{\pi b_0}\log\frac{\log Q^2/\Lambda^2}{\log Q_0^2/\Lambda^2}}}{(\log\frac{1}{x})^{3/2}}}\,. \end{aligned} \tag{1.180}$$

This is the high energy limit of the forward DIS amplitude/structure function. The dominate small-x behavior is a power dependence on $\left(\frac{1}{x}\right)^{n_0}$ with n_0 given by a saddle point approximation. It is obvious that the value of n_0, which determines the leading high energy and small-x behavior, is dictated by the right most pole on the complex n plane. In this forward case, it is at $n = 1$ and thus we arrive at the x^{-1} leading behavior of the gluon distribution under DLLA.

Chapter 2

Decomposition of the Non-forward Amplitude

In this chapter, we discuss in detail the tensorial decomposition of the non-forward Amplitude. First we define the amplitude under discussion by laying out the kinematics of the process; then we proceed to use current conservation and the symmetry properties of the amplitude itself to decompose it into various invariant components and define the corresponding invariant amplitudes in the non-forward case.

2.1 Kinematics

The process we consider is $\gamma^* + P \to \gamma^* + P$ (virtual photon + proton goes to virtual photon + proton) shown in fig. 12 with $\mathrm{T}_{\mu\nu}$ its amplitude. It is a double virtual Compton scattering in the sense that both the incoming photon q' and the outgoing photon q have, in general, non-vanishing invariant masses. There is a non-zero four-momentum transfer from the virtual photon to the target proton which we label as r. Thus the struck proton p' has a slightly different four-momentum than the original proton p. And we have

$$p' = p - r \ ; \quad q' = q - r \ . \tag{2.1}$$

Furthermore, the incoming virtual photon q' has a different invariant mass than the outgoing virtual photon q with

$$q'^2 = -Q_1^2, \ \ q^2 = -Q_2^2 \, . \tag{2.2}$$

We restrict our discussion to amplitudes. Cross sections are obtained from the square of the amplitudes. In this non-forward case there is not a simple optical theorem relating the imaginary part of an amplitude to a cross section [12]. Our main interest is in the high energy limit, so similar to [17] we take all but one of the light cone (LC) components of the proton momentum to be zero. We do the same for the momentum transfer r. We will choose the non-zero component of both as the plus component, which means

$$p = (p_+, 0, \underline{0}) \, , \qquad r = (r_+, 0, \underline{0}) \, , \tag{2.3}$$

and

$$p^2 = 0 = r^2 \equiv -t \, . \tag{2.4}$$

That is, we are in the zero nucleon mass and zero t limit.

The fact that the target remains a proton means that

$$p'^2 = 0 = (p - r)^2 = p^2 - 2p \cdot r + r^2 \, , \tag{2.5}$$

and thus

$$p \cdot r = 0\,, \tag{2.6}$$

which gives

$$p \cdot q' = p \cdot q\,. \tag{2.7}$$

And since

$$q'^2 = (q - r)^2 = q^2 - 2q \cdot r + r^2 = q^2 - 2q \cdot r\,, \tag{2.8}$$

which results in

$$2q \cdot r = q^2 - q'^2\,. \tag{2.9}$$

Together with the fact that r is proportional to the external momentum p, we have,

$$r = \zeta p\,, \qquad \zeta = \frac{q^2 - q'^2}{2p \cdot q}\,. \tag{2.10}$$

We always suppose $-q^2 = Q_2^2 \leq Q_1^2 = q'^2$ so that $\zeta \geq 0$.

It is worth noting that the process of a double virtual Compton scattering is not a physical one, however, it provides a general framework in which one can move the discussion continuously from one physical limit to another. In particular, DIS corresponds to $q = q'$ while DVCS corresponds to $q = 0$.

Also worth noting is that the kinematic limit we are discussing, namely, $r \neq 0$ but $t = 0$, is not a physical one, since for a physical non-forward process, t is bounded by

$$|t| \geq \frac{x^2 m^2}{1 - x}\,, \tag{2.11}$$

where x is the Bjorken x variable and m the target proton mass. However, our main interest lies in the high energy and small-x behavior of the amplitude and therefore

$t = 0$ is a good approximation. In principle we can always discuss finite t behavior by giving r a transverse component $\underline{r} \neq 0$. There the proportionality between r and p (see 2.10) becomes a proportionality between the plus components of the two momenta (i.e. $r_+ = \zeta p_+$)(see, for example, [23]).

2.2 The Amplitude

The amplitude of the process in fig. 12 in coordinate space is a matrix element between two proton states of a time ordered product of two electromagnetic currents. If we label the space time point of the incoming interaction as $x + z$ and the outgoing one as x, we have

$$\mathrm{T}^{coord}_{\mu\nu} = \langle p'| T\, j_\mu(x) j_\nu(x+z) |p\rangle\,. \tag{2.12}$$

As usual we want to Fourier transform the amplitude into the momentum space where the kinematics and the dynamics of the process can be much more readily discussed. The momentum space amplitude, which we label as $\mathrm{T}_{\mu\nu}$, depends on all the relevant independent kinematic variables (in the momentum space). In this non-forward case there are three kinematic degrees of freedom and sets of variables such as (p, p', q) or (p, q, r) or (p, q', r) are valid choices and are all equivalent to each other. We choose the set (p, q', q) for convenience of later discussion. Therefore, using the convention that an incoming momentum p at space time point x enters the Fourier transformation as an exponential of negative exponent $e^{-ip\cdot x}$ and an outgoing momentum as one with a positive exponent, we can write explicitly the amplitude of fig. 12 in momentum space as

$$\mathrm{T}_{\mu\nu} = \mathrm{T}_{\mu\nu}(p, q', q)$$

$$= \; i\int d^4x d^4z\, e^{-iq'\cdot(x+z)+iq\cdot x}\,\langle p'|T\, j_\mu(x) j_\nu(x+z)|p\rangle\,. \tag{2.13}$$

Integration over d^4x gives only an overall four dimensional momentum space δ-function. Ignoring it we can rewrite the amplitude, up to normalization factors like $(2\pi)^4$, as

$$\mathrm{T}_{\mu\nu}(p,q',q) = i\int d^4z\, e^{-iq'\cdot z}\,\langle p'|T\, j_\mu(0) j_\nu(z)|p\rangle\,. \tag{2.14}$$

It is obvious that this amplitude should be invariant under an overall translation in coordinate space, for example, $(z \to 0\ \&\ 0 \to -z)$. Explicitly, by using the translation operator $e^{i\hat{p}\cdot x}$ where $\hat{p}$ is the momentum operator and for any operator in coordinate space $\hat{O}(z)$ (see equations 1.18 and 1.19)

$$e^{i\hat{p}\cdot x}\hat{O}(z-x)e^{-i\hat{p}\cdot x} = \hat{O}(z)\,, \tag{2.15}$$

we have

$$\begin{aligned}
\mathrm{T}_{\mu\nu}(p,q',q) &= i\int d^4z\, e^{-iq'\cdot z}\,\langle p'|T\, j_\mu(0) j_\nu(z)|p\rangle \\
&= i\int d^4z\, e^{-iq'\cdot z}\,\langle p'|T\, e^{i\hat{p}\cdot z} j_\mu(-z) e^{-i\hat{p}\cdot z} e^{i\hat{p}\cdot z} j_\nu(0) e^{-i\hat{p}\cdot z}|p\rangle \\
&= i\int d^4z\, e^{-iq'\cdot z} e^{ip'\cdot z}\langle p'|T\, j_\mu(-z) j_\nu(0)|p\rangle e^{-ip\cdot z} \\
&= i\int d^4z\, e^{-i(q'-p'+p)\cdot z}\langle p'|T\, j_\mu(-z) j_\nu(0)|p\rangle \\
&= i\int d^4z\, e^{-iq\cdot z}\langle p'|T\, j_\mu(-z) j_\nu(0)|p\rangle\,.
\end{aligned} \tag{2.16}$$

Anticipating the operator product expansion analysis we want to recast the amplitude into the so-called light cone expansion where $(z \to \frac{1}{2}z\ \&\ 0 \to -\frac{1}{2}z)$. Similar to the above process we have

$$\mathrm{T}_{\mu\nu}(p,q',q) \;=\; i\int d^4z\, e^{-iq'\cdot z}\,\langle p'|T\, j_\mu(0) j_\nu(z)|p\rangle$$

$$\begin{aligned}
&= i\int d^4z\, e^{-iq'\cdot z}\,\langle p'|T\, e^{i\hat{p}\cdot\frac{1}{2}z} j_\mu(-\frac{1}{2}z)e^{-i\hat{p}\cdot\frac{1}{2}z}e^{i\hat{p}\cdot\frac{1}{2}z} j_\nu(\frac{1}{2}z)e^{-i\hat{p}\cdot\frac{1}{2}z}|p\rangle \\
&= i\int d^4z\, e^{-iq'\cdot z}e^{ip'\cdot\frac{1}{2}z}\langle p'|T\, j_\mu(-\frac{1}{2}z)j_\nu(\frac{1}{2}z)|p\rangle e^{-ip\cdot\frac{1}{2}z} \\
&= i\int d^4z\, e^{-i(q'+\frac{1}{2}(p-p'))\cdot z}\langle p'|T\, j_\mu(-\frac{1}{2}z)j_\nu(\frac{1}{2}z)|p\rangle \\
&= i\int d^4z\, e^{-i(q'+\frac{1}{2}r)\cdot z}\langle p'|T\, j_\mu(-z)j_\nu(0)|p\rangle\,. \qquad (2.17)
\end{aligned}$$

Therefore, we arrive at

$$\mathrm{T}_{\mu\nu}(p,q',q) = i\int d^4z\, e^{-i\overline{q}\cdot z}\,\langle p'|T\, j_\mu(-\frac{z}{2})j_\nu(\frac{z}{2})|p\rangle \qquad (2.18)$$

where we have defined

$$\overline{q} = q' + \frac{1}{2}r = q - \frac{1}{2}r = \frac{1}{2}(q'+q)\,. \qquad (2.19)$$

Because from (2.9) we have

$$q\cdot q' = q^2 - q\cdot r = \frac{1}{2}(q'^2+q^2) = -\frac{1}{2}(Q_1^2+Q_2^2)\,, \qquad (2.20)$$

the square of this newly defined $\overline{q}$ becomes

$$\overline{q}^2 = \frac{1}{4}(q'+q)^2 = \frac{1}{2}(q'^2+q^2) = q\cdot q'\,. \qquad (2.21)$$

If we define

$$\begin{aligned}
\overline{Q}^2 &\equiv -\overline{q}^2 = -\frac{1}{2}(q'^2+q^2) \\
&= \frac{1}{2}(Q_1^2+Q_2^2)\,, \qquad (2.22)
\end{aligned}$$

we will see in later discussions that $\overline{Q}^2$ is now the *natural* scale of the scattering process and it characterizes the hardness of the scattering.

2.3 Current Conservation

Similar to the case shown in the previous chapter for the forward scattering we will use the conservation of the electro–magnetic(e&m) current (2.23) to simplify the tensorial decomposition of the non-forward amplitude $\mathrm{T}_{\mu\nu}$. However, because now the two e&m currents carries different momenta, we need to be more careful in writing down the consequences of the current conservation in momentum space.

Because

$$\frac{\partial}{\partial z^{\mu}} j_{\mu}(z) = 0 \tag{2.23}$$

from (2.16) we have

$$\begin{aligned} 0 &= i\int d^4z\, e^{-iq'\cdot z}\,\langle p'|T\, j_\mu(0)\frac{\partial}{\partial z^\nu} j_\nu(z)|p\rangle \\ &= i\int d^4z\, e^{-iq'\cdot z}\,\frac{\partial}{\partial z^\nu}\langle p'|T\, j_\mu(0) j_\nu(z)|p\rangle \\ &= i\int d^4z\, \frac{\partial}{\partial z^\nu}(e^{-iq'\cdot z}\,\langle p'|T\, j_\mu(0) j_\nu(z)|p\rangle) \\ &\quad -i\int d^4z\,(-iq')_\nu e^{-iq'\cdot z}\langle p'|T\, j_\mu(0) j_\nu(z)|p\rangle) \\ &= 0 + iq'_\nu \mathrm{T}_{\mu\nu} \end{aligned} \tag{2.24}$$

and

$$\begin{aligned} 0 &= i\int d^4z\, e^{-iq\cdot z}\,\langle p'|T\,(\frac{\partial}{\partial z^\mu} j_\mu(-z)) j_\nu(0)|p\rangle \\ &= i\int d^4z\, e^{-iq\cdot z}\,\frac{\partial}{\partial z^\mu}\langle p'|T\, j_\mu(-z) j_\nu(0)|p\rangle \\ &= i\int d^4z\, \frac{\partial}{\partial z^\mu}(e^{-iq\cdot z}\,\langle p'|T\, j_\mu(0) j_\nu(z)|p\rangle) \\ &\quad -i\int d^4z\,(-iq)_\nu e^{-iq\cdot z}\langle p'|T\, j_\mu(-z) j_\nu(0)|p\rangle) \\ &= 0 + iq_\mu \mathrm{T}_{\mu\nu}\,. \end{aligned} \tag{2.25}$$

Thus in momentum space the conservation of the electro–magnetic current now requires the amplitude to satisfy (cf 1.22)

$$q^{\mu}\mathrm{T}_{\mu\nu} = 0; \quad q'^{\nu}\mathrm{T}_{\mu\nu} = 0. \tag{2.26}$$

2.4 Decomposition and Invariant Amplitudes

To obtain the complete tensorial decomposition of $\mathrm{T}_{\mu\nu}$ we also need the symmetry properties of the amplitude. From fig. 13 we can see clearly that the second diagram is simply the first diagram with a different momentum labeling and thus the two have identical values, both equal to $\mathrm{T}_{\mu\nu}$. Note that to be completely explicit, each diagram in fig. 13 should have its own corresponding cross diagram, and the symmetry exists for the sum of the cross diagram and the original one.

Therefore, while $\mathrm{T}_{\mu\nu}$ is no longer symmetric, it is invariant under the transformation $\mu \leftrightarrow \nu, \;\; q \leftrightarrow -q'$, that is,

$$\mathrm{T}_{\mu\nu}(p, q', q) = \mathrm{T}_{\nu\mu}(p, -q, -q')\,. \tag{2.27}$$

From Section 2.2 we know that $\mathrm{T}_{\mu\nu}$ depends on the momenta p, q and q', similar to the last chapter we can write the most general tensorial decomposition of $\mathrm{T}_{\mu\nu}$ as

$$\begin{aligned}\mathrm{T}_{\mu\nu}(p, q', q) \;\; = \;\; & a_0 g_{\mu\nu} + a_1 p_\mu p_\nu + a_2 q'_\mu q'_\nu + a_3 q_\mu q_\nu + a_4 p_\mu q'_\nu + a_5 q'_\mu p_\nu \\ & + a_6 p_\mu q_\nu + a_7 q_\mu p_\nu + a_8 q'_\mu q_\nu + a_9 q_\mu q'_\nu\,. \end{aligned} \tag{2.28}$$

As in the previous chapter, this is a spin averaged electron-scattering, thus we do not need to consider any helicity issues and hence there are no ϵ-tensor terms in the decomposition.

The a_is are invariant amplitudes depending again only on scalar products of the momenta. Because we have equations (2.6), (2.7), (2.9), and (2.20), we take the a_is to only depend on q'^2, q^2 and $p \cdot q$, that is

$$a_i = a_i(q'^2, q^2, p \cdot q), \qquad i = 0, 1, 2, ..., 9\,. \tag{2.29}$$

From (2.27) and (2.28) we have

$$\begin{aligned} \mathrm{T}_{\nu\mu}(p, -q, -q') &= a_0(-)g_{\mu\nu} + a_1(-)p_\mu p_\nu + a_2(-)q'_\mu q'_\nu + a_3(-)q_\mu q_\nu - a_4(-)p_\mu q'_\nu \\ &\quad -a_5(-)q'_\mu p_\nu - a_6(-)p_\mu q_\nu - a_7(-)q_\mu p_\nu + a_8(-)q'_\mu q_\nu + a_9(-)q_\mu q'_\nu \\ &= \mathrm{T}_{\mu\nu}(p, q', q) \end{aligned} \tag{2.30}$$

where we have used the short handed notation

$$a_i(-) = a_i(q \leftrightarrow -q') = a_i(q'^2, q^2, -p \cdot q)\,. \tag{2.31}$$

Equation 2.30 is true for any values of the momenta q', q and p, in particular, we can fix the scalar products but still vary arbitrarily the individual momentum. Thus the coefficients of the same momentum combinations must be identical, which leads to the following relationships among the invariant amplitudes:

$$\begin{gathered} a_0 = a_0(-)\,; a_1 = a_1(-)\,; a_8 = a_8(-)\,; a_9 = a_9(-)\,; \\ a_2 = a_3(-)\,; a_3 = a_2(-)\,; a_4 = -a_7(-)\,; a_5 = -a_6(-)\,. \end{gathered} \tag{2.32}$$

By using the conservation of the e&m current (2.26) we can establish another two sets of equations of the invariants. We have

$$q^\mu \mathrm{T}_{\mu\nu} = 0 \;\; = \;\; a_0 q_\nu + a_1 p \cdot q p_\nu + a_2 q \cdot q' q'_\nu + a_3 q^2 q_\nu + a_4 p \cdot q q'_\nu + a_5 p \cdot q' p_\nu$$

$$
\begin{aligned}
& && +a_6 p\cdot q q_\nu + a_7 q^2 p_\nu + a_8 q\cdot q' q_\nu + a_9 q^2 q'_\nu\,; \\
q'^\nu \mathrm{T}_{\mu\nu} = 0 &= && a_0 q'_\mu + a_1 p\cdot q' p_\mu + a_2 q'^2 q'_\mu + a_3 q\cdot q' q_\mu + a_4 q'^2 p_\mu + a_5 p\cdot q' q'_\mu \\
& && +a_6 q\cdot q' p_\mu + a_7 p\cdot q' q_\mu + a_8 q\cdot q' q'_\mu + a_9 q'^2 q_\mu\,. \qquad (2.33)
\end{aligned}
$$

Because again this is always true when we vary arbitrarily the momenta while keeping the scalar products fixed, the coefficients of the same combinations of momenta must identically vanish, leading to

$$
\begin{aligned}
a_0 + a_3 q^2 + a_6\, p\cdot q + a_8\, q\cdot q' &= 0 \\
a_7 q^2 + a_1\, p\cdot q + a_5\, q\cdot q' &= 0 \\
a_9 q^2 + a_4\, p\cdot q + a_2\, q\cdot q' &= 0 \\
a_0 + a_2 q'^2 + a_5\, p\cdot q' + a_8\, q\cdot q' &= 0 \\
a_4 q'^2 + a_1\, p\cdot q' + a_6\, q\cdot q' &= 0 \\
a_9 q'^2 + a_7\, p\cdot q' + a_3\, q\cdot q' &= 0\,. \qquad (2.34)
\end{aligned}
$$

As in the previous chapter, we will use these equations to reduce the number of independent invariant amplitudes. We first rewrite (2.34) by taking $a_{0,1,2,3,5}$ as given and expressing the rest of $a_i's$ in terms of them. We obtain

$$
\begin{aligned}
a_4 &= -\frac{q\cdot q'}{p\cdot q} a_2 - \frac{q^2}{p\cdot q} a_9 \\
&= -\frac{p\cdot q'}{q'^2} a_1 - \frac{q\cdot q'}{p\cdot q} a_2 + \frac{q\cdot q' q^2}{q'^2 p\cdot q} a_3 - \frac{q\cdot q'}{q'^2} a_5\,, \\
a_6 &= -\frac{p\cdot q}{q\cdot q'} a_1 - \frac{q'^2}{q\cdot q'} a_4 \\
&= +\frac{q'^2}{p\cdot q} a_2 - \frac{q^2}{p\cdot q} a_3 + a_5\,, \\
a_7 &= -\frac{p\cdot q}{q^2} a_1 - \frac{q\cdot q'}{q^2} a_5\,,
\end{aligned}
$$

$$\begin{aligned} a_8 &= -\frac{1}{q\cdot q'}a_0 - \frac{q'^2}{q\cdot q'}a_2 - \frac{p\cdot q'}{q\cdot q'}a_5\,, \\ a_9 &= -\frac{q\cdot q'}{q'^2}a_3 - \frac{p\cdot q'}{q'^2}a_7 \\ &= \frac{p\cdot q p\cdot q'}{q^2 q'^2}a_1 - \frac{q\cdot q'}{q'^2}a_3 + \frac{p\cdot q' q\cdot q'}{q^2 q'^2}a_5\,. \end{aligned} \tag{2.35}$$

We now use equation 2.32 to further simplify the above result. We have

$$\begin{aligned} a_8 &= a_8(-) \;\Rightarrow \\ & -\frac{1}{q\cdot q'}a_0 - \frac{q'^2}{q\cdot q'}a_2 - \frac{p\cdot q'}{q\cdot q'}a_5 = -\frac{1}{q\cdot q'}a_0(-) - \frac{q'^2}{q\cdot q'}a_2(-) + \frac{p\cdot q'}{q\cdot q'}a_5(-) \\ \Rightarrow\;& a_5 + a_5(-) = \frac{p\cdot q}{q'^2}(a_3 - a_2)\,, \\ a_9 &= a_9(-) \;\Rightarrow \\ & \frac{p\cdot q p\cdot q'}{q^2 q'^2}a_1 - \frac{q\cdot q'}{q'^2}a_3 + \frac{p\cdot q' q\cdot q'}{q^2 q'^2}a_5 \\ & \qquad = \frac{p\cdot q p\cdot q'}{q^2 q'^2}a_1(-) - \frac{q\cdot q'}{q'^2}a_3(-) - \frac{p\cdot q' q\cdot q'}{q^2 q'^2}a_5(-) \\ \Rightarrow\;& a_5 + a_5(-) = \frac{p\cdot q}{q^2}(a_3 - a_2)\,, \end{aligned} \tag{2.36}$$

where we have used $a_1 = a_1(-)$, $a_3(-) = a_2$ and $p\cdot q = p\cdot q'$. From the above, it is obvious that we must have

$$a_2 = a_3\,, \quad a_5 = -a_5(-)\,. \tag{2.37}$$

Furthermore,

$$\begin{aligned} a_5 &= -a_6(-) \;\Rightarrow \\ & a_5 = -(-\frac{q'^2}{p\cdot q}a_2(-) + \frac{q^2}{p\cdot q}a_3(-) + a_5(-)) \\ \Rightarrow\;& a_5 + a_5(-) = \frac{q'^2}{p\cdot q}a_3 - \frac{q^2}{p\cdot q}a_2\,, \end{aligned} \tag{2.38}$$

which, combined with (2.37), gives

$$a_2 = a_3 = 0\,. \tag{2.39}$$

At the same time,

$$\begin{aligned}
a_4 &= -a_7(-) \;\; \Rightarrow \\
& -\frac{p\cdot q'}{q'^2}a_1 - \frac{q\cdot q'}{p\cdot q}a_2 + \frac{q\cdot q' q^2}{q'^2 p\cdot q}a_3 - \frac{q\cdot q'}{q'^2}a_5 = -(\frac{p\cdot q}{q^2}a_1(-) - \frac{q\cdot q'}{q^2}a_5(-)) \\
\Rightarrow & \;\; a_5 = -\frac{p\cdot q}{q\cdot q'}a_1 \,. \qquad (2.40)
\end{aligned}$$

Therefore, going back to (2.35) we have

$$\begin{aligned}
a_4 &= -\frac{p\cdot q'}{q'^2}a_1 - \frac{q\cdot q'}{q'^2}(-\frac{p\cdot q}{q\cdot q'})a_1 \\
&= 0\,, \\
a_6 &= a_5 = -\frac{p\cdot q}{q\cdot q'}a_1\,, \\
a_7 &= -\frac{p\cdot q}{q^2}a_1 - \frac{q\cdot q'}{q^2}(-\frac{p\cdot q}{q\cdot q'})a_1 \\
&= 0\,, \\
a_8 &= -\frac{1}{q\cdot q'}a_0 - \frac{p\cdot q'}{q\cdot q'}(-\frac{p\cdot q}{q\cdot q'})a_1 \\
&= -\frac{1}{q\cdot q'}a_0 + \frac{(p\cdot q)^2}{(q\cdot q')^2}a_1\,, \\
a_9 &= \frac{p\cdot q p\cdot q'}{q^2 q'^2}a_1 + \frac{p\cdot q' q\cdot q'}{q^2 q'^2}(-\frac{p\cdot q}{q\cdot q'})a_1 \\
&= 0\,. \qquad (2.41)
\end{aligned}$$

It is now clear that only two independent invariant amplitudes, a_0 and a_1, remain in the end while all the other ones either vanish identically or can be expressed in terms of a_0 and a_1. In summary, we have

$$\begin{aligned}
a_2 = a_3 = a_4 = a_7 = a_9 = 0 \\
a_5 = a_6 = -\frac{p\cdot q}{q\cdot q'}a_1 \\
a_8 = -\frac{1}{q\cdot q'}a_0 + \frac{(p\cdot q)^2}{(q\cdot q')^2}a_1\,. \qquad (2.42)
\end{aligned}$$

Thus, substitute these back into (2.28) we have

$$\begin{aligned}
\mathrm{T}_{\mu\nu}(p,q',q) &= a_0 g_{\mu\nu} + a_1 p_\mu p_\nu - \frac{p\cdot q}{q\cdot q'} a_1 (q'_\mu p_\nu + p_\mu q_\nu) \\
&\quad +(-\frac{1}{q\cdot q'} a_0 + \frac{(p\cdot q)^2}{(q\cdot q')^2} a_1)\, q'_\mu q_\nu \\
&= \left(g_{\mu\nu} - \frac{q'_\mu q_\nu}{q\cdot q'}\right) a_0 \\
&\quad + \left(p_\mu p_\nu - \frac{p\cdot q}{q\cdot q'}(q'_\mu p_\nu + p_\mu q_\nu) + \frac{(p\cdot q)^2}{(q\cdot q')^2} q'_\mu q_\nu\right) a_1 \,. \qquad (2.43)
\end{aligned}$$

By defining two invariant amplitudes T_1 and T_2 we finally obtain the general tensorial decomposition of $\mathrm{T}_{\mu\nu}$ as

$$\begin{aligned}
\frac{1}{\sqrt{1-\zeta}} \mathrm{T}_{\mu\nu} &= \left(-g_{\mu\nu} + \frac{q'_\mu q_\nu}{\overline{q}^2}\right) \mathrm{T}_1 \\
&\quad + \frac{1}{M^2}\left(p_\mu p_\nu - \frac{p\cdot q}{\overline{q}^2}(p_\mu q_\nu + q'_\mu p_\nu) + (\frac{p\cdot q}{\overline{q}^2})^2 q'_\mu q_\nu\right) \mathrm{T}_2 \,. \qquad (2.44)
\end{aligned}$$

$\mathrm{T_i} = \mathrm{T_i}(q'^2, q^2, p\cdot q)$ are invariant amplitudes analogous to those in DIS (see section 1.3.3). They are even functions of $p\cdot q$ (because they are essentially a_0 and a_1).

We have pulled out from $\mathrm{T_i}$ an explicit factor of $\sqrt{1-\zeta}$, as was done in, e.g. ,[16] (see equation 2.10 for the definition of ζ). This factor comes from the external spinors of the proton as the following: The normalization convention we use for computing Feynman diagrams is such that an external incoming Fermion with momentum p and spin state r enters a Feynman diagram expression as $\frac{U_r(p)}{\sqrt{p}}$ where $U_r(p)$ is the standard 4-component basis spinor for a Fermion. Thus there is an explicit factor of $\sqrt{p'}^{-1} \equiv (\sqrt{1-\zeta}\sqrt{p})^{-1}$ in the expression of $\mathrm{T}_{\mu\nu}$. We pull it outside explicitly so that later on the expression of the lowest order Wilson coefficient will be simpler (see equation 4.60 in section 4.3).

It is worth noting that we still have only two independent invariant amplitudes even in the presence of three independent invariants, in contrast to the case of forward scattering where we only have two (see sections 1.2.1 and 1.3.3).

Chapter 3

Operator Product Expansion

In this chapter we will discuss the operator product expansion (OPE) of the non-forward amplitude $\mathrm{T}_{\mu\nu}$ (see section 2.2). We will identify the operators in the OPE, obtain their reduced matrix elements between (asymmetric) external states and define new moment variables. We will express $\mathrm{T}_{\mu\nu}$ and eventually the invariant amplitudes T_1 and T_2 (see equation 2.44) in terms of the reduced matrix elements and the corresponding Wilson coefficients.

3.1 Non-forward Operator Product Expansion

In the short distance limit, we can perform an OPE for $\mathrm{T}_{\mu\nu}$ as a sum of products of local operators and their corresponding Wilson coefficients [5]:

$$T\, j_\mu(-\frac{z}{2}) j_\nu(\frac{z}{2}) \xrightarrow{z_\mu \to 0} \hat{A}_{\mu\nu} \sum_{J=1}^{\infty} \sum_{n=0}^{J} \sum_{i=1}^{u_J} F^{(i)}_{J,n}(z^2) \hat{O}^{(i)(J,n)}_{\mu_1 \ldots \mu_J}(0) \frac{z^{\mu_1}}{2} \cdots \frac{z^{\mu_J}}{2} + \hat{B}_{\mu\nu\alpha\beta} \sum_{J=1}^{\infty} \sum_{n=0}^{J+2} \sum_{i=1}^{u_J} E^{(i)}_{J,n}(z^2) \hat{O}^{(i)(J+2,n)}_{\alpha\beta;\mu_1 \ldots \mu_J}(0) \frac{z^{\mu_1}}{2} \cdots \frac{z^{\mu_J}}{2}\,, \qquad (3.1)$$

where $\hat{A}_{\mu\nu}$ and $\hat{B}_{\mu\nu\alpha\beta}$ are conserved tensor structure operators. The summations should be regarded as such that in addition to the literal summation over J, n and i, all the indices except μ, ν are also symmetrized as usual.

To obtain the explicit expression of $\hat{B}_{\mu\nu\alpha\beta}$ we start with the most general form that an tensor operator in the position of $\hat{B}$ can have:

$$\begin{aligned}
\hat{O}_{\mu\nu\alpha\beta} &= a_1 g_{\mu\nu} g_{\alpha\beta} \Box + a_2 g_{\mu\alpha} g_{\nu\beta} \Box + a_3 g_{\mu\beta} g_{\nu\alpha} \Box \\
&\quad + a_4 g_{\mu\nu} \partial_\alpha \partial_\beta + a_5 g_{\mu\alpha} \partial_\nu \partial_\beta + a_6 g_{\mu\beta} \partial_\alpha \partial_\nu \\
&\quad + a_7 g_{\nu\beta} \partial_\mu \partial_\alpha + a_8 g_{\nu\alpha} \partial_\mu \partial_\beta + a_9 g_{\alpha\beta} \partial_\mu \partial_\nu \,. \qquad (3.2)
\end{aligned}$$

Current conservation requires

$$\partial_\mu \hat{O}_{\mu\nu\alpha\beta} = \partial_\nu \hat{O}_{\mu\nu\alpha\beta} = 0 \,. \qquad (3.3)$$

We have

$$\begin{aligned}
0 &= a_1 \partial_\nu g_{\alpha\beta} \Box + a_2 \partial_\alpha g_{\nu\beta} \Box + a_3 \partial_\beta g_{\nu\alpha} \Box + a_4 \partial_\nu \partial_\alpha \partial_\beta \\
&\quad + a_5 \partial_\alpha \partial_\nu \partial_\beta + a_6 \partial_\beta \partial_\alpha \partial_\nu + a_7 g_{\nu\beta} \partial_\alpha \Box + a_8 g_{\nu\alpha} \Box \partial_\beta + a_9 g_{\alpha\beta} \Box \partial_\nu \\
&= ((a_1 + a_9) g_{\alpha\beta} \partial_\nu + (a_2 + a_7) g_{\nu\beta} \partial_\alpha + (a_3 + a_8) g_{\nu\alpha} \partial_\beta) \Box \\
&\quad + (a_4 + a_5 + a_6) \partial_\nu \partial_\alpha \partial_\beta \,, \qquad (3.4)
\end{aligned}$$

and

$$\begin{aligned}
0 &= a_1 \partial_\mu g_{\alpha\beta} \Box + a_2 g_{\mu\alpha} \partial_\beta \Box + a_3 g_{\mu\beta} \partial_\alpha \Box + a_4 \partial_\mu \partial_\alpha \partial_\beta \\
&\quad + a_5 g_{\mu\alpha} \partial_\beta \Box + a_6 g_{\mu\beta} \partial_\alpha \Box + a_7 \partial_\beta \partial_\mu \partial_\alpha + a_8 \partial_\alpha \partial_\mu \partial_\beta + a_9 g_{\alpha\beta} \Box \partial_\mu \\
&= ((a_1 + a_9) g_{\alpha\beta} \partial_\mu + (a_2 + a_5) g_{\mu\alpha} \partial_\beta + (a_3 + a_6) g_{\mu\beta} \partial_\alpha) \Box \\
&\quad + (a_4 + a_7 + a_8) \partial_\mu \partial_\alpha \partial_\beta \,. \qquad (3.5)
\end{aligned}$$

Since the above two equations are tensor equations, the coefficient of each (different) tensor structure operator must vanish, which gives us

$$a_1 + a_9 = a_2 + a_7 = a_2 + a_5 = a_3 + a_8 = a_3 + a_6 = 0$$
$$a_4 + a_5 + a_6 = a_4 + a_7 + a_8 = 0 \,. \tag{3.6}$$

Thus we have

$$\begin{aligned} a_9 &= -a_1 \\ a_4 &= a_2 + a_3 \\ a_5 &= a_7 = -a_2 \\ a_6 &= a_8 = -a_3 \,, \end{aligned} \tag{3.7}$$

which, after substitution back into (3.2), leads us to

$$\begin{aligned} \hat{O}_{\mu\nu\alpha\beta} &= a_1 g_{\mu\nu} g_{\alpha\beta} \Box + a_2 g_{\mu\alpha} g_{\nu\beta} \Box + a_3 g_{\mu\beta} g_{\nu\alpha} \Box \\ &\quad + (a_2 + a_3) g_{\mu\nu} \partial_\alpha \partial_\beta - a_2 g_{\mu\alpha} \partial_\nu \partial_\beta - a_3 g_{\mu\beta} \partial_\alpha \partial_\nu \\ &\quad - a_2 g_{\nu\beta} \partial_\mu \partial_\alpha - a_3 g_{\nu\alpha} \partial_\mu \partial_\beta - a_1 g_{\alpha\beta} \partial_\mu \partial_\nu \\ &= (g_{\mu\nu} g_{\alpha\beta} \Box - g_{\alpha\beta} \partial_\mu \partial_\nu) a_1 + (g_{\mu\alpha} g_{\nu\beta} \Box + g_{\mu\nu} \partial_\alpha \partial_\beta \\ &\quad - g_{\mu\alpha} \partial_\nu \partial_\beta - g_{\nu\beta} \partial_\mu \partial_\alpha) a_2 + (g_{\mu\beta} g_{\nu\alpha} \Box \\ &\quad + g_{\mu\nu} \partial_\alpha \partial_\beta - g_{\mu\beta} \partial_\alpha \partial_\nu - g_{\nu\alpha} \partial_\mu \partial_\beta) a_3 \,. \end{aligned} \tag{3.8}$$

If we symmetrize the indices α and β we have

$$\begin{aligned} \hat{O}_{\mu\nu\alpha\beta} &= g_{\alpha\beta} (g_{\mu\nu} \Box - \partial_\mu \partial_\nu) a_1 \\ &\quad + (g_{\mu\alpha} g_{\nu\beta} \Box + g_{\mu\nu} \partial_\alpha \partial_\beta - g_{\mu\alpha} \partial_\nu \partial_\beta - g_{\nu\beta} \partial_\mu \partial_\alpha) a_2 \,. \end{aligned} \tag{3.9}$$

It is now obvious that $\hat{B}_{\mu\nu\alpha\beta}$ is the same as in the forward case (see equation 1.54), i.e. ,

$$\hat{B}_{\mu\nu\alpha\beta} = g_{\mu\alpha} g_{\nu\beta} \Box + g_{\mu\nu} \partial_\alpha \partial_\beta - g_{\mu\alpha} \partial_\nu \partial_\beta - g_{\nu\beta} \partial_\mu \partial_\alpha \,, \tag{3.10}$$

and corresponds to the tensor structure multiplying T_2 in (2.44).

We have not found the explicit form of $\hat{A}_{\mu\nu}$ which would generate the tensor structure corresponding to T_1. But, as in the forward case, once we know T_2, we can obtain T_1 by a Callen-Gross relationship, at least in leading logarithmic level (see equation 3.38). In the following discussion, we will leave $\hat{A}_{\mu\nu}$ as an unspecified general conserved tensor operator. Eventually we will obtain the necessary terms from the Callen-Gross relationship.

3.2 The Operators

The operators that enter into the OPE of an amplitude are composed of the fields involved in the interaction and derivatives. In the most general case, the derivatives can either be internal, bi-directional derivatives like the ones in equation 1.92, or overall derivatives that act outside the fields. These overall derivatives, when evaluated between external states, give the difference in momentum (in momentum space) between the incoming and outgoing states, or, equivalently, the net momentum inflow/outflow of the local vertex of the interaction. Thus operators with overall derivatives are identically zero in a forward scattering process, and we do not need to consider them in forward OPE (see section 1.3.3). However, in a non-forward case as the one we are discussing, because the momentum flowing into the local vertex is r instead of zero, we have to include in the expansion new sets of operators that have overall derivatives.

In leading twist (the physical consequences of leading twist in this case will be clear later) the operators for QCD are (cf. equation 1.92)

$$\begin{aligned}
\hat{O}^{(q)(J,n)}_{\mu_1 ... \mu_J} &= \partial_{\mu_1} ... \partial_{\mu_n} \, \tilde{q}(0) \gamma_{\mu_{n+1}} i \overleftrightarrow{D}_{\mu_{n+2}} ... i \overleftrightarrow{D}_{\mu_J} \, q(0) && (3.11) \\
\hat{O}^{(g)(J,n)}_{\mu_1 ... \mu_J} &= \partial_{\mu_1} ... \partial_{\mu_n} \, F_{\mu_{n+1}}{}^{\nu}(0) i \overleftrightarrow{\mathcal{D}}_{\mu_{n+1}} ... i \overleftrightarrow{\mathcal{D}}_{\mu_J} \, F_{\mu_J \nu}(0) \, , && (3.12)
\end{aligned}$$

where it is again understood that the indices are all symmetrized.

After taking the matrix elements between the asymmetric external states, and after taking a Fourier transform, the external derivatives would eventually be turned into factors of r_μ while the internal derivatives into either r_μ or $(p+p')_\mu \equiv (2p-r)_\mu$. We thus define the two moment variables in the non-forward case as

$$\omega = \frac{(2p-r) \cdot q}{\overline{Q}^2}, \qquad \nu = \frac{r \cdot q}{\overline{Q}^2} \; . \tag{3.13}$$

The forward case would be the limit $\nu = 0$ while DVCS corresponds to $\nu = 1$.

In a QCD-Parton picture, the diagrams contributing to $\mathrm{T}_{\mu\nu}$ are the so-called *hand-bag* diagrams as shown in figure 14. If we parametrize the momentum of the scattered parton as $k = xp + yr$ (see, e.g. [16]), where x and y are two Bjorken type scaling variables defined by

$$\begin{aligned}
x &= \frac{Q_1^2}{2p \cdot q} \equiv \frac{1}{\tilde{\omega}} \\
y &= \frac{Q_1^2}{2r \cdot q} \equiv \frac{1}{\tilde{\nu}} \, , && (3.14)
\end{aligned}$$

our moment variables are related to these quantities via

$$\begin{aligned}
\omega &= \frac{Q_1^2}{\overline{Q}^2} \, (x\tilde{\omega} + y\tilde{\nu} - \frac{1}{2}\tilde{\nu}) \\
\nu &= \frac{Q_1^2}{\overline{Q}^2} \, (\frac{1}{2}\tilde{\nu}) \, . && (3.15)
\end{aligned}$$

There will be mixing among operators with the same J but different n labels. Under evolution in the momentum scale, which as we shall see later, is characterized by $\overline{Q}^2$, the internal derivatives, with which the dynamics of the process lies, can be turned into either overall derivatives or again internal ones, and, as stated before, eventually give rise to factors of either $2p - r$ or r. On the other hand, the overall derivatives, which are in essence only involved with kinematics, can be turned into themselves only and eventually give factors of r. Therefore, evolution in $\overline{Q}^2$ will lead to mixing of these operators in only one direction in an upper-triangular fashion, namely,

$$\hat{O}^{(i)(J,n)} \rightarrow \hat{O}^{(i)(J,n')}\,, \qquad 0 \leq n \leq n' \leq J\,. \tag{3.16}$$

However, we do still have the freedom to choose, for simplification, at a factorization scale μ_0, that all internal derivatives give factors of $(2p - r)$. Thus we can write

$$\langle p'|\hat{O}^{(i)(J,n)}_{\mu_1...\mu_J}|p\rangle_{(\mu_0)} = r_{\mu_1}...r_{\mu_n}(2p-r)_{\mu_{n+1}}...(2p-r)_{\mu_J}\,\langle p'||\hat{O}^{(i)(J,n)}||p\rangle_{(\mu_0)}\,. \tag{3.17}$$

This is also the definition of the reduced matrix elements of $\hat{O}^{(i)(J,n)}$. It is clear that from the above discussion, these reduced matrix elements will depend only on $J - n$, since the overall derivatives give (in certain sense) trivial kinematic factors of r, and we have

$$\langle p'||\hat{O}^{(i)(J,n)}||p\rangle = \langle p'||\hat{O}^{(i)(J-n,0)}||p\rangle\ . \tag{3.18}$$

The choice of (3.17) is essentially the same as taking the leading twist approximation. Analogous to the discussion in the forward case (1.3.3), the indices of the asymmetric matrix elements of the operators in the OPE must be made from $g_{\mu_i\mu_j}$ or $p_{\mu_i}(p'_{\mu_i})$. After Fourier transformation the factors of z_{μ_i} are turned into momentum factors, namely,

$$z_{\mu_i} \Rightarrow \frac{\overline{q}_{\mu_i}}{\overline{q}^2}\,. \tag{3.19}$$

And we have

$$g_{\mu_i\mu_j}\frac{\overline{q}_{\mu_i}}{\overline{q}^2}\frac{\overline{q}_{\mu_j}}{\overline{q}^2} \sim \frac{1}{\overline{q}^2}\,, \tag{3.20}$$

while (see 3.30)

$$p_{\mu_i}p_{\mu_j}\frac{\overline{q}_{\mu_i}}{\overline{q}^2}\frac{\overline{q}_{\mu_j}}{\overline{q}^2} = (\frac{p\cdot q}{\overline{q}^2})^2 \sim \mathrm{O}(1)\,. \tag{3.21}$$

Therefore the leading contribution from the operators in the OPE should all give factors of the external momentum, while $g_{\mu_i\mu_j}$ type of contributions are small (by factors of $\frac{1}{\overline{q}^2}$). It is clear that this also makes the leading twist operators dominate (cf. [5]). Non-leading twist terms in this case are suppressed by at least a power of $\frac{1}{\overline{Q}^2}$. And $\overline{q}$ is now the measure of the momentum scale of the scattering.

3.3 Wilson Coefficients and Explicit OPE

To obtain the explicit expression of the OPE of the amplitude, we rewrite equation 3.1 by explicit substitution of the tensor structure operators as

$$\begin{aligned} T\, j_\mu(-\frac{z}{2})j_\nu(\frac{z}{2}) \quad &\xrightarrow{z_\mu\to 0} \quad \hat{A}_{\mu\nu}\sum_{J=1}^{\infty}\sum_{n=0}^{J}\sum_{i=1}^{u_J} F^{(i)}_{J,n}(z^2)\hat{O}^{(i)(J,n)}_{\mu_1\ldots\mu_J}(0)\frac{z^{\mu_1}}{2}\cdots\frac{z^{\mu_J}}{2} \\ &+ \quad (g_{\mu\alpha}g_{\nu\beta}\Box + g_{\mu\nu}\partial_\alpha\partial_\beta - g_{\mu\alpha}\partial_\nu\partial_\beta - g_{\nu\beta}\partial_\mu\partial_\alpha) \\ &\qquad \sum_{J=1}^{\infty}\sum_{n=0}^{J+2}\sum_{i=1}^{u_J} E^{(i)}_{J,n}(z^2)\hat{O}^{(i)(J+2,n)}_{\alpha\beta;\mu_1\ldots\mu_J}(0)\frac{z^{\mu_1}}{2}\cdots\frac{z^{\mu_J}}{2}\,. \end{aligned} \tag{3.22}$$

We evaluate the above time-ordered product between the asymmetric external proton states and expand it at an arbitrarily chosen factorization scale μ_0 into products of reduced matrix elements, corresponding Wilson coefficients, and the kinematic factors coming from the tensor structure operators. We obtain

$$\begin{aligned}
&\langle p'|T\, j_\mu(-\frac{z}{2}) j_\nu(\frac{z}{2})|p\rangle \longrightarrow \\
&\sum_{J=1}^{\infty}\sum_{n=0}^{J}\sum_{i=1}^{u_J} \langle p'||\hat{O}^{(i)(J,n)}||p\rangle_{(\mu_0)} \hat{A}_{\mu\nu} F^{(i)}_{J,n}(z^2)(2)^{-J}((2p{-}r)\cdot z)^{J-n}(r\cdot z)^n \\
&\quad + \sum_{J=1}^{\infty}\sum_{n=0}^{J+2}\sum_{i=1}^{u_J} \langle p'||\hat{O}^{(i)(J+2,n)}||p\rangle_{(\mu_0)} \\
&\quad \cdot \hat{B}_{\mu\nu} E^{(i)}_{J,n}(z^2)(2)^{-J}((2p{-}r)\cdot z)^{J-n}(r\cdot z)^{n-2}\mathcal{C}_{\alpha\beta}(2p{-}r, r)\,.
\end{aligned} \tag{3.23}$$

Because the α and β indices can either come from $2p{-}r$ or r when going to the reduced matrix elements we have included a factor of $\mathcal{C}_{\alpha\beta}$ that incorporates all the possible combinations in a symmetric fashion. Explicitly, it is the completely symmetrized sum of the following three terms:

$$\begin{aligned}
&(2p-r)_\alpha(2p-r)_\beta r_{\mu_i} r_{\mu_j}\ , \\
&(2p-r)_\alpha r_\beta (2p-r)_{\mu_i} r_{\mu_j}\ , \\
&r_\alpha r_\beta (2p-r)_{\mu_i}(2p-r)_{\mu_j}\ ,
\end{aligned} \tag{3.24}$$

together with a normalization factor that we need to put in to compensate for the double counting of terms because, as stated before, the summation signs have already implied a complete symmetrization of all indices except μ and ν.

The first term of equation (3.23) becomes

$$\sum_{J=1}^{\infty}\sum_{n=0}^{J}\sum_{i=1}^{u_J} \langle p'||\hat{O}^{(i)(J,n)}||p\rangle_{(\mu_0)}(2)^{-J}(2-\zeta)^{J-n}\zeta^n \hat{A}_{\mu\nu}\, F^{(i)}_{J,n}(z^2)(p\cdot z)^J\,, \tag{3.25}$$

where we have used equation (2.10). The second term, after a shift of labeling $J{+}2 \rightarrow J$, becomes

$$\begin{aligned} &\sum_{J=1}^{\infty}\sum_{n=0}^{J}\sum_{i=1}^{u_J}\langle p'||\hat{O}^{(i)(J,n)}||p\rangle_{(\mu_0)}\,(g_{\mu\alpha}g_{\nu\beta}\Box+g_{\mu\nu}\partial_\alpha\partial_\beta-g_{\mu\alpha}\partial_\nu\partial_\beta-g_{\nu\beta}\partial_\mu\partial_\alpha)\\ &\qquad\qquad\cdot E^{(i)}_{J-2,n}(z^2)2^{-(J-2)}(2-\zeta)^{J-n-2}\zeta^{n-2}(p\cdot z)^{J-4}\mathcal{C}_{\alpha\beta}(2p-r,r)\\ =\;&\sum_{J=1}^{\infty}\sum_{n=0}^{J}\sum_{i=1}^{u_J}\langle p'||\hat{O}^{(i)(J,n)}||p\rangle_{(\mu_0)}\;2^{-(J-2)}\;(2-\zeta)^{J-n-2}\,\zeta^{n-2}\;\mathcal{C}\\ &\quad\cdot(((2p-r)_\mu(2p-r)_\nu\Box+g_{\mu\nu}((2p-r)\cdot\partial)^2\\ &\qquad\qquad-((2p-r)_\mu\partial_\nu+\partial_\mu(2p-r)_\nu)\,(2p-r)\cdot\partial)\,(r\cdot z)^2\\ &\qquad+(((2p-r)_\mu r_\nu\Box+g_{\mu\nu}(2p-r)\cdot\partial\,r\cdot\partial-(2p-r)_\mu\partial_\nu\,r\cdot\partial-\partial_\mu r_\nu(2p-r)\cdot\partial)\\ &\qquad\qquad+(r_\mu(2p-r)_\nu\Box+g_{\mu\nu}(2p-r)\cdot\partial\,r\cdot\partial\\ &\qquad\qquad\quad-\partial_\mu(2p-r)_\nu\,r\cdot\partial-r_\mu\partial_\nu(2p-r)\cdot\partial))\;\;(2p-r)\cdot z\;r\cdot z\\ &\qquad+(r_\mu r_\nu\Box+g_{\mu\nu}(r\cdot\partial)^2\\ &\qquad\qquad-(r_\mu\partial_\nu+\partial_\mu r_\nu)\,r\cdot\partial)\;\;((2p-r)\cdot z)^2)\;\;(p\cdot z)^{J-4}\;E^{(i)}_{J-2,n}(z^2)\,, \end{aligned} \tag{3.26}$$

where $\mathcal{C}$ is the normalization factor that will compensate for the double counting. After using equation (2.10), it is straight forward to see that the four terms coming from $\mathcal{C}_{\alpha\beta}(2p-r,r)$ all give the same value. This means that we should set $\mathcal{C}=\frac{1}{4}$ in the above expression, which now becomes

$$\begin{aligned} &\sum_{J=1}^{\infty}\sum_{n=0}^{J}\sum_{i=1}^{u_J}\langle p'||\hat{O}^{(i)(J,n)}||p\rangle_{(\mu_0)}\;2^{-(J-2)}\;(2-\zeta)^{J-n}\;\zeta^{n}\\ &\qquad(p_\mu p_\nu\Box+g_{\mu\nu}(p\cdot\partial)^2-(p_\mu\partial_\nu+\partial_\mu p_\nu)\,p\cdot\partial)\;(p\cdot z)^{J-2}\;E^{(i)}_{J-2,n}(z^2)\,. \end{aligned} \tag{3.27}$$

Therefore we have, in summary, for the asymmetric matrix elements,

$$\begin{aligned} &\langle p'|T\,j_\mu(-\frac{z}{2})j_\nu(\frac{z}{2})|p\rangle\longrightarrow\\ &\sum_{J=1}^{\infty}\sum_{n=0}^{J}\sum_{i=1}^{u_J}\langle p'||\hat{O}^{(i)(J,n)}||p\rangle_{(\mu_0)}\Bigg(\frac{(2-\zeta)^{J-n}\zeta^n}{2^J}\hat{A}_{\mu\nu}\;F^{(i)}_{J,n}(z^2)\;(p\cdot z)^J\\ &\quad+\frac{(2-\zeta)^{J-n}\zeta^n}{2^{J-2}}(p_\mu p_\nu\Box+g_{\mu\nu}(p\cdot\partial)^2-(p_\mu\partial_\nu+\partial_\mu p_\nu)\,p\cdot\partial)\;E^{(i)}_{J-2,n}(z^2)\;(p\cdot z)^{J-2}\Bigg)\,. \end{aligned}$$

The amplitude $\mathrm{T}_{\mu\nu}$ is the Fourier transform of the matrix elements with momentum $\overline{q}$ (see equation 2.18). Under the Fourier transform, we have, as in the forward case (see section 1.3.3, note the difference in sign convention) $z_\mu \Rightarrow i\frac{\partial}{\partial \overline{q}_\mu}$. Therefore, by taking the logarithmic derivative of $\overline{q}^2$ of the Fourier transformed Wilson coefficients, we have

$$\begin{aligned}
&\frac{1}{2^J}\int d^4z\, e^{-i\bar{q}\cdot z}(p\cdot z)^J E^{(i)}_{J,n}(z^2) \\
&= \frac{1}{2^J}(ip_\mu\frac{\partial}{\partial \overline{q}_\mu})^J \int d^4z\, e^{-i\bar{q}\cdot z} E^{(i)}_{J,n}(z^2) \\
&= ((p\cdot\overline{q})\frac{\partial}{\partial \overline{q}^2})^J\, \tilde{e}^{(i)}_{J,n}(\overline{q}^2) \\
&= \left(\frac{p\cdot\overline{q}}{\overline{Q}^2}\right)^J \left(-i\overline{q}^2\frac{\partial}{\partial\overline{q}^2}\right)^J \tilde{e}^{(i)}_{J,n}(\overline{q}^2)\,, \qquad (3.28)
\end{aligned}$$

where $\tilde{e}^{(i)}_{J,n}$ is the Fourier transform of $E^{(i)}_{J,n}$. We also have a similar equation for F.

At the same time, by a integration by parts it is straight forward to show that, as in the forward case, the derivatives in the tensor structure operator will simply turn into factors of the $\overline{q}$ momentum, namely, $\partial_\mu \Rightarrow -i\overline{q}_\mu$. We will drop the $\hat{A}_{\mu\nu}$ terms and concentrate on the explicit calculation of the structure function T_2. Again the necessary terms generated from $\hat{A}_{\mu\nu}$ and thus the structure function T_1 will be obtained from a Callen-Gross relationship once we know T_2. We obtain the following results for the expression of the amplitude in the short distance limit:

$$\begin{aligned}
\mathrm{T}_{\mu\nu}(p,q',q) &= i\int d^4z\, e^{-i\overline{q}\cdot z}\, \langle p'|T\, j_\mu(-\frac{z}{2})j_\nu(\frac{z}{2})|p\rangle \\
&= -\, i\sum_{J,n,i}\langle p'||\hat{O}^{(i)(J,n)}||p\rangle_{(\mu_0)}\,(2-\zeta)^{J-n}\,\zeta^n \\
&\quad\cdot(\hat{A}_{\mu\nu} terms + \ (p_\mu p_\nu\overline{q}^2 + g_{\mu\nu}(p\cdot\overline{q})^2 - (p_\mu\overline{q}_\nu + \overline{q}_\mu p_\nu)\, p\cdot\overline{q}) \\
&\qquad\cdot\left(\frac{p\cdot\overline{q}}{\overline{Q}^2}\right)^{J-2}\left(-i\overline{q}^2\frac{\partial}{\partial\overline{q}^2}\right)^{J-2}\tilde{e}^{(i)}_{J-2,n}(\overline{q}^2)\,)\,. \qquad (3.29)
\end{aligned}$$

From equations 2.6 and 2.19 it is obvious

$$p \cdot q' = p \cdot \overline{q} = p \cdot q \,. \tag{3.30}$$

Thus for the tensor structure generated by $\hat{B}_{\mu\nu}$ (3.10) in equation 3.29 we have

$$\begin{aligned}
&p_\mu p_\nu \overline{q}^2 + g_{\mu\nu}(p \cdot \overline{q})^2 - (p_\mu \overline{q}_\nu + \overline{q}_\mu p_\nu) p \cdot \overline{q} \\
&\quad = \overline{q}^2 \left(p_\mu p_\nu + g_{\mu\nu} \frac{(p \cdot q)^2}{\overline{q}^2} - \frac{p \cdot q}{\overline{q}^2} \left(p_\mu \overline{q}_\nu + \overline{q}_\mu p_\nu \right) \right) \\
&\quad = \overline{q}^2 \left(p_\mu p_\nu - \frac{p \cdot q}{\overline{q}^2} (p_\mu q_\nu - \frac{\zeta}{2}\, p_\mu p_\nu + q'_\mu p_\nu + \frac{\zeta}{2} p_\mu p_\nu) \right. \\
&\qquad\qquad \left. + \; g_{\mu\nu} \frac{(p \cdot q)^2}{\overline{q}^2} + q'_\mu q_\nu \frac{(p \cdot q)}{\overline{q}^2}^2 - q'_\mu q_\nu \frac{(p \cdot q)}{\overline{q}^2}^2 \right) \\
&\quad = \overline{q}^2 \left(p_\mu p_\nu - \frac{p \cdot q}{\overline{q}^2} \left(p_\mu q_\nu + q'_\mu p_\nu \right) + \frac{(p \cdot q)}{\overline{q}^2}^2 q'_\mu q_\nu \right) + \frac{(p \cdot q)^2}{(\overline{q}^2)^2} \left(g_{\mu\nu} - \frac{q'_\mu q_\nu}{\overline{q}^2} \right) .
\end{aligned} \tag{3.31}$$

Also from equation 3.13 we have

$$\omega = (2 - \zeta) \frac{p \cdot q}{\overline{Q}^2}, \qquad \nu = \zeta \frac{p \cdot q}{\overline{Q}^2} \;. \tag{3.32}$$

Therefore, we can rewrite the expression of $\mathrm{T}_{\mu\nu}$ (3.29) as

$$\begin{aligned}
\mathrm{T}_{\mu\nu}(p, q', q) &= i \int d^4 z \, e^{-i\overline{q} \cdot z} \, \langle p' | T \, j_\mu(-\frac{z}{2}) j_\nu(\frac{z}{2}) | p \rangle \\
&= -i \sum_{J,n,i} \langle p' || \hat{O}^{(i)(J,n)} || p \rangle_{(\mu_0)} \, \omega^{J-n} \, \nu^n \; (\; \hat{A}_{\mu\nu} terms \\
&\quad + \; (\frac{\overline{Q}^2}{p \cdot q})^2 \overline{q}^2 \left(\frac{(p \cdot q)^2}{(\overline{q}^2)^2} \left(g_{\mu\nu} - \frac{q'_\mu q_\nu}{\overline{q}^2} \right) \right. \\
&\qquad \left. + \left(p_\mu p_\nu - \frac{p \cdot q}{\overline{q}^2} (p_\mu q_\nu + q'_\mu p_\nu) + \frac{(p \cdot q)}{\overline{q}^2}^2 q'_\mu q_\nu \right) \left(-i \overline{q}^2 \frac{\partial}{\partial \overline{q}^2} \right)^{J-2} \tilde{e}^{(i)}_{J-2,n}(\overline{q}^2) \; \right) .
\end{aligned} \tag{3.33}$$

Note we could not obtain the explicit expression of the $\hat{A}_{\mu\nu}$ terms because of its general form. However, there is a term of the correct tensor structure (see equation

2.44) generated from the second term. This is another indication of the Callen-Gross relationship that relates the two invariant amplitudes.

Now let us define the Wilson coefficients in momentum space $\tilde{E}$ as

$$\frac{p\cdot q}{\overline{Q}^2} i(\frac{1}{\overline{q}^2})^2 \tilde{E}^{(i)}_{J,n} = (-i\overline{q}^2\frac{\partial}{\partial\overline{q}^2})^J \int d^4z\, e^{-i\overline{q}\cdot z} E^{(i)}_{J,n}, \tag{3.34}$$

where we have pulled out an explicit factor of $\frac{p\cdot q}{\overline{Q}^2}$ to make the form of $\tilde{E}$ simple (see equation 4.60). We will have

$$\begin{aligned} \mathrm{T}_{\mu\nu} \;=\; & i^2 \sum_{J,n,i} \langle p'||\hat{O}^{(i)(J,n)}||p\rangle_{(\mu_0)} \omega^{J-n}\nu^n \left((-g_{\mu\nu} + \frac{q'_\mu q_\nu}{\overline{q}^2})\, \frac{p\cdot q}{\overline{Q}^2}\, \tilde{F'}^{(i)}_{J,n}(\alpha_s, \frac{\overline{Q}}{\mu_0}) \right. \\ & \left. + \frac{1}{p\cdot q} \left(p_\mu p_\nu - \frac{p\cdot q}{\overline{q}^2}(p_\mu q_\nu + q'_\mu p_\nu) + (\frac{p\cdot q}{\overline{q}^2})^2 q'_\mu q_\nu \right) \tilde{E}^{(i)}_{J-2,n}(\alpha_s, \frac{\overline{Q}}{\mu_0}) \right) . \end{aligned} \tag{3.35}$$

$\tilde{F'}$ should be a linear combination of $\tilde{E}$ and a similarly defined $\tilde{F}$. We have explicitly indicated their dependence on the factorization scale μ_0.

By comparing with equation 2.44 we finally obtain the explicit expression of the invariant amplitudes, in terms of reduced matrix elements and their corresponding Wilson coefficients, as the following:

$$\sqrt{1-\zeta}\,\mathrm{T}_1 = -\sum_{J,n,i} \langle p'||\hat{O}^{(i)(J,n)}||p\rangle_{(\mu_0)} \omega^{J-n}\nu^n \tilde{F'}^{(i)}_{J,n}(\alpha_s, \frac{\overline{Q}}{\mu_0}) \tag{3.36}$$

$$\frac{p\cdot q}{M^2}\sqrt{1-\zeta}\,\mathrm{T}_2 = -\sum_{J,n,i} \langle p'||\hat{O}^{(i)(J,n)}||p\rangle_{(\mu_0)} \omega^{J-n}\nu^n \tilde{E}^{(i)}_{J-2,n}(\alpha_s, \frac{\overline{Q}}{\mu_0}) \ . \tag{3.37}$$

We will concentrate on the discussion of T_2 from now on. We always regard T_1 as being obtained from T_2 from a Callen-Gross relationship that can readily be extracted from the above discussion:

$$\mathrm{T}_1 = \frac{p\cdot q}{M^2}\mathrm{T}_2\,. \tag{3.38}$$

Chapter 4

Renormalization Group Analysis

In this chapter, We perform a renormalization group (RG) analysis of the non-forward amplitude. We write down the renormalization group equation (RGE) of the operators and their corresponding Wilson coefficients of the operator product expansion (OPE) of the amplitude, and find the formal solution to the RGE. Then we compute the evolution kernels that will generate the anomalous dimensions and discuss their properties. Finally we will calculate explicitly the lowest order Wilson coefficients.

4.1 Renormalization Group Equation

As stated in the last chapter, we will concentrate on the invariant amplitude T_2, and T_1 can be obtained from the Callen-Gross relationship (3.38). T_2 is analogous to the structure function F_2 ($\nu \mathrm{W}_2$) (see section 1.2.2), however, the analogy is not complete because of the subtleties in the discussion of dispersion relationship later on.

Recall that T_2 is given, after OPE, as

$$\frac{p\cdot q}{M^2}\sqrt{1-\zeta}\,\mathrm{T}_2 = -\sum_{J,n,i}\langle p'||\hat{O}^{(i)(J,n)}||p\rangle_{(\mu_0)}\omega^{J-n}\nu^n\tilde{E}^{(i)}_{J-2,n}(\alpha_s,\frac{\overline{Q}}{\mu_0}) \ . \tag{4.1}$$

It can be regarded as a double distribution function. In the usual language it is the series sum of "double" moments in terms of the two new moment variable ω and ν that we defined earlier (see equation 3.13):

$$\mathrm{T}_2 = \sum_{J,n}\omega^{J-n}\nu^n\mathrm{T}_2^{(J,n)}, \tag{4.2}$$

with the double moments $\mathrm{T}_2^{(J,n)}$ defined as

$$\sqrt{1-\zeta}\frac{p\cdot q}{M^2}\mathrm{T}^{(J,n)}_{2,(\overline{Q},\mu_0)} = -\sum_i\langle p'||\hat{O}^{(i)(J,n)}||p\rangle_{(\mu_0)}\tilde{E}^{(i)}_{J-2,n}(\alpha_s,\frac{\overline{Q}}{\mu_0}) \tag{4.3}$$

We will eventually analytically continue in J, but leave n as discrete. The discussion of analyticity properties of T_2 is almost identical to that presented for the forward case (see section 1.3.3) with minor adjustments. Explicitly writing out the spin sum of $\mathrm{T}_{\mu\nu}$ we have (see equations 2.16 and 2.18)

$$\begin{aligned}
\mathrm{T}_{\mu\nu}(p,q',q) &= i\frac{1}{2}\sum_s\int d^4z\, e^{-i\overline{q}\cdot z}\,\langle p',s|T\,j_\mu(-\frac{z}{2})j_\nu(\frac{z}{2})|p,s\rangle \\
&= i\int d^4z\, e^{-iq\cdot z}\langle p'|T\,j_\mu(-z)j_\nu(0)|p\rangle \\
&\overset{z\to -z}{\equiv} i\int d^4z\, e^{iq\cdot z}\langle p'|T\,j_\mu(z)j_\nu(0)|p\rangle \\
&= i\sum_r\int d^4z\, e^{iq\cdot z}\{\,\theta(z_0)\langle p'|j_\mu(z)|r\rangle\langle r|j_\nu(0)|p\rangle \\
&\quad +\theta(-z_0)\langle p'|j_\nu(0)|r\rangle\langle r|j_\mu(z)|p\rangle\,\} \\
&= i\sum_r\int d^4z\, e^{iq\cdot z}\{\,\theta(z_0)e^{i(p'-p_r)\cdot z}\langle p'|j_\mu(0)|r\rangle\langle r|j_\nu(0)|p\rangle \\
&\quad +\theta(-z_0)e^{-i(p-p_r)\cdot z}\langle p'|j_\nu(0)|r\rangle\langle r|j_\mu(0)|p\rangle\,\}\,,
\end{aligned} \tag{4.4}$$

where we have used the translation operators (see equation 1.18) in the last step. Performing the spacial integration yields a 3-d δ-function and we have, similar to the forward case,

$$\begin{aligned} \mathrm{T}_{\mu\nu} &= i(2\pi)^3 \sum_r \int dz_0 (\delta^3(\vec{q} + \vec{p}\,' - \vec{p}_r) e^{i(q_0 + p'_0 - p_{r,0}) \cdot z_0} \theta(z_0) \langle p' | j_\mu(0) | r \rangle \langle r | j_\nu(0) | p \rangle \\ &\quad + \delta^3(\vec{q} - \vec{p} + \vec{p}_r) e^{i(q_0 - p_0 + p_{r,0}) \cdot z_0} \theta(-z_0) \langle p' | j_\nu(0) | r \rangle \langle r | j_\mu(0) | p \rangle) \\ &= -(2\pi)^3 \sum_r \left(\frac{\delta^3(\vec{q} + \vec{p}\,' - \vec{p}_r) \langle p' | j_\mu(0) | r \rangle \langle r | j_\nu(0) | p \rangle}{q_0 + p'_0 - p_{r,0} + i\epsilon} \right. \\ &\quad \left. - \frac{\delta^3(\vec{q} - \vec{p} + \vec{p}_r) \langle p' | j_\nu(0) | r \rangle \langle r | j_\mu(0) | p \rangle}{q_0 - p_0 + p_{r,0} - i\epsilon} \right) . \end{aligned} \tag{4.5}$$

Again by looking at the poles in q_0 we can discuss the analytic properties of the amplitude. The first term gives rise to poles at

$$q_0 + p'_0 = p_{r,0} \equiv E_r \,, \tag{4.6}$$

which means that (cf 1.3.3)

$$(q_0 + p'_0)^2 = E_r^2 = M_r^2 + \vec{p}_r^{\,2} = M_r^2 + (\vec{q} + \vec{p}\,')^2 \,, \tag{4.7}$$

where in the last step we have used the spacial δ-function. M_r is again the invariant mass of the intermediate state r. Thus we have, recall equations 2.1 and 2.7,

$$M_r^2 = (q + p')^2 = (q' + p)^2 = q'^2 + 2p \cdot q \,. \tag{4.8}$$

Rewrite q'^2 as

$$q'^2 = \frac{1}{2}(q'^2 + q^2) - \frac{1}{2}(q^2 - q'^2) \tag{4.9}$$

and use equations 2.9 and 3.13, we arrive at

$$\begin{aligned} M_r^2 &= (q' + p)^2 = \overline{q}^2 + (2p - r) \cdot q \\ &= \overline{q}^2 (1 - \omega) \,, \end{aligned} \tag{4.10}$$

and thus the poles in ω are at

$$\omega_r = 1 + \frac{M_r^2}{\overline{Q}^2}\,. \tag{4.11}$$

The second term of the spectral expansion of the amplitude gives poles at

$$q_0 - p_0 = -p_{r,0} = -E_r \tag{4.12}$$

and after similar discussion results in

$$\begin{aligned} M_r^2 &= (q-p)^2 = q^2 - 2p\cdot q = \overline{q}^2 - (2p-r)\cdot q \\ &= \overline{q}^2(1+\omega)\,, \end{aligned} \tag{4.13}$$

which means that there are also poles in ω at

$$\omega_r = -1 - \frac{M_r^2}{\overline{Q}^2}\,. \tag{4.14}$$

We see again that the momentum scale is set naturally by $\overline{Q}^2 = -\overline{q}^2$.

It is clear that in this non-forward case, $\mathrm{T}_{\mu\nu}$ also has an analytic circle of unit radius in the ω-plane and has a Taylor expansion in ω. The discussion on the dispersion relationship for T_2 then follows almost identically to that of the forward case (see section 1.3.3)

We can indeed use the dispersion relationship to analytically extend the discussion from the region of $\omega \leq 1$ where $\mathrm{T}_{\mu\nu}$ is analytic but kinematically unphysical to the physical region of the Bjorken limit where $0 \leq (\omega)^{-1} \leq 1$ in the same manner as in the forward case. And we find that the inverse of the expansion 4.2 is

$$\begin{aligned} \mathrm{T}_2^{(J,n)} &= \int_c \frac{d\omega}{\omega} \int_c \frac{d\nu}{\nu} (\frac{1}{\omega})^{J-n} (\frac{1}{\nu})^n \mathrm{T}_2(\overline{Q}^2, \omega, \nu) \\ &= 2i \int_1^\infty \frac{d\omega}{\omega} \int_c \frac{d\nu}{\nu} (\frac{1}{\omega})^{J-n} (\frac{1}{\nu})^n \mathrm{W}_2(\overline{Q}^2, \omega, \nu)\,, \end{aligned} \tag{4.15}$$

where c is any contour in the $\omega(\nu)$ plane that encloses the origin and $\mathrm{W}_2 = \mathrm{Im}\,\mathrm{T}_2$. Note that in obtaining the second equality we have used the fact that T_2 is even in ω. Also note that the moment index of ω in this case is, in our notation, $J - n$.

T_2 now as a double distribution function is given by a Mellin transformation:

$$\mathrm{T}_2(\overline{Q}^2, \omega, \nu) = \int_c \frac{dJ}{2\pi i} \sum_{n=0}^{\infty} e^{(J-n)\log\omega}\, \nu^n\, \mathrm{T}_2^{(J,n)} \tag{4.16}$$

with the contour c lying parallel to the imaginary axis in the $J - n$ plane and to the right of all singularities in that plane.

Under renormalization, the operators scale according to a renormalization group equation

$$\mu^2 \frac{d}{d\mu^2} \hat{O}^{(i)(J,n)} = \sum_{n'=n}^{J} \sum_{i'} \tilde{\gamma}^{ii'}_{nn'} \hat{O}^{(i')(J,n')} \tag{4.17}$$

where $\tilde{\gamma}$ is the anomalous dimension matrix which acts in the product space $J \otimes u_J$. Because the mixing of these operators is such that $n \to n' \geq n$ (see equation 3.16 the matrix $\tilde{\gamma}$ is upper triangular in the J dimensional n-space.

Since T_2 is a physical quantity it should not depend on the factorization scale. The Wilson coefficients obey a similar renormalization group equation thus cancelling out the scale dependence of the operators:

$$\mu^2 \frac{d}{d\mu^2} \tilde{E}^{(i)}_{J,n}(\alpha_s(\mu), \frac{\overline{Q}}{\mu}) = -\sum_{n'=0}^{n} \sum_{i'} \tilde{E}^{(i')}_{J,n'}\ \ \tilde{\gamma}^{i'i}_{n'n}. \tag{4.18}$$

We can write the solution to this RG equation as

$$\tilde{E}^{(i)}_{J,n}(\alpha_s(\mu), \frac{\overline{Q}}{\mu}) = \sum_{n'=0}^{n} \sum_{i'} \tilde{E}^{(i')}_{J,n'}(\alpha_s(\overline{Q}), 1)\ \ \mathrm{M}^{i'i}_{n'n}(\alpha(\overline{Q}), \frac{\overline{Q}}{\mu}), \tag{4.19}$$

where M is a path ordered exponential of the anomalous dimension matrix, formally given as

$$\mathrm{M}^{i'i}_{n'n}(\alpha_s(\overline{Q}), \frac{\overline{Q}}{\mu}) = \left(\mathcal{P}\exp(-\int_{\mu^2}^{\bar{Q}^2} \tilde{\gamma}(\alpha_s(\lambda^2)) \frac{d\lambda^2}{\lambda^2})\right)^{i'i}_{n'n}. \tag{4.20}$$

Therefore by substitution the double moments are given by

$$\sqrt{1-\zeta}\frac{p\cdot q}{M^2}\mathrm{T}^{(J,n)}_{2,(\overline{Q},\mu_0)} = -\sum_{n'=0}^{n}\sum_{i'i}\tilde{E}^{(i')}_{J-2,n'}(\alpha_s(\overline{Q}),1)\ \mathrm{M}^{i'i}_{n'n}(\alpha_s(\overline{Q}),\frac{\overline{Q}}{\mu_0})\ \langle p'||\hat{O}^{(i)(J,n)}||p\rangle_{(\mu_0)}\,. \tag{4.21}$$

In principle, for the high energy scattering process, if we know the reduced matrix elements at a given momentum scale μ, we can calculate the Wilson coefficients order by order in perturbation theory at some high momentum scale Q, at which the scattering actually takes place and perturbative QCD is valid, then use the anomalous dimensions, also computed to some fixed order in perturbation theory, to evolve along the momentum scale to μ so that we can use these reduced matrix elements and obtain (theoretical predictions of) the invariant amplitude, the amplitude and eventually the cross section.

The matrix elements can come from two sources in general. They can either come from a first-principle calculation in lattice QCD, or from a high energy experiment. The former is phenomenologically not quite feasible yet. For the latter, we actually measure the cross section of the high energy scattering at a certain high momentum scale, use the above mentioned procedure in reverse and phenomenological models to extract the matrix elements, and then by RG analysis we can predict the cross section at any other high momentum scale where factorization is valid and compare the results with actual experimental data at the scale.

In the next two sections, we calculate the anomalous dimension (or rather, evolution kernels from which they can be extracted) and Wilson coefficients in the lowest non-trivial order.

4.2 Anomalous Dimensions

We will use the light cone (LC) gauge and compute the lowest order anomalous dimensions in terms of light cone variables. In the LC gauge, the lowest order anomalous dimensions of the quark and gluon operators are generated by the usual triangle diagrams. These diagrams are shown collectively in figure 15. The graphs where we have more than two quark/gluon lines meeting at the vertex are zero because of the gauge choice. The price we pay is the presence of extra end point colinear divergences in the integration of light cone variables, which, as we will see, will cancel out eventually after we include the self-energy graphs.

We define the momentum fraction variables (which are the moment variables) as

$$\omega = \frac{k_+}{k_{1+}}, \quad \nu = \frac{r_+}{k_{1+}}\,. \tag{4.22}$$

The conventions (cf. ref. [14]) we use for the quark and gluon vertices are

$$\begin{aligned} O_q^{(J,n)(+)}(k, k-r) &= \gamma_+(2k-r)_+^{J-n-1} r_+^n \\ O_g^{(J,n)\alpha\beta}(k, k-r) &= 2g_{\alpha\beta}\, n\cdot k\, n\cdot(k-r)\,(2k-r)_+^{J-n-2} r_+^n\,, \end{aligned} \tag{4.23}$$

where n is the LC null vector (see equation 1.119).

4.2.1 Quark-Quark Anomalous Dimension

The detailed version of the first diagram in figure 15 is shown in figure 16, where α and β are Lorentz indices; a, a', b and b' are color indices for the quarks while i labels the gluon line. In contrast to figure 9 of the forward case, the gluon is not on-shell.

Using standard Feynman rules and the LC gauge we can write down its value as

$$\gamma_{qq}^{+,1} = (ig)^2 \sum_{i,b} (T^i_{ab} T^i_{ba'}) \int \frac{d^4k}{(2\pi)^4} \gamma_\alpha \frac{i}{\not{k} - \not{r}} O_q^{(J,n)(+)}(k, k-r) \frac{i}{\not{k}} \gamma_\beta \frac{-iD^{\alpha\beta}(k_1 - k)}{(k_1-k)^2}\,, \tag{4.24}$$

which, after the color algebra (see section 1.4.3) gives (where we have taken the incoming and outgoing quark to have the same color index)

$$\gamma_{qq}^{+,1} = (ig)^2 C_F \int \frac{d^4k}{(2\pi)^4} \gamma_\alpha \frac{i}{\not{k} - \not{r}} O_q^{(J,n)(+)}(k, k-r) \frac{i}{\not{k}} \gamma_\beta \frac{-iD^{\alpha\beta}(k_1 - k)}{(k_1-k)^2}\,, \tag{4.25}$$

where the quark vertex O_q is given in (4.23) and the LC gluon projector (numerator of the LC gluon propagator) (see equation 1.121)

$$D^{\alpha\beta}(k) = g^{\alpha\beta} - \frac{n_\alpha k_\beta + k_\alpha n_\beta}{n \cdot k}\,. \tag{4.26}$$

Again n is the LC null vector. Writing the integral of the loop momentum in terms of light-cone variables (see section 1.4.3) we get

$$\gamma_{qq}^{+,1} = -i\frac{g^2 C_F}{(2\pi)^4} \int d^2\underline{k}\, dk_+ dk_- \frac{(2k-r)_+^{J-n-1} r_+^n A(k,r)}{(k^2+i\epsilon)((k-r)^2+i\epsilon)((k_1-k)^2+i\epsilon)}\,, \tag{4.27}$$

where

$$A(k,r) = \gamma_\alpha(\not{k} - \not{r})\gamma_+ \not{k} \gamma_\beta D^{\alpha\beta}(k_1 - k)\,. \tag{4.28}$$

We perform the k_- integration first, in terms of the poles of the integrand, which come from the denominator factors

$$\begin{aligned} k^2 + i\epsilon &= 2k_+k_- - \underline{k}^2 + i\epsilon\,, \\ (k-r)^2 + i\epsilon &= 2(k-r)_+k_- - \underline{k}^2 + i\epsilon\,, \\ (k_1-k)^2 + i\epsilon &= 2(k-k_1)_+k_- - \underline{k}^2 + i\epsilon\,. \end{aligned} \tag{4.29}$$

However, the positions of poles in k_- are different when k_+ is in different regions. More specifically, there are four regions we need to consider:

1) $k_+ < 0$, by setting the denominators to zero we find that the poles are at, respectively,

$$k_- = \frac{\underline{k}^2}{2k_+} + i\epsilon\,, \quad k_- = \frac{\underline{k}^2}{2(k-r)_+} + i\epsilon\,, \quad k_- = \frac{\underline{k}^2}{2(k-k_1)_+} + i\epsilon\,. \tag{4.30}$$

All three poles lie above the real axis of the complex k_- plane (see figure 17 (a)). We can complete the contour of integration of k_- in the lower-half-plane and clockwise, enclosing no poles and thus the integration gives zero value.

2) $0 < k_+ < r_+ (< k_{1+})$, the poles are at, respectively,

$$k_- = \frac{\underline{k}^2}{2k_+} - i\epsilon\,, \quad k_- = \frac{\underline{k}^2}{2(k-r)_+} + i\epsilon\,, \quad k_- = \frac{\underline{k}^2}{2(k-k_1)_+} + i\epsilon\,. \tag{4.31}$$

The first pole is below the real axis while the other two are above it (see figure 17 (b)). We can complete the integration contour again in the lower-half-plane and clockwise, picking up the pole at $k_- = \frac{\underline{k}^2}{2k_+} - i\epsilon \equiv k_-^{(1)}$. The value of the integration would then be the residue of the $k_-^{(1)}$ pole with a factor of $-2\pi i$.

3) $r_+ < k_+ < k_{1+}$, the poles are at, respectively,

$$k_- = \frac{\underline{k}^2}{2k_+} - i\epsilon\,, \quad k_- = \frac{\underline{k}^2}{2(k-r)_+} - i\epsilon\,, \quad k_- = \frac{\underline{k}^2}{2(k-k_1)_+} + i\epsilon\,. \tag{4.32}$$

The first two poles are below the real axis while the third one is above it (see figure 17 (c)). We can complete the integration contour now in the upper-half-plane and counterclockwise, picking up the pole at $k_- = \frac{\underline{k}^2}{2(k_+ - k_{1+})} + i\epsilon \equiv k_-^{(2)}$. The value of the integration would then be the residue of the $k_-^{(2)}$ pole with a factor of $2\pi i$.

4) $k_+ > k_{1+}$, the poles are at, respectively,

$$k_- = \frac{\underline{k}^2}{2k_+} - i\epsilon\,, \quad k_- = \frac{\underline{k}^2}{2(k-r)_+} - i\epsilon\,, \quad k_- = \frac{\underline{k}^2}{2(k-k_1)_+} - i\epsilon\,. \tag{4.33}$$

All three poles lie below the real axis of the k_- plane (see figure 17 (d)). We can complete the contour of integration of k_- in the upper-half-plane and counterclockwise, again enclosing no poles and the integration gives zero value.

Thus the k_+ integration should be performed in two regions:

$$\begin{aligned}\gamma_{qq}^{+,1} &= -i\frac{g^2C_F}{(2\pi)^4}\int d^2\underline{k}\left(\int_0^{r_+}dk_+ + \int_{r_+}^{k_{1+}}dk_+\right)\int\frac{dk_-\ (2k-r)_+^{J-n-1}r_+^n\ A(k,r)}{(k^2+i\epsilon)((k-r)^2+i\epsilon)((k_1-k)^2+i\epsilon)}\\ &= -i\frac{\alpha_sC_F}{(2\pi)^2}\int d\underline{k}^2\left(\int_0^{r_+}dk_+ + \int_{r_+}^{k_{1+}}dk_+\right)(2k-r)_+^{J-n-1}r_+^n\\ &\quad\cdot\int dk_-\frac{A(k,r)}{(2k_+k_- - \underline{k}^2+i\epsilon)(2(k-r)_+k_- - \underline{k}^2+i\epsilon)(2(k-k_1)_+k_- - \underline{k}^2+i\epsilon)}\,.\end{aligned} \tag{4.34}$$

Evaluating the contour integral of k_- at the poles specified above, we have

$$\begin{aligned}\gamma_{qq}^{+,1} =& -i\frac{\alpha_sC_F}{(2\pi)^2}\int d\underline{k}^2\left[\int_0^{r_+}dk_+\frac{(-2\pi i)}{2k_+}\left(\frac{(2k-r)_+^{J-n-1}r_+^n\,A(k,r)}{(2(k-r)_+k_- - \underline{k}^2)(2(k-k_1)_+k_- - \underline{k}^2)}\right)\Bigg|_{k_-=k_-^{(1)}}\right.\\ &\left.+\int_{r_+}^{k_{1+}}dk_+\frac{(2\pi i)}{2(k-k_1)_+}\left(\frac{(2k-r)_+^{J-n-1}r_+^n\,A(k,r)}{(2k_+k_- - \underline{k}^2)(2(k-r)_+k_- - \underline{k}^2)}\right)\Bigg|_{k_-=k_-^{(2)}}\right].\end{aligned} \tag{4.35}$$

Substituting the pole values into the denominators and simplify them first, we get, for the pole $k_-^{(1)} = \frac{\underline{k}^2}{2k_+}$,

$$\begin{aligned}&(2(k-r)_+k_- - \underline{k}^2)(2(k-k_1)_+k_- - \underline{k}^2)\Big|_{k_-=k_-^{(1)}}\\ &= \left(\underline{k}^2\frac{(k-r)_+}{k_+} - \underline{k}^2\right)\left(\underline{k}^2\frac{(k-k_1)_+}{k_+} - \underline{k}^2\right) = (\underline{k}^2)^2\left(1 - \frac{(k-r)_+}{k_+}\right)\left(1 + \frac{(k_1-k)_+}{k_+}\right)\\ &= \frac{1}{\underline{k}^4}\,\frac{k_+}{(k_+ - (k-r)_+)(k_+ + (k-r)_+)} = \frac{1}{\underline{k}^4}\,\frac{k_+}{r_+k_{1+}}\,,\end{aligned} \tag{4.36}$$

while for the pole $k_-^{(2)} = \frac{\underline{k}^2}{2(k_+ - k_{1+})}$,

$$
\begin{aligned}
&(2k_+k_- - \underline{k}^2)(2(k-r)_+k_- - \underline{k}^2)\Big|_{k_-=k_-^{(2)}} \\
&= (\underline{k}^2\frac{k_+}{(k-k_1)_+} - \underline{k}^2)(\underline{k}^2\frac{(k-r)_+}{(k-k_1)_+} - \underline{k}^2) = (\underline{k}^2)^2\left(1+\frac{k_+}{(k_1-k)_+}\right)\left(1+\frac{(k-r)_+}{(k_1-k)_+}\right) \\
&= \frac{1}{\underline{k}^4}\frac{(k_1-k)_+}{((k_1-k)_+ + k_+)((k_1-k)_+ + (k-r)_+)} = \frac{1}{\underline{k}^4}\frac{(k_1-k)_+}{(k-r)_+k_{1+}} \,. \qquad (4.37)
\end{aligned}
$$

Thus we arrive at

$$
\begin{aligned}
\gamma_{qq}^{+,1} = -i\frac{\alpha_s C_F}{(2\pi)^2}\int\frac{d^2\underline{k}}{\underline{k}^4}\Bigg(&\frac{-2\pi i}{2r_+k_{1+}}\int_0^{r_+} dk_+k_+A^{(1)}(2k-r)_+^{J-n-1}r_+^n \\
&+\frac{2\pi i}{-2k_{1+}(k_1-r)_+}\int_{r_+}^{k_{1+}} dk_+(k_1-k)_+A^{(2)}(2k-r)_+^{J-n-1}r_+^n\Bigg)\,, \qquad (4.38)
\end{aligned}
$$

where $A^{(1)}$ and $A^{(2)}$ are $A(k,r)$ evaluated at the poles $k_-^{(1)}$ and $k_-^{(2)}$, respectively.

We are interested in the logarithmic divergence in the graphs so in computing the Dirac spinors we keep only the leading powers of $\underline{k}$. This leading power turns out to be quadratic, which means we will keep in the following calculation of $A^{(1)}$ and $A^{(2)}$ only terms that are proportional to $\underline{k}^2$.

Defining ω and ν as before in (4.22), we have, at the poles,

$$
\begin{aligned}
k_-^{(1)} = \frac{\underline{k}^2}{2k_+} = \frac{\underline{k}^2}{2k_{1+}\omega} \qquad &\& \quad k^2 = 2k_+k_- - \underline{k}^2 = 0, \\
k_-^{(2)} = \frac{\underline{k}^2}{2(k_+-k_{1+})} = \frac{-\underline{k}^2}{2k_{1+}(1-\omega)} \qquad &\& \quad k^2 = -\frac{\underline{k}^2}{(1-\omega)}.
\end{aligned}
$$

Recall that $A(k,r)$ can be written into two terms:

$$
\begin{aligned}
A(k,r) &= \gamma_\alpha(\not{k} - \not{r})\gamma_+\not{k}\gamma_\beta D^{\alpha\beta}(k_1-k) \\
&= \gamma_\alpha\not{k}\gamma_+\not{k}\gamma_\beta D^{\alpha\beta}(k_1-k) - \gamma_\alpha\not{r}\gamma_+\not{k}\gamma_\beta D^{\alpha\beta}(k_1-k) \\
&\equiv A_I - A_{II} \qquad (4.39)
\end{aligned}
$$

Using the anti-commutation relationships of the Dirac matrices in light cone variables (see equations 1.115 and 1.117) we can reduce the first term (A_I) to

$$\gamma_\alpha \not{k}\gamma_+ \not{k}\gamma_\beta D^{\alpha\beta}(k_1-k) \;=\; 2k_+\gamma_\alpha \not{k}\gamma_\beta D^{\alpha\beta}(k_1-k) - k^2\gamma_\alpha\gamma_+\gamma_\beta D^{\alpha\beta}(k_1-k). \tag{4.40}$$

At the pole $k_-^{(1)}$, because $k^2=0$, we have

$$A_I = 2k_+\gamma_\alpha \not{k}\gamma_\beta D^{\alpha\beta}(k_1-k) \equiv A_I^{(1)}\,. \tag{4.41}$$

Expanding $D^{\alpha\beta}(k_1-k)$ and again using anti-commutation rules and the fact that $k^2=0$ we have

$$\begin{aligned} A_I^{(1)} &= 2k_+\gamma_\alpha \not{k}\gamma^\alpha - \frac{2k_+}{k_{1+}(1-\omega)}(\gamma_+ \not{k}(\not{k}_1 - \not{k}) + (\not{k}_1 - \not{k}) \not{k}\gamma_+) \\ &= -4k_+ \not{k} - \frac{2\omega}{1-\omega}(\gamma_+ \not{k} \not{k}_1 + \not{k}_1 \not{k}\gamma_+ - 2\gamma_+ k^2) \\ &= -4k_+(k_+\gamma_- + k_-\gamma_+ - \underline{k}\cdot\underline{\gamma}) - \frac{2\omega}{1-\omega}(\gamma_+ \not{k} \not{k}_1 + \not{k}_1 \not{k}\gamma_+)\,. \end{aligned} \tag{4.42}$$

As stated before, we want to keep only the leading power in $\underline{k}$. It is clear that in (4.42) the leading power of $\underline{k}$ is quadratic and can only come from a term that contains the factor k_-. In addition, since we are computing the quark-quark anomalous dimension, only terms that have similar vertex structure as the original quark vertex O_q in (4.23) contribute. Thus, we keep in (4.42) only terms that are proportional to $\gamma_+ k_-$. By expanding the dot products and inspecting the result, it is straight forward to see that only the second term in the first parenthesis survives and we have

$$A_I^{(1)} = -4k_+k_-\gamma_+ = -2\underline{k}^2\gamma_+\,, \tag{4.43}$$

where we have also used (1.116).

The second term of (4.39) can be similarly reduced by keeping only the terms proportional to $\gamma_+ k_-$:

$$\begin{aligned}
A_{II}^{(1)} &= \gamma_\alpha \not{r}\gamma_+ \not{k}\gamma_\beta D^{\alpha\beta}(k_1 - k) \\
&= \gamma_\alpha \not{r}\gamma_+ \not{k}\gamma^\alpha - \frac{1}{(k_1 - k)_+}(\gamma_+ \not{r}\gamma_+ \not{k}(\not{k}_1 - \not{k}) + (\not{k}_1 - \not{k}) \not{r}\gamma_+ \not{k}\gamma_+) \\
&= 0 - \frac{1}{(k_1 - k)_+}(2r_+\gamma_+(\not{k} \not{k}_1 - k^2) + (\not{k}_1 \not{r} - \not{k} \not{r})2k_+\gamma_+) \\
&= -\frac{1}{(k_1 - k)_+}(2r_+\gamma_+(\gamma_- k_+ \not{k}_1 - 0) + (0 - \not{k} \not{r})2k_+\gamma_+) \\
&= -\frac{1}{(k_1 - k)_+}(0 - 2k_+ \not{k} \not{r}\gamma_+) = \frac{1}{(k_1 - k)_+}2k_+k_-r_+\gamma_+\gamma_-\gamma_+ \\
&= \frac{2r_+}{(k_1 - k)_+}\underline{k}^2\gamma_+ = \frac{2\nu}{1-\omega}\underline{k}^2\gamma_+ \,. \qquad (4.44)
\end{aligned}$$

Note some of the zeros above do not mean numerically zero but rather the contributions of those terms are zero. We use zero (0) this way through out the discussion unless otherwise stated.

Combining the results we have

$$A^{(1)} = -2\underline{k}^2\gamma_+(1 + \frac{\nu}{1-\omega}) \,. \qquad (4.45)$$

The evaluation of $A^{(2)}$ goes parallel but is a little bit more involved since we no longer have $k^2 = 0$ but rather $k^2 = -\frac{\underline{k}^2}{(1-\omega)}$. We outline the main steps in the following:

$$\begin{aligned}
A_I^{(2)} &= \gamma_\alpha \not{k}\gamma_+ \not{k}\gamma_\beta D^{\alpha\beta}(k_1 - k) \\
&= 2k_+\gamma_\alpha \not{k}\gamma_\beta D^{\alpha\beta}(k_1 - k) - k^2\gamma_\alpha\gamma_+\gamma_\beta D^{\alpha\beta}(k_1 - k) \\
&= 2k_+\gamma_\alpha \not{k}\gamma^\alpha - \frac{2k_+}{k_{1+}(1-\omega)}(\gamma_+ \not{k}(\not{k}_1 - \not{k}) + (\not{k}_1 - \not{k}) \not{k}\gamma_+) \\
&\quad -k^2\gamma_\alpha\gamma_+\gamma^\alpha + \frac{k^2}{k_{1+}(1-\omega)}(\gamma_+\gamma_+(\not{k}_1 - \not{k}) + (\not{k}_1 - \not{k})\gamma_+\gamma_+) \\
&= -4k_+ \not{k} - \frac{2\omega}{1-\omega}(\gamma_+ \not{k} \not{k}_1 + \not{k}_1 \not{k}\gamma_+ - 2\gamma_+k^2) + 2k^2\gamma_+ + 0 \,.
\end{aligned}$$

Using identical arguments as those in calculation of $A^{(1)}$, plus the fact that now the leading contribution (of $\underline{k}^2$) can also come from a k^2 factor in addition to k_-, we arrive at

$$
\begin{aligned}
A_I^{(2)} &= -4k_+k_-\gamma_+ - \frac{2\omega}{1-\omega}(-2\gamma_+k^2) + 2k^2\gamma_+ \\
&= -4k_+\frac{-\underline{k}^2}{2(k_+-k_{1+})}\gamma_+ + (\frac{4\omega}{1-\omega}+2)\gamma_+(-\frac{\underline{k}^2}{(1-\omega)}) \\
&= \frac{\underline{k}^2}{(1-\omega)^2}\gamma_+(2\omega(1-\omega) - 4\omega - 2(1-\omega)) \\
&= -\frac{2\underline{k}^2\gamma_+}{(1-\omega)^2}(1+\omega^2)\,. \qquad (4.46)
\end{aligned}
$$

On the other hand,

$$
\begin{aligned}
A_{II}^{(2)} &= \gamma_\alpha \not{r}\gamma_+ \not{k}\gamma_\beta D^{\alpha\beta}(k_1-k) \\
&= \gamma_\alpha \not{r}\gamma_+ \not{k}\gamma^\alpha - \frac{1}{(k_1-k)_+}(\gamma_+ \not{r}\gamma_+ \not{k}(\not{k}_1 - \not{k}) + (\not{k}_1 - \not{k}) \not{r}\gamma_+ \not{k}\gamma_+) \\
&= 0 - \frac{1}{(k_1-k)_+}(2r_+\gamma_+(\not{k}\,\not{k}_1 - k^2) + (\not{k}_1 \not{r} - \not{k}\,\not{r})2k_+\gamma_+) \\
&= -\frac{1}{(k_1-k)_+}(2r_+\gamma_+(\gamma_-k_+ \not{k}_1 - k^2) + (0 - \not{k}\,\not{r})2k_+\gamma_+) \\
&= -\frac{1}{(k_1-k)_+}(-2r_+\gamma_+k^2 - 2k_+k_-r_+\gamma_+\gamma_-\gamma_+) \\
&= \frac{2r_+}{(k_1-k)_+}\gamma_+\frac{-\underline{k}^2}{(1-\omega)} + \frac{4k_+}{(k_1-k)_+}\gamma_+r_+\frac{\underline{k}^2}{2(k_+-k_{1+})} \\
&= -2\underline{k}^2\gamma_+\left(\frac{r_+}{k_{1+}(1-\omega)^2} + \frac{k_+r_+}{k_{1+}^2(1-\omega)^2}\right) \\
&= -\frac{2\nu\underline{k}^2\gamma_+}{(1-\omega)^2}(1+\omega)\,. \qquad (4.47)
\end{aligned}
$$

Again combining the results we have

$$
A^{(2)} = -\frac{2\underline{k}^2\gamma_+}{(1-\omega)^2}(1+\omega^2-\nu(1+\omega))\,. \qquad (4.48)
$$

Thus we have

$$\gamma_{qq}^{+,1} = \frac{\alpha_s}{2\pi} C_F \int \frac{d\underline{k}^2}{\underline{k}^2} \gamma_+ k_{1+}^{J-1} \nu^n \left(\frac{1}{\nu} \int_0^{\nu} d\omega\, (2\omega - \nu)^{J-n-1}\, \omega\, (1 + \frac{\nu}{1-\omega}) \right. \\ \left. + \frac{1}{1-\nu} \int_{\nu}^{1} d\omega\, (2\omega - \nu)^{J-n-1} \frac{1 + \omega^2 - \nu(1+\omega)}{1-\omega} \right) . \quad (4.49)$$

The ω integration is divergent at $\omega = 1$. This divergence is cancelled by the self-energy graphs, the value of which in LC gauge can be readily taken from [47] as

$$Z_F(x) = 1 + \frac{\alpha_s C_F}{2\pi} \frac{2}{\epsilon} (-2I_0 - 2\log|x| + \frac{3}{2}) , \quad (4.50)$$

where I_0 is the colinear divergence

$$I_0 = \int_0^1 \frac{dz}{z} = \int_0^1 \frac{dz}{1-z} . \quad (4.51)$$

This divergence can be viewed as essentially an artifact coming from the choice of using light cone calculations. For a detailed treatment please refer to [47]. Here we simply state and use their conclusions.

Z_F depends on the longitudinal momentum fraction x. So in our case the self-energy contribution from the $k_1 - r$ line should have $x = \frac{n\cdot(k_1 - r)}{n\cdot k_1} = 1 - \nu$ while the k_1 line has simply $x = 1$. Adding $\frac{1}{2}(Z_F(1) + Z_F(1-\nu))$ to (4.49) and identifying the logarithmic divergence $\int \frac{d\underline{k}^2}{\underline{k}^2} = \frac{2}{\epsilon}$ we will obtain the expression of γ_{qq} as in (4.53), namely,

$$\gamma_{qq}^{(+)} = \frac{\alpha_s}{2\pi} C_F\, \gamma_+ k_{1+}^{J-1} \nu^n \int_0^1 d\omega\, (2\omega - \nu)^{J-n-1} \Big(\Theta(\nu - \omega) \frac{1}{\nu}\, \omega\, (1 + \frac{\nu}{1-\omega}) \\ + \Theta(\omega - \nu) \frac{1}{1-\nu} \frac{1 + \omega^2 - \nu(1+\omega)}{1-\omega} + \delta(1-\omega)(-2I_0 - \log(1-\nu) + \frac{3}{2}) \Big) . \quad (4.52)$$

Note there are no other diagrams contributing in LC gauge. Recall this is because the gauge choice makes diagrams that have more than two lines meeting at the vertex vanish.

4.2.2 Anomalous Dimension Kernels

While the details of the computation of the other three anomalous dimensions are shown in the Appendices, we list the result in the following:

$$
\begin{aligned}
\gamma^{(+)}_{qq} &= \frac{\alpha_s}{2\pi} C_F\, \gamma_+ k_{1+}^{J-1} \nu^n \int_0^1 d\omega\, (2\omega - \nu)^{J-n-1} \Big(\Theta(\nu-\omega)\, \frac{1}{\nu}\, \omega\, (1 + \frac{\nu}{1-\omega}) \\
&\qquad + \Theta(\omega-\nu) \frac{1}{1-\nu} \frac{1+\omega^2-\nu(1+\omega)}{1-\omega} + \delta(1-\omega)(-2I_0 - \log(1-\nu) + \frac{3}{2}) \Big) \ , \\
\gamma^{(+)}_{gq} &= \frac{\alpha_s}{2\pi} C_F\, \gamma_+ k_{1+}^{J-1} \nu^n \int_0^1 d\omega (2\omega-\nu)^{J-n-2} \Big(\Theta(\nu-\omega) \frac{1}{\nu} (\omega^2-2\omega) \\
&\qquad - \Theta(\omega-\nu) \frac{1}{1-\nu} (1+(1-\omega)^2-\nu) \Big) \ , \\
\gamma^{ij}_{qg} &= \frac{\alpha_s}{2\pi} \frac{1}{2} 2 g^{ij} k_{1+}^{J} \nu^n \int_0^1 d\omega (2\omega-\nu)^{J-n-1} \Big(-\Theta(\nu-\omega) \frac{1}{\nu} \omega (2\omega-1-\nu) \\
&\qquad + \Theta(\omega-\nu) \frac{1}{1-\nu} (\omega^2+(1-\omega)^2-\omega\nu) \Big) \ , \\
\gamma^{ij}_{gg} &= \frac{\alpha_s C_A}{2\pi} 2 g^{ij} k_{1+}^{J} \nu^n \int_0^1 d\omega (2\omega-\nu)^{J-n-2} \Big(\Theta(\nu-\omega) \frac{1}{2} (\frac{1}{\nu} (4\omega^3-\omega^2+4\omega) \\
&\quad + (\frac{\omega^3+\omega^2-2\omega^2\nu}{1-\omega} - 3(\omega^2+\omega))) + \frac{\Theta(\omega-\nu)}{1-\nu} (\frac{2(1-\omega+\omega^2)^2}{1-\omega} + \nu \frac{\nu(1+\omega^2)-2(1+\omega^3)}{1-\omega}) \\
&\qquad + (1-\nu)\delta(1-\omega)(-2I_0 - \log(1-\nu) + \frac{b_0}{2C_A}) \Big) \ .
\end{aligned}
\tag{4.53}
$$

We have included virtual corrections coming from the self-energy graphs that will cancel the colinear singularities at end point in the ω integration (I_0 terms, see equation 4.51 for the definition of I_0) and give correct constant terms in the anomalous dimensions. Θ is the usual step function and

$$
b_0 = 11 - \frac{2}{3} n_f \tag{4.54}
$$

is the leading coefficient of the QCD β-function.

4.2.3 Comments on Anomalous Dimensions

Some comments are in order. It is straight forward to show that in the forward limit where $\nu = 0$, equation 4.53 reduces to the conventional Altarelli-Parisi splitting functions [13]. However, because of the non-forwardness, there is no simple probability interpretation for these evolution kernels as splitting functions.

The reason we have two terms for each triangle graph is that for different k_+ integration regions pinching of the k_- pole is different (see section 4.2.1). Thus there is no straight forward optical-theorem type dispersion relationship between the cross section and the imaginary part of the amplitude. Nonetheless, we can still do analytic continuation in J and relate the matrix elements to "double parton distributions" [12] which now may not have a direct probability interpretation. The second terms of our γ differ from those of [20] only because the factor $n \cdot k \, n \cdot (k - r)$ in our convention for the gluon vertex (see equation 4.23) is not included in the definition of their gluon vertex. Once we take this into account and use the same vertices, their results are identical to the second terms of the corresponding kernels of ours. [1]

After we perform the ω (i.e. k_+) integral, we can in principle obtain the corresponding anomalous dimension matrix in the moment space. At first sight, the $(1-\nu)^{-1}$ factor in the second terms may generate a series of infinite sums over powers of ν, which will spoil the locality of the vertex thus invalidate the operator product expansion. Detailed calculations show that because the lower bound of integration is now ν instead of zero, there will also always be at least one power of $(1-\nu)$ coming out from the integral. Furthermore, the highest powers of ν cancel completely between the first two terms of each γ, leaving the highest surviving ν powers to be ν^{J-1} for the quark sector and ν^J for the gluonic sector, as they must be to make the OPE valid. All the I_0 and $\log(1-\nu)$ terms that might potentially spoil the operator product expansion also cancel completely between each triangle graph and its corresponding self-energy graphs.

---

[1] Simultaneous to our work, Blumlein *et al* [48] also calculated the evolution kernels of what they call "twist 2 light-ray operators for unpolarized and polarized DIS". It would be useful to make a detailed comparison to see whether the seeming deviation between our results is a nontrivial disagreement or mainly due to different notations and approaches. Because of these differences in notation and approaches, together with the fact that their main results *eqs.*(15) – (18) were stated but not derived, we are not able to make such a comparison at the present time.

Because of the mixing among different n moments, as well as between quark and gluon sectors, it is very difficult to read off the anomalous dimensions of the mixing between two operators (with same J) of definite n moments. [2] However, the form of the anomalous dimension matrix simplifies greatly when the high energy limit is taken, and we will be able to make connection between this general formalism and the conventional leading logarithmic approximation (LLA) analysis results.

4.3 Lowest Order Wilson Coefficients

The lowest order Wilson coefficients can be calculated in perturbation theory from tree level Born diagrams as shown in figure 18, where we have explicitly included the cross diagram.

Using standard Feynman rules we can write the value of the Born diagrams, after spin average, as

---

[2] After the completion of this work, through private communications with Professors X. Ji and A. V. Radyushkin, the author learned that by adopting a basis of linear combinations of the operators in 3.11 using Gegenbauer polynomials as coefficients we might be able to explicitly diagonalize the anomalous dimension matrix in the moment space.

$$\begin{aligned} \mathrm{T}^{(0)}_{\mu\nu} &= \frac{1}{2}\sum_s (ie_q)^2 \overline{U}_s(p') \left(\gamma_\mu \frac{i(\not{p} + \not{q}')}{(p+q')^2 + i\epsilon}\gamma_\nu \right. \\ &\quad \left. + \gamma_\nu \frac{i(\not{p} - \not{q})}{(p-q)^2 + i\epsilon}\gamma_\mu \right) U_s(p)\,. \end{aligned} \tag{4.55}$$

By using equations 2.10, 4.10 and 4.13 we find

$$\begin{aligned} \mathrm{T}^{(0)}_{\mu\nu} &= -ie_q^2 \frac{\sqrt{1-\zeta}}{2\overline{q}^2} \left(\frac{1}{1-\omega} \sum_s \overline{U}_s(p)\gamma_\mu(\not{p} + \not{q}')\gamma_\nu U_s(p) \right. \\ &\quad \left. + \frac{1}{1+\omega} \sum_s \overline{U}_s(p)\gamma_\nu(\not{p} - \not{q})\gamma_\mu U_s(p) \right) \\ &= -ie_q^2 \frac{\sqrt{1-\zeta}}{2\overline{q}^2} \sum_{l=0}^{\infty} \omega^l (Tr(\not{p}\gamma_\mu(\not{p} + \not{q}')\gamma_\nu) + (-1)^l Tr(\not{p}\gamma_\nu(\not{p} - \not{q})\gamma_\mu)) \\ &= -2ie_q^2 \frac{\sqrt{1-\zeta}}{\overline{q}^2} \sum_{l=0}^{\infty} \omega^l (p_\mu(p+q')_\nu + p_\nu(p+q')_\mu - g_{\mu\nu} p\cdot(p+q')) \\ &\quad + (-1)^l (p_\mu(p-q)_\nu + p_\nu(p-q)_\mu - g_{\mu\nu} p\cdot(p-q)) \\ &= -2ie_q^2 \frac{\sqrt{1-\zeta}}{\overline{q}^2} \sum_{l=0}^{\infty} \omega^l ((p_\mu p_\nu + p_\nu p_\mu + p_\mu q'_\nu + p_\nu q'_\mu - g_\mu p\cdot q) \\ &\quad + (-1)^l (p_\mu p_\nu + p_\nu p_\mu - p_\mu q_\nu - p_\nu q_\mu + g_{\mu\nu} p\cdot q)) \\ &= -2ie_q^2 \frac{\sqrt{1-\zeta}}{\overline{q}^2} \sum_{l=0}^{\infty} \omega^{2l} (2(p_\mu p_\nu + p_\nu p_\mu) - p_\mu r_\nu - r_\mu p_\nu \\ &\quad + \omega(p_\mu(q'+q)_\nu + (q'+q)_\mu p_\nu - 2g_{\mu\nu} p\cdot q)) \\ &\equiv -2ie_q^2 \frac{\sqrt{1-\zeta}}{\overline{Q}^2} \sum_{l=0}^{\infty} \omega^{2l} C_{\mu\nu}\,, \end{aligned} \tag{4.56}$$

where

$$C_{\mu\nu} = (2p-r)_\mu r_\nu + r_\mu (2p-r)_\nu + 2\omega(p_\mu \overline{q}_\nu + \overline{q}_\mu p_\nu - g_{\mu\nu} p\cdot q)\,. \tag{4.57}$$

By similar derivations that lead to equation (3.31), as well as the definition of the moment variables (see equation 3.13) we find

$$\begin{aligned}
C_{\mu\nu} &= 2(2-\zeta)p_\mu p_\nu - 2(2-\zeta)\frac{p\cdot q}{\overline{q}^2}(p_\mu \overline{q}_\nu + \overline{q}_\mu p_\nu - g_{\mu\nu} p\cdot q) \\
&= 2(2-\zeta)(p_\mu p_\nu - \frac{p\cdot q}{\overline{q}^2}(p_\mu q_\nu + q'_\mu p_\nu) + \frac{(p\cdot q)^2}{\overline{q}^2} g_{\mu\nu}) \\
&= 2(2-\zeta)\left(p_\mu p_\nu - \frac{p\cdot q}{\overline{q}^2}(p_\mu q_\nu + q'_\mu p_\nu) + (\frac{p\cdot q}{\overline{q}^2})^2 q'_\mu q_\nu + \frac{(p\cdot q)^2}{\overline{q}^2}(g_{\mu\nu} - \frac{q'_\mu q_\nu}{\overline{q}^2})\right) \\
&= 2\omega\left(\frac{\overline{Q}^2}{p\cdot q}(p_\mu p_\nu - \frac{p\cdot q}{\overline{q}^2}(p_\mu q_\nu + q'_\mu p_\nu) + (\frac{p\cdot q}{\overline{q}^2})^2 q'_\mu q_\nu) + p\cdot q(-g_{\mu\nu} + \frac{q'_\mu q_\nu}{\overline{q}^2})\right) .
\end{aligned} \tag{4.58}$$

Thus we obtain the value of the Born diagrams as

$$\begin{aligned}
\mathrm{T}^{(0)}_{\mu\nu} &= -4ie_q^2\sqrt{1-\zeta}\sum_{l=0}^{\infty}\omega^{2l+1}\left(\frac{p\cdot q}{\overline{Q}^2}(-g_{\mu\nu} + \frac{q'_\mu q_\nu}{\overline{q}^2})\right. \\
&\left. + \frac{1}{p\cdot q}\left(p_\mu p_\nu - \frac{p\cdot q}{\overline{q}^2}(p_\mu q_\nu + q'_\mu p_\nu) + (\frac{p\cdot q}{\overline{q}^2})^2 q'_\mu q_\nu\right)\right) .
\end{aligned} \tag{4.59}$$

Comparing to (3.35) we obtain

$$\begin{aligned}
\tilde{E}^{(q)(0)}_{J,n=0}(\alpha_s(\overline{Q}),1) &= 4ie_q^2 , \qquad J \ odd,\ J \geq 3 , \\
\tilde{E}^{(q)(0)}_{J,n}(\alpha_s(\overline{Q}),1) &= 0 , \qquad J, n \ otherwise .
\end{aligned} \tag{4.60}$$

Thus for each J value, only the Wilson coefficient of $n = 0$ has a non-zero value at the tree level. This means that at leading order there is no dependence on the non-forwardness in the Wilson coefficients.

Chapter 5

High Energy Limit

In this chapter we will solve the renormalization group (RG) equation for the Wilson coefficient (4.18) in the high energy limit, and discuss the relationship between our general results and that of the usual forward double leading logarithmic approximation (DLLA) [17].

5.1 Reduction of the Path Ordered Exponential

Since we have computed the lowest order Wilson coefficients (4.60), and the reduced matrix elements are regarded as input from experimental data, we need now to evaluate the path ordered exponential $\mathrm{M}^{i'i}_{n'n}(\alpha(\overline{Q}), \frac{\overline{Q}}{\mu})$ in equation (4.19) in order to obtain the double moments $\mathrm{T}^{(J,n)}_{2,(\overline{Q},\mu_0)}$ and then the invariant amplitude T_2.

Recall that formally we have (4.20)

$$\mathrm{M}^{i'i}_{n'n}(\alpha_s(\overline{Q}), \frac{\overline{Q}}{\mu}) = \left(\mathcal{P}\exp(-\int_{\mu^2}^{\bar{Q}^2} \tilde{\gamma}(\alpha_s(\lambda^2))\frac{d\lambda^2}{\lambda^2})\right)^{i'i}_{n'n}, \tag{5.1}$$

thus to explicitly solve the RG equation we need to diagonalize the anomalous dimension matrix in both flavor (i) and moment (n) indices. However, in the high energy limit the situation simplifies.

Similar to the forward case, after analytical continuation in J, at high energy, the dominant contributions to T_2 come from the leading (right most) poles in the J-plane, which, as we will see, means that the diagonal elements in the moment space (n space) of the anomalous dimension matrix dominate the evolution. And because the gluon anomalous dimension has the right most pole (one unit higher than that of the quark sector) (see equation 5.4), together with the fact that gluons have bigger color charge, in flavor space (i space) at high energy the gluon anomalous dimension dominates.

However, gluons only have color charge and thus can not interact directly with the color-neutral virtual photon. The scattering must happen via a quark loop, which means we have to force at least one gluon-quark transition at the end of the evolution. Thus in each term of the expansion of the path ordered exponential (4.20), all the factors of the product of anomalous dimensions are $\tilde{\gamma}^{gg}$ except the first one, which should be $\tilde{\gamma^{qg}}$ due to the quark-gluon transition.

Therefore we need only to evaluate M between a quark and a gluon state and thus collapse the two flavor sums in (4.21) to $i' = q$ and $i = g$. Formally, if we now label $\gamma^{ii'}$ as the anomalous dimension matrix in the moment space, by expanding the path ordered exponential, we have

$$
\begin{aligned}
\mathrm{M}^{i'i}_{n'n}(\alpha_s(\overline{Q}), \frac{\overline{Q}}{\mu}) \quad & \xrightarrow{i'=q\,,i=g} \quad \left(\mathcal{P}\exp(-\int_{\mu^2}^{\bar{Q}^2}\tilde{\gamma}(\alpha_s(\lambda^2))\frac{d\lambda^2}{\lambda^2})\right)^{qg}_{n'n} \\
& = \quad \left(\sum_{l=0}^{\infty}\frac{1}{l!}\mathcal{P}\int_{\mu^2}^{\bar{Q}^2}\prod_{k=l}^{1}(\frac{d\lambda_k^2}{\lambda_k^2}(-\tilde{\gamma}(\alpha_s(\lambda_k^2))))\right)^{qg}_{n'n}
\end{aligned}
$$

$$
\begin{aligned}
&= \quad \Bigg(\sum_{l=0}^{\infty}\int_{\mu^2}^{\bar{Q}^2}\frac{d\lambda_l^2}{\lambda_l^2}(-\tilde{\gamma}(\alpha_s(\lambda_l^2)))\int_{\mu^2}^{\lambda_l^2}\frac{d\lambda_{l-1}^2}{\lambda_{l-1}^2}(-\tilde{\gamma}(\alpha_s(\lambda_{l-1}^2)))\cdot ...\\
&\qquad\qquad \cdot\int_{\mu^2}^{\lambda_2^2}\frac{d\lambda_1^2}{\lambda_1^2}(-\tilde{\gamma}(\alpha_s(\lambda_1^2)))\Bigg)^{qg}_{n'n}\\
&\xrightarrow{hi-E} \Bigg(\sum_{l=0}^{\infty}\int_{\mu^2}^{\bar{Q}^2}\frac{d\lambda_l^2}{\lambda_l^2}(-\tilde{\gamma}^{qg}(\alpha_s(\lambda_l^2)))\int_{\mu^2}^{\lambda_l^2}\frac{d\lambda_{l-1}^2}{\lambda_{l-1}^2}(-\tilde{\gamma}^{gg}(\alpha_s(\lambda_{l-1}^2)))\cdot ...\\
&\qquad\qquad \cdot\int_{\mu^2}^{\lambda_2^2}\frac{d\lambda_1^2}{\lambda_1^2}(-\tilde{\gamma}^{gg}(\alpha_s(\lambda_1^2)))\Bigg)_{n'n}\\
&\equiv \quad \Bigg(-\int_{\mu_0^2}^{\bar{Q}^2}\frac{d\lambda^2}{\lambda^2}\gamma^{qg}(\alpha_s(\lambda^2))\mathcal{P}\exp(-\int_{\mu_0^2}^{\lambda^2}\frac{d\lambda'^2}{\lambda'^2}\gamma^{gg}(\alpha_s(\lambda'^2)))\Bigg)_{n'n}\,,
\end{aligned}
\tag{5.2}
$$

which can be rewritten as

$$
\mathrm{M}^{qg}_{n'n}(\alpha_s(\overline{Q}),\frac{\overline{Q}}{\mu_0}) = \Bigg(-\int_{\mu_0^2}^{\bar{Q}^2}\frac{d\lambda^2}{\lambda^2}\gamma^{qg}(\alpha_s(\lambda^2))\Bigg)_{n'l}\Bigg(\mathcal{P}\exp(-\int_{\mu_0^2}^{\lambda^2}\frac{d\lambda'^2}{\lambda'^2}\gamma^{gg}(\alpha_s(\lambda'^2)))\Bigg)_{ln}\,. \tag{5.3}
$$

The leading pole terms of the relevant anomalous dimensions can be parametrized as (suppressing the J label), after integration of (4.53),

$$
\gamma^{qg}_{nn'}(\lambda^2) \;=\; -\frac{n_f}{2\pi}\alpha_s(\lambda^2)\;\overline{\gamma}^{qg}_{nn'}\,, \qquad \text{with} \quad \overline{\gamma}^{qg}_{nn'} = \frac{1+c^{qg}_{n',n}}{J-n}\;; \tag{5.4}
$$

$$
\gamma^{gg}_{nn'}(\lambda^2) \;=\; -\frac{C_A}{\pi}\alpha_s(\lambda^2)\;\overline{\gamma}^{gg}_{nn'}\,, \qquad \text{with} \quad \overline{\gamma}^{gg}_{nn'} = \frac{1+c^{gg}_{n',n}}{J-n-1}\;. \tag{5.5}
$$

where $0 \leq n \leq n' \leq J$ for $\overline{\gamma}_{nn'}$ and more importantly

$$
c^{i'i}_{n'=n,n} = 0\,. \tag{5.6}
$$

This means that only the diagonal elements have leading pole contributions.

Because of the upper triangular structure and the leading pole positions (recall that the right most pole dominates) of these $\overline{\gamma}$ matrices, we have, for $m \leq l \leq n$,

$$
\sum_l \overline{\gamma}_{ml}\,\overline{\gamma}_{ln} = \overline{\gamma}_{mn}\,\overline{\gamma}_{nn} \equiv \overline{\gamma}_{mn}\,a_n\,, \tag{5.7}
$$

where we have labeled the diagonal terms of the anomalous dimensions (leading pole terms) a_n. This means in a product of $\overline{\gamma}$ matrices we can, in leading logarithmic approximation (LLA), take only diagonal entries in all the factors except for the first one (see equation 5.11).

On the other hand, because we have the QCD running coupling (see, eg, [2])

$$\alpha(\lambda^2) = \frac{4\pi}{b_0 \log(\lambda^2/\Lambda^2)} \,, \tag{5.8}$$

where $\Lambda \equiv \Lambda_{QCD}$ is the QCD scale and b_0 the leading QCD β-function coefficient (see (4.54)), the integration of the anomalous dimension at high energy gives

$$\begin{aligned} -\int_{\mu_0^2}^{\lambda^2} \frac{d\lambda'^2}{\lambda'^2}\gamma^{gg}(\alpha_s(\lambda'^2))) &= \frac{C_A}{\pi}\overline{\gamma}^{gg}\int_{\mu_0^2}^{\lambda^2} \frac{d\lambda'^2}{\lambda'^2}\alpha(\lambda'^2) \\ &= \frac{4C_A}{\pi}\overline{\gamma}^{gg}\int_{\mu_0^2}^{\lambda^2} \frac{d\log(\lambda'^2/\Lambda^2)}{\log(\lambda'^2/\Lambda^2)} \\ &\equiv A(\lambda^2)\overline{\gamma}^{gg} \,, \end{aligned} \tag{5.9}$$

where we define the usual double-log factor

$$A(\lambda^2) = A \equiv \frac{4C_A}{b_0}\log\left(\frac{\log(\lambda^2/\Lambda^2)}{\log(\mu_0^2/\Lambda^2)}\right) \,. \tag{5.10}$$

Thus the path ordered exponential of γ^{gg} (the second factor on the right hand side of 5.3) becomes

$$\begin{aligned} \left(\mathcal{P}\exp(-\int_{\mu_0^2}^{\lambda^2} \frac{d\lambda'^2}{\lambda'^2}\gamma^{gg}(\alpha_s(\lambda'^2)))\right)_{ln} &= \left(\mathcal{P}e^{A\lambda^2\overline{\gamma}^{gg}}\right)_{ln} \\ &= (1 + A\overline{\gamma} + \frac{1}{2!}A^2\overline{\gamma}\overline{\gamma} + ...)_{ln} \\ &= (1 + A\overline{\gamma}_{ln} + \frac{A^2}{2!}a_n\overline{\gamma}_{ln} + ...) \\ &= \mathrm{I} + (A + \frac{A^2}{2!}a_n + ...)\overline{\gamma}^{gg}_{ln} \\ &= \mathrm{I} + \frac{1}{a_n^g}(e^{A(\lambda^2)a_n^g} - 1)\ \overline{\gamma}^{gg}_{ln} \\ &= \frac{1}{a_n^g}e^{A(\lambda^2)a_n^g}\ \overline{\gamma}^{gg}_{ln}\ +\ \text{non}-\text{leading terms} \,. \end{aligned} \tag{5.11}$$

Therefore, (5.3) reduces to

$$\begin{aligned}
\mathrm{M}^{qg}_{n'n}(\alpha_s(\overline{Q}),\frac{\overline{Q}}{\mu_0}) &= \left(-\int_{\mu_0^2}^{\bar{Q}^2}\frac{d\lambda^2}{\lambda^2}\gamma^{qg}(\alpha_s(\lambda^2))\right)_{n'l}\frac{1}{a_n^g}e^{A(\lambda^2)a_n^g}\ \overline{\gamma}^{gg}_{ln} \\
&= \frac{n_f}{2\pi}\int_{\mu_0^2}^{\bar{Q}^2}\frac{d\lambda^2}{\lambda^2}\alpha_s(\lambda^2)\ e^{A(\lambda^2)a_n}\overline{\gamma}^{qg}_{n'm}\ \overline{\gamma}^{gg}_{ln}\frac{1}{a_n^g} \\
&= \frac{n_f}{2\pi}\int_{\mu_0^2}^{\bar{Q}^2}\frac{d\lambda^2}{\lambda^2}\alpha_s(\lambda^2)\ e^{A(\lambda^2)a_n}\ \overline{\gamma}^{qg}_{n'n}\,, \qquad (5.12)
\end{aligned}$$

where in the last step we have again used (5.7).

The n' summation in the expression for the double moments (4.21) also collapses in LLA to the diagonal element of $\overline{\gamma}_{qg}$, and we have the final expression of the double moments as

$$\begin{aligned}
\frac{p\cdot q}{M^2}\sqrt{1-\zeta}\mathrm{T}^{(J,n)}_{2,(\overline{Q},\mu_0)} &= -\sum_{n'=0}^{n}\sum_{i'i}\tilde{E}^{(i')}_{J-2,n'}(\alpha_s(\overline{Q}),1)\ \mathrm{M}^{i'i}_{n'n}(\alpha_s(\overline{Q}),\frac{\overline{Q}}{\mu_0})\ \langle p'||\hat{O}^{(i)(J,n)}||p\rangle_{(\mu_0)} \\
&= -\sum_{n'=0}^{n}\tilde{E}^{(q)}_{J-2,n'}(\alpha_s(\overline{Q}),1)\ \mathrm{M}^{qg}_{n'n}(\alpha_s(\overline{Q}),\frac{\overline{Q}}{\mu_0})\ \langle p'||\hat{O}^{(g)(J,n)}||p\rangle_{(\mu_0)} \\
&= -\sum_{n'=0}^{n}\tilde{E}^{(q)}_{J-2,n'}(\alpha_s(\overline{Q}),1)\ \frac{n_f}{2\pi}\int_{\mu_0^2}^{\bar{Q}^2}\frac{d\lambda^2}{\lambda^2}\alpha_s(\lambda^2)\ e^{A(\lambda^2)a_n} \\
&\qquad\cdot\ \overline{\gamma}^{qg}_{n'n}\ \langle p'||\hat{O}^{(g)(J,n)}||p\rangle_{(\mu_0)} \\
&= -\tilde{E}^{(q)}_{J-2,n}(\alpha_s(\overline{Q}),1)\ \frac{n_f}{2\pi}\int_{\mu_0^2}^{\bar{Q}^2}\frac{d\lambda^2}{\lambda^2}\alpha_s(\lambda^2)\ e^{A(\lambda^2)a_n} \\
&\qquad\cdot\ \overline{\gamma}^{qg}_{nn}\ \langle p'||\hat{O}^{(g)(J,n)}||p\rangle_{(\mu_0)} \\
&= -\tilde{E}^{(q)}_{J-2,n}(\alpha_s(\overline{Q}),1)\ \langle p'||\hat{O}^{(g)(J,n)}||p\rangle_{(\mu_0)} \\
&\qquad\cdot\ \frac{n_f}{2\pi}\int_{\mu_0^2}^{\bar{Q}^2}\frac{d\lambda^2}{\lambda^2}\alpha_s(\lambda^2)\ e^{\frac{A(\lambda^2)}{J-n-1}}\frac{1}{J-n}\,. \qquad (5.13)
\end{aligned}$$

Now the invariant amplitude T_2 (4.16), after analytically continuation in J, becomes

$$\begin{aligned}
-\frac{p\cdot q}{M^2}\sqrt{1-\zeta}\mathrm{T}_2(\overline{Q}^2,\omega,\nu) = \int\frac{dJ}{2\pi i}\sum_{n=0}^{\infty}\nu^n\int_{\mu_0^2}^{\bar{Q}^2}\frac{d\lambda^2}{\lambda^2}\ \alpha_s(\lambda^2)\frac{n_f}{2\pi}\ \tilde{E}^{(q)}_{J-2,n}(\alpha_s(\overline{Q}),1) \\
\cdot\ \langle p'||\hat{O}^{(g)(J,n)}||p\rangle_{(\mu_0)}\frac{1}{J-n}\exp\left((J-n)\log\omega+\frac{A(\lambda^2)}{J-n-1}\right)\,. \qquad (5.14)
\end{aligned}$$

In the high energy limit we can make a saddle point approximation for the J integration because the structure of the exponent in the expression. Define

$$f(J) = (J-n)\log\omega + \frac{A}{J-n-1}\,, \tag{5.15}$$

we obtain the first and second order derivative of f as

$$\begin{aligned} f'(J) &= \log\omega - \frac{A}{(J-n-1)^2} \\ f''(J) &= \frac{2A}{(J-n-1)^3}\,. \end{aligned} \tag{5.16}$$

Setting $f'(J) = 0$ we obtain the saddle point J_s as

$$J_s = 1 + n + \sqrt{\frac{A}{\log\omega}}\,, \tag{5.17}$$

and at the saddle point we have

$$\begin{aligned} f(J_s) &= \log\omega + 2\sqrt{A\log\omega} \\ f''(J_s) &= \frac{2\log^{3/2}\omega}{A^{1/2}}\,, \end{aligned} \tag{5.18}$$

where indeed we can see that at J_s we have $f'' > 0$. Now the invariant amplitude becomes

$$\begin{aligned} -\frac{p\cdot q}{M^2}\sqrt{1-\zeta}\,\mathrm{T}_2 = {} & \sum_{n=0}^{\infty}\nu^n \int_{\mu_0^2}^{\bar{Q}^2}\frac{d\lambda^2}{\lambda^2}\,\alpha_s(\lambda^2)\frac{n_f}{2\pi}\;\tilde{E}^{(q)}_{J-2,n}(\alpha_s(\overline{Q}),1) \\ & \cdot\,\langle p'||\hat{O}^{(g)(J,n)}||p\rangle_{(\mu_0)}\frac{1}{1+\sqrt{\frac{A(\lambda^2)}{\log\omega}}}\sqrt{\frac{\pi A(\lambda^2)^{1/2}}{\log^{3/2}\omega}}\,e^{\log\omega}e^{2\sqrt{A(\lambda^2)\log\omega}} \\ = {} & \frac{n_f}{2\pi}\sqrt{\frac{\pi}{\log^{3/2}\omega}}\,\omega\int_{\mu_0^2}^{\bar{Q}^2}\frac{d\lambda^2}{\lambda^2}\alpha_s(\lambda^2)A(\lambda^2)^{1/4}\frac{1}{1+\sqrt{\frac{A(\lambda^2)}{\log\omega}}} \\ & \cdot e^{2\sqrt{A(\lambda^2)\log\omega}}\;\sum_{n=0}^{\infty}\nu^n\,\tilde{E}^{(q)}_{J-2,n}(\alpha_s(\overline{Q}),1)\langle p'||\hat{O}^{(g)(J,n)}||p\rangle_{(\mu_0)}\,. \end{aligned} \tag{5.19}$$

Because the saddle point fixes the value $J-n=J_s-n=1+\sqrt{\frac{A}{\log\omega}}$, which fixes the number of internal derivatives inside the gluon operator, recall (3.18), it is clear that the reduced matrix element in (5.19) is the conventional (forward) gluon distribution function in moment space with a shifted moment label J_s-n. Despite the formal summation over n, which is essentially the only difference introduced in this limit by the non-forwardness, only one non-perturbative input is needed, which is the gluon density. This means that although in general there will be complicated dependence on ν in the double distributions, (in high energy and hard scattering limit) it is nontheless perturbative.

At leading order the n summation collapses when we take the lowest order Wilson coefficients given in (4.60) with the flavor averaged quark charge $n_f<e_q^2>=\sum_f e_q^2$, which actually could be done even before taking the saddle point approximation. We went to the saddle point first in order to make manifest the fact that there would be only one soft input, the forward gluon density, even with the formal n summation. Thus, subsitute the result from equation 4.60 we have

$$-\frac{p\cdot q}{M^2}\sqrt{1-\zeta}\,\mathrm{T}_2(\overline{Q}^2,\omega,\nu)=\frac{2in_f<e_q^2>}{\pi}\sqrt{\frac{\pi}{\log^{3/2}\omega}}\,\omega\,\langle p'||\hat{O}^{(g)(J_s-n,0)}||p\rangle_{(\mu_0)} \cdot\int_{\mu_0^2}^{\bar{Q}^2}\frac{d\lambda^2}{\lambda^2}\alpha_s(\lambda^2)A(\lambda^2)^{1/4}\frac{1}{1+\sqrt{\frac{A(\lambda^2)}{\log\omega}}}\,e^{2\sqrt{A(\lambda^2)\log\omega}}. \quad (5.20)$$

Because of the pinching of J_s-n, the final evaluation of (5.20) is the same in both forward and non-forward cases, in the high energy limit under leading logarithmic approximation. Define the integral

$$I=\int_{\mu_0^2}^{\bar{Q}^2}\frac{d\lambda^2}{\lambda^2}\alpha_s(\lambda^2)A(\lambda^2)^{1/4}\frac{1}{1+\sqrt{\frac{A(\lambda^2)}{\log\omega}}}\,e^{2\sqrt{A(\lambda^2)\log\omega}}, \quad (5.21)$$

with $A(\lambda^2)$ given in equation 5.10, which is not very big even for very hard processes due to the double logarithm. Therefore in the high energy limit where ω is very big, we have

$$\frac{1}{1+\sqrt{\frac{A(\lambda^2)}{\log\omega}}} \to 1\,. \tag{5.22}$$

We make the change of variable

$$\log(\lambda^2/\Lambda^2) = \log(\overline{Q}^2/\Lambda^2) - u\,, \tag{5.23}$$

which implies that

$$u = \log(\overline{Q}^2/\Lambda^2) - \log(\lambda^2/\Lambda^2) = \log(\overline{Q}^2/\lambda^2)\,. \tag{5.24}$$

It is clear that $u > 0$ and is not very large either. To simplify the notation, let

$$\begin{aligned} a &= \frac{4C_A}{b_0} \\ b &= \log(\overline{Q}^2/\Lambda^2) \\ c &= \log\left(\frac{\log(\overline{Q}^2/\Lambda^2)}{\log(\mu_0^2/\Lambda^2)}\right)\,, \end{aligned}$$

where we make the note that a and b can be seen as moderate compared with the double logarithms u and c, which are not big. Then

$$\begin{aligned} A(\lambda^2) &= \log\left(\frac{\log(\lambda^2/\Lambda^2)}{\log(\mu_0^2/\Lambda^2)}\right) = \log\left(\frac{\log(\overline{Q}^2/\Lambda^2)}{\log(\mu_0^2/\Lambda^2)}\,(1-\frac{u}{\log(\overline{Q}^2/\Lambda^2)})\right) \\ &= c + \log(1-\frac{u}{b}) \\ &= c - \frac{u}{b} = c\left(1-\frac{u}{bc}\right)\,. \end{aligned} \tag{5.25}$$

And the integral I becomes

$$\begin{aligned} I &= \int_0^{\log\frac{\overline{Q}^2}{\mu_0^2}} \frac{du}{b_0(b-u)}(ac)^{1/4}(1-\frac{u}{4bc})\,e^{2\sqrt{\log\omega\,ac\,(1-\frac{u}{bc})}} \\ &= \int_0^{\log\frac{\overline{Q}^2}{\mu_0^2}} \frac{du}{b_0(b-u)}(ac)^{1/4}(1-\frac{u}{4bc})\,e^{2\sqrt{\log\omega\,ac}(1-\frac{u}{2bc})} \\ &\equiv \frac{(ac)^{1/4}}{b_0}e^{2\sqrt{ac\log\omega}}\cdot II\,, \end{aligned} \tag{5.26}$$

where the second integral II is defined as

$$II = \int_0^{\log \frac{\overline{Q}^2}{\mu_0^2}} du \frac{1}{b-u}(1 - \frac{u}{4bc})\, e^{-\frac{u}{b}\sqrt{\frac{\log \omega a}{c}}} \,. \tag{5.27}$$

As we have discussed earlier, because the relative size of a and c and that $\log \omega$ is considered big, $\sqrt{\frac{\log \omega a}{c}}$ is actually a big number.

Rewrite (5.27) as

$$\begin{aligned} II &= \frac{1}{b}\int_0^{\log \frac{\overline{Q}^2}{\mu_0^2}} du \frac{1}{1-\frac{u}{b}}(\frac{1}{4c}(1-\frac{u}{b}) + 1 - \frac{1}{4c})e^{-\frac{u}{b}\sqrt{\frac{\log \omega a}{c}}} \\ &= \frac{1}{b}\int_0^{\log \frac{\overline{Q}^2}{\mu_0^2}} du(\frac{1}{4c} + \frac{1}{1-\frac{u}{b}}(1-\frac{1}{4c}))e^{-\frac{u}{b}\sqrt{\frac{\log \omega a}{c}}} \,, \end{aligned} \tag{5.28}$$

and let

$$z = \frac{u}{b} = \frac{\log(\overline{Q}^2/\lambda^2)}{\log(\overline{Q}^2/\Lambda^2)} \,, \tag{5.29}$$

we have

$$II = \int_0^{\frac{\log \overline{Q}^2/\mu_0^2}{b}} dz \left(\frac{1}{4c} e^{-z\sqrt{\log \omega a/c}} + (1 - \frac{1}{4c}) \frac{e^{-z\sqrt{\log \omega a/c}}}{1-z} \right) . \tag{5.30}$$

The z integral can be evaluated using the formula of special integral exponential function $IE_i(z)$

$$\int_0^{\alpha} \frac{dz}{1-z} e^{-\lambda z} = e^{-\lambda}(IE_i(\lambda) - IE_i(\lambda - \alpha\lambda)) \,, \tag{5.31}$$

where the $IE_i(z)$ function has a branch cut discontinuity at the negative real axis in the z-plane (i.e., cut at $z \in (-\infty, 0]$), and has the following properties:

$$\begin{aligned} IE_i(0) &= -\infty \,, \\ IE_i(0.3725) &= 0 \,, \end{aligned}$$

and while

$$\partial_z IE_i(z) = \frac{e^z}{z} \,, \tag{5.32}$$

we have

$$\frac{IE_i(z)}{e^z} << 1 \tag{5.33}$$

when $z \geq \mathrm{O}(1)$.

Therefore the second term in equation 5.30 does not contribute in the leading logarithmic approximation, and we have

$$\begin{aligned} II &= \frac{1}{4c}\frac{1}{\sqrt{\log \omega a/c}}\left(1 - e^{-\sqrt{\log \omega a/c \frac{\log \overline{Q}^2/\mu_0^2}{\log \overline{Q}^2/\Lambda^2}}}\right) \\ &= \frac{1}{4\sqrt{\log \omega ac}} \,, \end{aligned} \tag{5.34}$$

where in the last step we have used the fact that $\sqrt{\log \omega a/c}$ is a big number.

Subsititute these results into the expression of the invariant amplitude (5.20) we have

$$\begin{aligned} -\frac{p\cdot q}{M^2}\sqrt{1-\zeta}\,\mathrm{T}_2(\overline{Q}^2,\omega,\nu) &= \frac{2in_f < e_q^2 >}{\pi}\,\langle p'||\hat{O}^{(g)(J_s-n,0)}||p\rangle_{(\mu_0)} \\ &\quad \sqrt{\frac{\pi}{\log^{3/2}\omega}}\,\omega\frac{(ac)^{1/4}}{b_0}e^{2\sqrt{ac\log\omega}}\frac{1}{4\sqrt{\log \omega ac}} \,. \end{aligned} \tag{5.35}$$

Since $ac = \frac{4C_A}{b_0}A(\overline{Q}^2)$, the final expression of the double distribution T_2 is

$$\begin{aligned} \frac{p\cdot q}{M^2}\sqrt{1-\zeta}\,\mathrm{T}_2(\overline{Q}^2,\omega,\nu) = -\frac{in_f < e_q^2 >}{2\sqrt{\pi}b_0}\left(\frac{1}{\frac{4C_A}{b_0}\log^5 \omega A(\overline{Q}^2)}\right)^{\frac{1}{4}} \\ \cdot\,\langle p'||\hat{O}^{(g)(J_s-n,0)}||p\rangle_{(\mu_0)}\ \omega\ e^{2\sqrt{\frac{4C_A}{b_0}A(\bar{Q}^2)\log\omega}} \,. \end{aligned} \tag{5.36}$$

In the very high energy and hard scattering limit, the asymmetric reduced matrix element becomes the forward gluon density

$$\begin{aligned} \langle p'||\hat{O}^{(g)(J_s-n,0)}||p\rangle &\to \sqrt{1-\zeta}\,\langle p||\hat{O}^{(g)(J_s-n,0)}||p\rangle \\ &\propto \sqrt{1-\zeta}\,(xg(x))\,, \end{aligned} \tag{5.37}$$

with $x=\frac{1}{\omega}$. We can see clearly that the leading high energy (small-x) behavior is exactly the same as that is given by a forward direction leading logarithmic analysis (see section 1.5.3).

Chapter 6

Conclusions and Open Problems

We have formulated a general operator product expansion of a non-forward and unequal mass virtual Compton scattering amplitude. We have found that, because of the non zero momentum transfer, the expansion now should be done in double moments with respect to the moment variables defined in equation 3.13. The double moments can be parametrized as products of the Wilson coefficients, which can be computed perturbatively at the hard scattering scale, and the non-forward matrix elements of new sets of quark and gluon operators, namely, the double distribution functions. These operators in general have total derivatives and they mix among themselves under renormalization. They obey a well-defined renormalization group equation that can be solved formally. We have calculated the (equivalent to) evolution kernels of the double distribution from which the anomalous dimension matrix of these operators can be extracted.

In the high energy limit, we have found the leading contributions of the anomalous dimension and used them to solve the resulting renormalization group equation. We have recovered the conventional double leading logarithmic (DLL) analysis results and have made the connection that in fact the double distributions are proportional to the conventional forward gluon density in this limit. This justifies previous analyses using DLL approximation and forward gluon density on non-forward processes like the exclusive diffractive vector meson production (e.g. [17]). Furthermore, we have developed the formalism to proceed in principle beyond DLL to next to leading order and/or relaxing the leading logarithmic approximations.

Our analysis is quite general. We can go from deeply inelastic scattering (DIS) (the forward case, corresponding to $\nu = 0$) to deeply virtual Compton scattering (DVCS) (corresponding to $\nu = 1$) in principle by continuously varying the moment variable ν. However, the analyticity/analytical continuation in ν, as well as the relevant dispersion relationships, still need further investigation. At the same time, diffractive vector meson production (ν close to 1) can also be studied by replacing the second quark-photon vertex with effectively a (light cone) wave function of the vector meson [17, 31]. Also, here because the time-like four-momentum of the vector meson, further study is needed in extending the discussion to the time-like region.

There are also other details of the analysis still need to be filled in. The explicit form of the anomalous dimension matrix is not yet written down, although as mentioned earlier, a change of base to a linear combination of the quark and gluon operators 3.11 using Gegenbauer polynomial as coefficient might simplify the situation. The dispersion relationship between the invariant amplitudes T_i and the double structure functions need also to be clarified. For example, the exact physical meaning of the fact that different k_- poles are being picked out at different k_+ integration regions. The relationship between this and the transition/interpolation between Altarelli-Parisi evolution and Brodsky-Lepage evolution is also an important topic to study in depth.

More experimental data are to come from DESY, Jefferson Lab, SLAC, CERN and RHIC, the study of the physics of non-forward processes will continue to be an interesting and important field.

Bibliography

[1] C. Itzykson and J. Zuber, *Quantum Field Theory*, McGraw-Hill International Book Co., 1980.

[2] M.E. Peskin and D.V. Schroeder, *Introduction to Quantum Field Theory*, Addison-Wesley Pub., 1995.

[3] G. Sterman, *An Introduction to Quantum Field Theory*, Cambridge University Press, 1993.

[4] T.P. Cheng and L.F. Li, *Gauge Theory of Elementary Particle Physics*, Oxford University Press, 1984.

[5] J.C. Collins, *Renormalization*, Cambridge University Press, 1984.

[6] M. Hamermesh, *Group Theory and Its Application to Physical Problems*, Dover Pub., 1989.

[7] W. Greiner and A. Schafer, *Quantum Chromodynamics*, Springer-Verlag, 1994.

[8] *Perturbative Quantum Chromodynamics*, A.H. Mueller, ed., World Scientific, Singapore, 1989.

[9] R.D. Field, *Applications of Perturbative QCD*, Addison-Wesley Pub., 1989.

[10] K. Wilson, Phys. Rev. D10 (1974) 2445., *New Phenomena in Subnuclear Physics*, A. Zichichi ed., Plenum Press, 1977.

[11] H. Abramowicz and A. Caldwell, *HERA Collider Physics*, DESY98-192, December 1998. To appear in Rev. Mod. Phys..

[12] N. Christ, B. Hasslacher, A.H. Mueller, Phys. Rev. D6 (1972) 3543.

[13] G. Altarelli and G. Parisi, Nucl. Phys. B126 (1977) 298.

[14] A.H. Mueller, Phys. Rev. D18 (1978) 3705.

[15] X. Ji, Phys. Rev. D55 (1997) 7114.

[16] A.V. Radyushkin, Phys. Lett. B380 (1996) 417.

[17] S.J. Brodsky, L. Frankfurt, J.F. Gunion, A.H. Mueller, M. Strikman, Phys. Rev. D50 (1994) 3134.

[18] A.V. Radyushkin, Phys. Lett. B385 (1996) 333.

[19] J.C. Collins, L. Frankfurt, M. Strikman, Phys. Rev. D56 (1997) 2982.

[20] L. Frankfurt, A. Freund, V. Guzey, M. Strikman, Phys. Lett. B418 (1998) 345.

[21] X. Ji, Phys. Rev. Lett. 78 (1997) 610.

[22] X. Ji, W. Melnitchouk, X. Song, Phys. Rev. D56 (1997) 5511.

[23] A.V. Radyushkin, Phys. Rev. D56 (1997) 5524.

[24] I.V. Musatov and A.V. Radyushkin, hep-ph 9905376.

[25] Z. Chen, Nucl. Phys. B525 (1998) 369.

[26] A.H. Compton, Phys. Rev. 22 (1923) 409.

[27] F. Low, Phys. Rev. 96 (1954) 1954.
M. Gell-Mann and M.L. Goldberger, Phys. Rev. 96 (1954) 1433.

[28] F.J. Federspiel, *et al*, Phys. Rev. Lett. 67 (1991) 1511.
A. Zieger, *et al*, Phys. Lett. B278 (1992) 34.
E. Hallin, *et al*, Phys. Rev. C48 (1993) 1497.

[29] B. Geyer, D. Robaschik, M. Bordag, J. Horejsi, Z. Phys. C26 (1985) 591.
T. Braunshweig, B. Geyer, J. Horejsi, D. Robaschik, Z. Phys. C33 (1987) 275.
F.M. Dittes, *et al* , Phys. Lett. B209 (1988) 325.

[30] D. Mueller, *et al* , Fortschr. Phys. 42 (1994) 101.

[31] G.P. Lepage and S.J. Brodsky, Phys. Lett. B87 (1979) 79.
G.P. Lepage and S.J. Brodsky, Phys. Rev. D22 (1980) 2157.

[32] P. Jain and J.P. Ralston, in *The Proceedings of the Workshop on Future Directions in Particle and Nuclear Physics at Multi-Gev Hadron Beam Facilities*, Brookhaven National Laboratory, 1993.

[33] J. Ashman, *et al*, Nucl. Phys. B32 (1989) 1.
B. Adeva, *et al* , Phys. Lett. B302 (1993) 533.
P.L. Anthony, *et al* , Phys. Rev. Lett. 71 (1993) 959.
K. Abe, *et al* , Phys. Rev. Lett. 75 (1995) 25.

[34] A. Chodos, R.L. Jaffe, K. Johnson, C.B. Thorn, V. Weisskopf, Phys. Rev. D9 (1974) 3471.

[35] Y.Y. Balitsky and L.N. Lipatov, Sov. J. Nuc. Phys. 28 (1978) 822.
E.A. Kuraev, L.N. Lipatov, V.S. Fadin, Sov. Phys. JETP 45 (1977) 199.
L.N. Lipatov, Sov. Phys. JETP 63 (1986) 904.

[36] M.G. Ryskin, Z. Phys. C57 (1993) 89.

[37] L. Frankfurt, W. Koepf, M. Strikman, Phys. Rev. D54 (1996) 3194.

[38] M. Derrick, *et al*, Phys. Lett. B356 (1995) 601.
T. Ahmed, *et al*, Phys. Lett. B338 (1994) 507.
M. Derrick, *et al*, Phys. Lett. B350 (1995) 120.

[39] S. Catani, M. Ciafaloni, F. Hautmann, Phys. Lett. B242 (1990) 97.
S. Catani, M. Ciafaloni, F. Hautmann, Nucl. Phys. B366 (1991) 135.
S. Catani, M. Ciafaloni, F. Hautmann, Phys. Lett. B307 (1993) 147.
S. Catani and F. Hautmann, Nucl. Phys. B427 (1994) 475.

[40] J.R. Forshaw and M.G. Ryskin, Z. Phys. C68 (1995) 589.
M.G. Ryskin, R.G. Roberts, A.D. Martin, E.M. Levin, Z. Phys. C76 (1997) 231.

[41] X. Ji and J. Osborne, Phys. Rev. D58 (1998) 094018.

[42] K. Watanabe, Prog. Theo. Phys. 67 (1982) 1834.

[43] A.V. Efremov and A.V. Radyushkin, Phys. Lett. B94 (1980) 245.
S. Brodsky, Y. Frishman, G. Lepage, C. Sachrajda, Phys. Lett. B91 (1980) 239.

[44] A.V. Belitsky and D. Mueller, Phys. Lett. B417 (1998) 129.
D. Mueller, Phys. Rev. D58 (1998) 054005.

[45] A.H. Mueller, *Notes for Particle Physics*, Columbia University Graduate Course G8070.

[46] J.J. Sakurai, *Modern Quantum Mechanics*, Addison-Wesley Pub., 1994.

[47] G. Curci, W. Furmanski, R. Petronzio, Nucl. Phys. B175 (1980) 27.

[48] J. Blumlein, B. Geyer, D. Robaschik, Phys. Lett. B406 (1997) 161.

Appendix A

Quark-Gluon Anomalous Dimension

In this appendix we show the details of the calculation that leads to our form of the quark-gluon anomalous dimension in equation 4.53.

In the LC gauge, the value of the quark-gluon diagram, shown in figure 20, is

$$\gamma_{qg}^{\alpha\beta} = (-ig)^2 \sum_{a,b} (T^i_{ab} T^{i'}_{ba}) \int \frac{d^4k}{(2\pi)^4} (-1) Tr \left(\gamma_\alpha \frac{i}{\not{k} - \not{r}} O_q^{(J,n)(+)}(k, k-r) \frac{i}{\not{k}} \gamma_\beta \frac{i}{\not{k}_1 - \not{k}} \right) , \tag{A.1}$$

where O_q is the quark vertex from (4.23) and there is an extra factor of (-1) coming from the fermion loop. After the color algebra we obtain

$$\gamma_{qg}^{\alpha\beta} = -ig^2 \frac{1}{2} \delta_{ii'} \int \frac{d^4k}{(2\pi)^4} \frac{(2k-r)_+^{J-n-1} r_+^n B_{\alpha\beta}(k,r)}{(k^2+i\epsilon)((k_1-k)^2+i\epsilon)((k-r)^2+i\epsilon)} , \tag{A.2}$$

where we have defined

$$B_{\alpha\beta}(k,r) = Tr(\gamma_\alpha (\not{k} - \not{r}) \gamma_+ \not{k} \gamma_\beta (\not{k}_1 - \not{k})) \ . \tag{A.3}$$

Almost identical to the calculation of the gluon-quark section, after the k_- integral we have

$$\begin{aligned} \gamma_{qg}^{\alpha\beta} &= -\frac{\alpha_s}{2\pi}\frac{1}{4}\int\frac{d\underline{k}^2}{\underline{k}^4}\left(\int_0^{r_+}\frac{dk_+k_+}{r_+k_{1+}}(2k-r)_+^{J-n-1}r_+^nB_{\alpha\beta}^{(1)}\right. \\ &\qquad \left.+\int_{r_+}^{k_{1+}}\frac{dk_+(k_1-k)_+}{k_{1+}(k_1-r)_+}(2k-r)_+^{J-n-1}r_+^nB_{\alpha\beta}^{(2)}\right) . \qquad \text{(A.4)} \end{aligned}$$

where $B^{(1)}$ and $B^{(2)}$ are $B_{\alpha\beta}(k,r)$ evaluated at the poles $k_-^{(1)}$ and $k_-^{(2)}$, respectively, and again with

$$k_-^{(1)} = \frac{\underline{k}^2}{2k_+} = \frac{\underline{k}^2}{2k_{1+}\omega} \qquad \& \quad k^2 = 2k_+k_- - \underline{k}^2 = 0,$$

$$k_-^{(2)} = \frac{\underline{k}^2}{2(k_+ - k_{1+})} = \frac{-\underline{k}^2}{2k_{1+}(1-\omega)} \qquad \& \quad k^2 = -\frac{\underline{k}^2}{(1-\omega)}.$$

We have

$$\begin{aligned} B_{\alpha\beta}(k,r) &= Tr(\gamma_\alpha(\not{k} - \not{r})\gamma_+ \not{k}\gamma_\beta(\not{k}_1 - \not{k})) \\ &= Tr(\gamma_\alpha \not{k}\gamma_+ \not{k}\gamma_\beta(\not{k}_1 - \not{k})) - Tr(\gamma_\alpha \not{r}\gamma_+ \not{k}\gamma_\beta(\not{k}_1 - \not{k})) \\ &\equiv B_{I,\alpha\beta} - B_{II,\alpha\beta} \ . \qquad \text{(A.5)} \end{aligned}$$

Same as in the main text, we will pick up only the terms that are logarithmic divergent in the transverse momentum ($\underline{k}^2$ terms) and thus we need only look at the transverse components ($\alpha\beta = i, j = 1, 2$ terms).

For the first term, we have

$$\begin{aligned} B_{I,\alpha\beta} &= Tr(\gamma_\alpha \not{k}\gamma_+ \not{k}\gamma_\beta(\not{k}_1 - \not{k})) \\ &= Tr(\gamma_\alpha(\gamma_+, \not{k} - \gamma_+ \not{k}) \not{k}\gamma_\beta(\not{k}_1 - \not{k})) \\ &= 2k_+Tr(\gamma_\alpha \not{k}\gamma_\beta(\not{k}_1 - \not{k})) - k^2Tr(\gamma_\alpha\gamma_+\gamma_\beta(\not{k}_1 - \not{k})) \\ &= 8k_+(k_\alpha(k_1-k)_\beta + k_\beta(k_1-k)_\alpha - g_{\alpha\beta}k\cdot(k_1-k)) \\ &\quad -k^2Tr(\gamma_-\gamma_+\gamma_-\gamma_+(k_1-k)_-) - k^2Tr(\underline{\gamma}\gamma_+\underline{\gamma}\gamma_-(k_1-k)_+)\, , \qquad \text{(A.6)} \end{aligned}$$

where we have used the identity of gamma matrices [2]

$$Tr(\not{A}\,\not{B}\,\not{C}\,\not{D}) = 4((A\cdot B)(C\cdot D) - (B\cdot D)(A\cdot C) + (A\cdot D)(B\cdot C))\,. \tag{A.7}$$

At the pole $k_-^{(1)}$ where $k^2 = 0$, we have, for the ij component,

$$\begin{aligned} B^{(1)}_{I,ij} &= 8k_+(-k_ik_j - k_jk_i - g_{ij}k\cdot k_1) \\ &= 8k_+(g_{ij}\underline{k}^2 - g_{ij}\frac{\underline{k}^2}{2\omega}) \\ &= 8k_+g_{ij}\underline{k}^2(1 - \frac{1}{2\omega}) \\ &= 4k_{1+}g_{ij}\underline{k}^2(2\omega - 1)\,, \end{aligned} \tag{A.8}$$

while at the pole $k_-^{(2)}$, where factors of transverse momentum $\underline{k}$ can come from either k_-, k^2 or k_ik_j terms, we have

$$\begin{aligned} B^{(2)}_{I,ij} &= 8k_+(-k_ik_j - k_jk_i - g_{ij}k\cdot(k_1 - k)) + 4k^2g_{ij}(k_1 - k)_+ \\ &= 8k_+(g_{ij}\underline{k}^2 - \frac{1}{2}g_{ij}\frac{\underline{k}^2}{1-\omega}) - \frac{4\underline{k}^2}{1-\omega}g_{ij}k_{1+}(1-\omega) \\ &= 4k_{1+}g_{ij}\underline{k}^2(2\omega - \frac{\omega}{1-\omega} - 1) \\ &= -4k_{1+}g_{ij}\underline{k}^2\frac{\omega^2 + (1-\omega)^2}{1-\omega}\,. \end{aligned} \tag{A.9}$$

On the other hand, for the second term we have, by using the anti-commutation relationships of the Dirac matrices (1.117),

$$\begin{aligned} B_{II,\alpha\beta} &= Tr(\gamma_\alpha\,\not{r}\gamma_+\,\not{k}\gamma_\beta(\not{k}_1 - \not{k})) \\ &= 2\gamma_\alpha Tr(\gamma_+\,\not{k}\gamma_\beta(\not{k}_1 - \not{k})) - Tr(\not{r}\gamma_\alpha\gamma_+\,\not{k}\gamma_\beta(\not{k}_1 - \not{k})) \\ &= 2\gamma_\alpha Tr(\gamma_+\,\not{k}\gamma_\beta(\not{k}_1 - \not{k})) - 2g_{\alpha+}Tr(\not{r}\,\not{k}\gamma_\beta(\not{k}_1 - \not{k})) \\ &\quad + Tr(\not{r}\gamma_+\gamma_\alpha\,\not{k}\gamma_\beta(\not{k}_1 - \not{k})) \\ &= \ldots \end{aligned}$$

$$\begin{aligned} &= 2\gamma_\alpha Tr(\gamma_+ \not k \gamma_\beta(\not k_1 - \not k)) - 2g_{\alpha+} Tr(\not r \not k \gamma_\beta(\not k_1 - \not k)) \\ &\quad +2k_\alpha Tr(\not r \gamma_+ \gamma_\beta(\not k_1 - \not k)) - 2g_{\alpha\beta} Tr(\not r \gamma_+ \not k(\not k_1 - \not k)) \\ &\quad +2(k_1 - k)_\alpha Tr(\not r \gamma_+ \not k \gamma_\beta) - Tr(\not r \gamma_+ \not k \gamma_\beta(\not k_1 - \not k)\gamma_\alpha)\,, \end{aligned} \qquad \text{(A.10)}$$

which gives

$$\begin{aligned} B_{II,\alpha\beta} &= \gamma_\alpha Tr(\gamma_+ \not k \gamma_\beta(\not k_1 - \not k)) - g_{\alpha+} Tr(\not r \not k \gamma_\beta(\not k_1 - \not k)) \\ &\quad +k_\alpha Tr(\not r \gamma_+ \gamma_\beta(\not k_1 - \not k)) - g_{\alpha\beta} Tr(\not r \gamma_+ \not k(\not k_1 - \not k)) \\ &\quad +(k_1 - k)_\alpha Tr(\not r \gamma_+ \not k \gamma_\beta)\,. \end{aligned} \qquad \text{(A.11)}$$

At the pole $k_-^{(1)}$, the first term of (A.11), after taking only the logarithmic divergent contribution and only the transverse component, becomes

$$\gamma_\alpha Tr(\gamma_+(\gamma_- k_+ - \underline{\gamma} \cdot \underline{k})\gamma_\beta(\gamma_+(k_1 - k)_- + \gamma_-(k_1 - k)_+ - \underline{\gamma} \cdot (\underline{k_1} - \underline{k}))) \xrightarrow{i,j} 0\,; \qquad \text{(A.12)}$$

the second term is always zero because $g_{i+} = 0$; the third term now becomes

$$4k_\alpha(r_+(k_1 - k)_\beta + g_{\beta+} r \cdot (k_1 - k) - r_\beta(k_1 - k)_+ \xrightarrow{i,j} -4k_i k_j r_+ = 2g_{ij}\underline{k}^2 r_+\,; \qquad \text{(A.13)}$$

the fourth term is now

$$\begin{aligned} & -4g_{ij}(r_+ k \cdot (k_1 - k) + r \cdot (k_1 - k)k_+ - k \cdot r(k_1 - k)_+) \\ &= -4g_{ij}(r_+ \frac{\underline{k}^2}{2\omega} - r_+ \frac{\underline{k}^2}{2} - \frac{\underline{k}^2}{2k_+} r_+(k_1 - k)_+) \\ &= -2g_{ij} r_+ \underline{k}^2 (\frac{1}{\omega} - 1 - \frac{1 - \omega}{\omega}) = 0\,; \end{aligned} \qquad \text{(A.14)}$$

and the last term is

$$-\,k_i Tr(\not r \gamma_+ \not k \gamma_j) = -4k_i(r_+ k_j) = 2g_{ij}\underline{k}^2 r_+\,. \qquad \text{(A.15)}$$

Thus we obtain

$$B^{(1)}_{II,ij} = 4g_{ij}\underline{k}^2 r_+ = 4k_{1+}g_{ij}\underline{k}^2\nu\,. \tag{A.16}$$

At the pole $k_-^{(2)}$, the first and second terms are both zero for the same reason; the third and fifth terms will take the same value as they do at the $k_-^{(1)}$ pole; for the fourth term we now have

$$\begin{aligned}
&-4g_{ij}(r_+k\cdot(k_1{-}k) + r\cdot(k_1{-}k)k_+ - k\cdot r(k_1{-}k)_+)\\
&= -4g_{ij}(r_+(-\frac{\underline{k}^2}{2(1{-}\omega)} + \frac{\underline{k}^2}{(1{-}\omega)}) + \frac{r_+k_+\underline{k}^2}{2k_{1+}(1{-}\omega)} - r_+(-\frac{\underline{k}^2}{2k_{1+}(1{-}\omega)})(k_1{-}k)_+\\
&= -2g_{ij}\underline{k}^2 r_+(\frac{1}{1{-}\omega} + \frac{\omega}{1{-}\omega} + 1)\\
&= -4g_{ij}\underline{k}^2 r_+\frac{1}{1{-}\omega}\,.
\end{aligned} \tag{A.17}$$

And we obtain

$$\begin{aligned}
B^{(2)}_{II,ij} &= 4k_{1+}g_{ij}\underline{k}^2\nu - 4k_{1+}g_{ij}\underline{k}^2\frac{\nu}{1{-}\omega}\\
&= -4k_{1+}g_{ij}\underline{k}^2\frac{\omega\nu}{1{-}\omega}\,.
\end{aligned} \tag{A.18}$$

Collecting terms we have

$$\begin{aligned}
B^{(1)}_{ij} &= B^{(1)}_{I,ij} - B^{(1)}_{II,ij} = 4k_{1+}g_{ij}\underline{k}^2(2\omega{-}1{-}\nu)\\
B^{(2)}_{ij} &= B^{(2)}_{I,ij} - B^{(2)}_{II,ij} = -4k_{1+}g_{ij}\underline{k}^2\,\frac{\omega^2{+}(1{-}\omega)^2{-}\omega\nu}{1-\omega}\,.
\end{aligned} \tag{A.19}$$

Substitute this back into (A.4) we have

$$\begin{aligned}
\gamma^{ij}_{qg} &= -\frac{\alpha_s}{2\pi}g_{ij}\int\frac{d\underline{k}^2}{\underline{k}^2}\Bigg(\int_0^{r_+}\frac{dk_+k_+}{r_+}(2k-r)_+^{J-n-1}r_+^n(2\omega{-}1{-}\nu)\\
&\qquad -\int_{r_+}^{k_{1+}}\frac{dk_+(k_1-k)_+}{(k_1-r)_+}(2k-r)_+^{J-n-1}r_+^n\frac{\omega^2{+}(1{-}\omega)^2{-}\omega\nu}{1-\omega}\Bigg)\\
&= \frac{\alpha_s}{2\pi}\frac{1}{2}2g_{ij}k_{1+}^J\nu^n\int\frac{d\underline{k}^2}{\underline{k}^2}\,(\int_0^{\nu}\frac{d\omega\omega}{\nu}(2\omega-\nu)^{J-n-1}(2\omega{-}1{-}\nu)\\
&\qquad -\int_{\nu}^{1}\frac{d\omega}{1{-}\nu}(2\omega-\nu)^{J-n-1}(\omega^2{+}(1{-}\omega)^2{-}\omega\nu)\Bigg)\,.
\end{aligned} \tag{A.20}$$

After factoring out the logarithmic divergence in $\underline{k}^2$ we again recover the result from (4.53)

$$\begin{aligned}\gamma_{qg}^{ij} &= \frac{\alpha_s}{2\pi}\frac{1}{2}2g^{ij}k_{1+}^{J}\nu^n\int_0^1 d\omega(2\omega-\nu)^{J-n-1}\Big(-\Theta(\nu-\omega)\frac{1}{\nu}\omega(2\omega-1-\nu) \\ &\qquad +\Theta(\omega-\nu)\frac{1}{1-\nu}(\omega^2+(1-\omega)^2-\omega\nu)\Big) \ . \qquad \text{(A.21)}\end{aligned}$$

Appendix B

Gluon-Gluon Anomalous Dimension

The computation of the gluonic sector follows similarly. In this appendix we will show in some detail the calculation of the gluon-gluon anomalous dimension.

Using standard Feynman rules and light-cone (LC) gauge we can write down the value of the gluon-gluon triangle diagram (figure 21) as

$$\begin{aligned}\gamma_{gg}^{+,1} &= (-i)^3 g^2 \sum_{ij}(f_{ijk}f_{ik'j}) \int \frac{d^4k}{(2\pi)^4} \frac{\Gamma_{\gamma\alpha\delta}\, D_{\gamma\delta}(k_1-k)\,\Gamma_{\delta\nu\beta}}{(k^2+i\epsilon)((k_1-k)^2+i\epsilon)((k-r)^2+i\epsilon)} \\ &\quad \cdot D_{\beta\beta'}(k-r)\, O^{(J,n)}_{g,\beta'\alpha'}(k,k-r)\, D_{\alpha'\alpha}(k) \qquad \text{(B.1)}\end{aligned}$$

where the gluon vertex $O_g^{(J,n)}$ is again taken from (4.23) and Γ denotes the triple-gluon vertex.

Performing the color algebra with

$$\sum_{ij}(f_{ijk}f_{ik'j}) = -C_A\delta_{k,k'} \qquad \text{(B.2)}$$

and setting the index $k = k'$ we arrive at

$$\gamma_{gg}^{+,1} = -ig^2 C_A \int \frac{d^4k}{(2\pi)^4} \frac{2(2k-r)_+^{J-n-2} r_+^n A^g_{\mu\nu}(k,r)}{(k^2+i\epsilon)((k_1-k)^2+i\epsilon)((k-r)^2+i\epsilon)} \tag{B.3}$$

where

$$A^g_{\mu\nu} = \Gamma_{\gamma\alpha\delta}\, D_{\gamma\delta}(k_1-k)\, \Gamma_{\delta\nu\beta} D_{\beta\beta'}(k-r)\, V^g_{\beta'\alpha'}(k,k-r)\, D_{\alpha'\alpha}(k) \tag{B.4}$$

with

$$D_{\gamma\delta}(k_1-k) = g_{\gamma\delta} - \frac{n_\gamma (k_1-k)_\delta + (k_1-k)_\gamma n_\delta}{n\cdot(k_1-k)} \tag{B.5}$$

and similar $D_{\beta\beta'}(k-r)$ and $D_{\alpha'\alpha}(k)$. Again the LC null four vector n is defined with properties $n^2 = 0$ and $n \cdot v = v_+$ for any four vector v.

The one point we need to pay extra attention to is the form of the gluon vertex in (4.23) and thus in (B.4). The full tensorial structure of the LC gluon vertex in the non-forward case, generalized from that in [14], is

$$V^g_{\beta'\alpha'}(k,k-r) = g_{\beta'\alpha'} n{\cdot}k\, n{\cdot}(k-r) - n{\cdot}(k-r)\, k'_\beta n'_\alpha - n{\cdot}k\, n'_\beta (k-r)'_\alpha + k{\cdot}(k-r)\, n'_\beta n'_\alpha \,, \tag{B.6}$$

and it is to be contracted with gluon lines $D_{\alpha'\alpha}(k)$ and $D_{\beta\beta'}(k-r)$.

We are now going to show explicitly that

$$\begin{aligned} D_{\beta\beta'}(k-r)\, V^g_{\beta'\alpha'}(k,k-r)\, D_{\alpha'\alpha}(k) &\equiv V^g_{\beta\alpha}(k,k-r) \\ D_{\beta\beta'}(k-r)\, g_{\beta'\alpha'} n{\cdot}k\, n{\cdot}(k-r)\, D_{\alpha'\alpha}(k) &\equiv V^g_{\beta\alpha}(k,k-r), \end{aligned} \tag{B.7}$$

thus we can substitute the full gluon vertex by its first term because of the LC projectors of the two gluon lines connected to the vertex and use (4.23) rather than the full expression in (B.4).

We split $V^g_{\beta'\alpha'}(k,k-r)$ into two parts, with the $g_{\beta'\alpha'}$ term being one part and the remaining three terms being the other. That is, we write

$$D_{\beta\beta'}(k-r)\,V^g_{\beta'\alpha'}(k,k-r)\,D_{\alpha'\alpha}(k) \equiv V^{(g,1)}_{\beta\alpha} - V^{(g,2)}_{\beta\alpha}\,, \tag{B.8}$$

where

$$\begin{aligned}
V^{(g,1)}_{\beta\alpha} &= D_{\beta\beta'}(k-r)\,g_{\beta'\alpha'}n{\cdot}kn{\cdot}(k-r)\,D_{\alpha'\alpha}(k)\,,\\
V^{(g,2)}_{\beta\alpha} &= D_{\beta\beta'}(k-r)\,(n{\cdot}(k-r)k_{\beta'}n_{\alpha'} + n{\cdot}kn_{\beta'}(k-r)_{\alpha'} - k{\cdot}(k-r)n_{\beta'}n_{\alpha'})\,D_{\alpha'\alpha}(k)\,.
\end{aligned}$$

We have

$$\begin{aligned}
V^{(g,1)}_{\beta\alpha} &= \left(g_{\beta\beta'} - \frac{n_\beta(k{-}r)_{\beta'} + (k{-}r)_\beta n_{\beta'}}{n\cdot(k-r)}\right)(g_{\beta'\alpha'}\,n{\cdot}kn{\cdot}(k{-}r))\,D_{\alpha'\alpha}(k)\\
&= (n{\cdot}kn{\cdot}(k{-}r)g_{\beta\alpha'} - n{\cdot}k(n_\beta(k{-}r)_{\alpha'} + (k{-}r)_\beta n_{\alpha'}))\left(g_{\alpha'\alpha} - \frac{n_{\alpha'}k_\alpha + k_{\alpha'}n_\alpha}{n\cdot k}\right)\\
&= n{\cdot}kn{\cdot}(k{-}r)g_{\beta\alpha} - n{\cdot}k(n_\beta(k{-}r)_\alpha + (k{-}r)_\beta n_\alpha) - n{\cdot}(k{-}r)(n_\beta k_\alpha + k_\beta n_\alpha)\\
&\quad + n{\cdot}(k{-}r)n_\beta k_\alpha + k{\cdot}(k-r)n_\beta n_\alpha + n^2(k{-}r)_\beta k_\alpha + n{\cdot}k(k{-}r)_\beta n_\alpha\\
&= g_{\beta\alpha}n{\cdot}kn{\cdot}(k-r) - n{\cdot}(k-r)k_\beta n_\alpha - n{\cdot}kn_\beta(k-r)_\alpha + k{\cdot}(k-r)n_\beta n_\alpha\\
&\equiv V^g_{\beta\alpha}(k,k-r)\,,
\end{aligned} \tag{B.9}$$

and

$$\begin{aligned}
V^{(g,2)}_{\beta\alpha} &= \left(g_{\beta\beta'} - \frac{n_\beta(k{-}r)_{\beta'} + (k{-}r)_\beta n_{\beta'}}{n\cdot(k-r)}\right)(n{\cdot}(k-r)k_{\beta'}n_{\alpha'} + n{\cdot}kn_{\beta'}(k-r)_{\alpha'}\\
&\qquad -k{\cdot}(k{-}r)n_{\beta'}n_{\alpha'})\,D_{\alpha'\alpha}(k)\\
&= (n{\cdot}(k{-}r)k_\beta n_{\alpha'} + n{\cdot}kn_\beta(k{-}r)_{\alpha'} - 2k{\cdot}(k{-}r)n_\beta n_{\alpha'} - n{\cdot}k(k{-}r)_\beta n_{\alpha'}\\
&\quad - \frac{n{\cdot}k}{n{\cdot}(k{-}r)}(n{\cdot}(k{-}r)n_\beta(k{-}r)_{\alpha'} + n^2(k{-}r)_\beta(k{-}r)_{\alpha'})\\
&\quad + \frac{k{\cdot}(k{-}r)}{n{\cdot}(k{-}r)}(n{\cdot}(k{-}r)n_\beta n_{\alpha'} + n^2(k{-}r)_\beta n_{\alpha'}))\,D_{\alpha'\alpha}(k)
\end{aligned}$$

$$\begin{aligned}
&= \; (n\cdot(k-r)k_\beta n_{\alpha'} + n\cdot k n_\beta (k-r)_{\alpha'} - k\cdot(k-r)n_\beta n_{\alpha'} - k\cdot(k-r)n_\beta n_{\alpha'} \\
&\quad -n\cdot k(k-r)_\beta n_{\alpha'} - n\cdot k n_\beta (k-r)_{\alpha'} + k\cdot(k-r)n_\beta n_{\alpha'})\, D_{\alpha'\alpha}(k) \\
&= \; (n\cdot(k-r)k_\beta n_{\alpha'} - 2k\cdot(k-r)n_\beta n_{\alpha'} - n\cdot k(k-r)_\beta n_{\alpha'} + k\cdot(k-r)n_\beta n_{\alpha'}) \\
&\qquad\qquad \cdot \left(g_{\alpha'\alpha} - \frac{n_{\alpha'}k_\alpha + k_{\alpha'}n_\alpha}{n\cdot k} \right) \\
&= \; (n\cdot(k-r)k_\beta n_\alpha - 2k\cdot(k-r)n_\beta n_\alpha - n\cdot k(k-r)_\beta n_\alpha + k\cdot(k-r)n_\beta n_\alpha \\
&\quad -\frac{n\cdot(k-r)}{n\cdot k}(n^2 k_\beta k_\alpha + n\cdot k k_\beta k_\alpha) + \frac{2k\cdot(k-r)}{n\cdot k}(n^2 k_\beta k_\alpha + n\cdot k k_\beta k_\alpha) \\
&\quad +(n^2(k-r)_\beta k_\alpha + n\cdot k(k-r)_\beta n_\alpha) - \frac{k\cdot(k-r)}{n\cdot k}(n^2 n_\beta n_\alpha + n\cdot k n_\beta n_\alpha)) \\
&= \; n\cdot(k-r)k_\beta n_\alpha - k\cdot(k-r)n_\beta n_\alpha - n\cdot k(k-r)_\beta n_\alpha - n\cdot(k-r)k_\beta n_\alpha \\
&\quad +2k\cdot(k-r)n_\beta n_\alpha + n\cdot k(k-r)_\beta n_\alpha - k\cdot(k-r)n_\beta n_\alpha \\
&\equiv \; 0\ . \qquad\qquad (B.10)
\end{aligned}$$

It is now clear that we have established (B.7), and explicitly we have

$$\begin{aligned}
&D_{\beta\beta'}(k-r)V^g_{\beta'\alpha'}(k,k-r)\,D_{\alpha'\alpha}(k) = V^g_{\beta\alpha}(k,k-r) \\
&\quad = g_{\beta\alpha} n\cdot k n\cdot(k-r) - n\cdot(k-r)k_\beta n_\alpha - n\cdot k n_\beta (k-r)_\alpha + k\cdot(k-r)n_\beta n_\alpha\, .
\end{aligned} \qquad (B.11)$$

The other factors in the definition of $A^g_{\mu\nu}$ (equation B.4) are the triple-gluon vertices involved in the diagram:

$$\begin{aligned}
\Gamma_{\gamma\alpha\mu} &= g_{\gamma\alpha}(2k-k_1)_\mu - g_{\alpha\mu}(k_1+k)_\gamma + g_{\mu\gamma}(2k_1-k)_\alpha \equiv -\Gamma_{\gamma\mu\alpha}\, , \\
\Gamma_{\delta\nu\beta} &= g_{\delta\nu}(2k_1-k-r)_\beta - g_{\nu\beta}(2r-k_1-k)_\delta + g_{\beta\delta}(2k-k_1-r)_\nu \equiv -\Gamma_{\beta\nu\delta}\, . \quad (B.12)
\end{aligned}$$

Thus we can rewrite (B.4) as, since $D_{\gamma\delta} = D_{\delta\gamma}$,

$$A^g_{\mu\nu} \;=\; \Gamma_{\gamma\alpha\mu}\, D_{\gamma\delta}(k_1-k)\,\Gamma_{\delta\nu\beta} D_{\beta\beta'}(k-r)\, V^g_{\beta'\alpha'}(k,k-r)\, D_{\alpha'\alpha}(k)$$

$$
\begin{aligned}
&= \Gamma_{\gamma\alpha\delta}\, D_{\gamma\delta}(k_1-k)\, \Gamma_{\delta\nu\beta} V^g_{\beta\alpha}(k,k-r) \\
&= (-\Gamma_{\delta\nu\beta}) D_{\delta\gamma}(-\Gamma_{\gamma\alpha\mu}) V^g_{\beta\alpha} \equiv \Gamma_{\beta\nu\delta} D_{\delta\gamma} \Gamma_{\gamma\mu\alpha} V^g_{\beta\alpha}\,.
\end{aligned}
\tag{B.13}
$$

Expanding the above we obtain the full expression of the "gluon triangle" as

$$
\begin{aligned}
& A^g_{\mu\nu} \equiv \Gamma_{\beta\nu\delta} D_{\delta\gamma} \Gamma_{\gamma\mu\alpha} V^g_{\beta\alpha} \\
& = n\cdot k n\cdot (k-r)\,[\,(\,(k-2k_1)\cdot(k-2k_1+r) + (k_1+k)\cdot(k_1+k-2r)\,)\, g_{\mu\nu} \\
& \quad + (k_1+k-2r)_\mu (k-2k_1)_\nu + (k-2k_1+r)_\mu (k_1+k)_\nu \\
& \quad + (k-2k_1)_\mu (k_1-2k+r)_\nu + (k_1+k)_\mu (k_1-2k+r)_\nu \\
& \quad + (k_1-2k)_\mu (k_1+k-2r)_\nu + (k_1-2k)_\mu (k-2k_1+r)_\nu + 4(k_1-2k)_\mu (k_1-2k+r)_\nu\,] \\
& - \frac{n\cdot k n\cdot (k-r)}{n\cdot(k_1-k)}\,[((k_1+k)\cdot(k_1-k) n\cdot(k_1+k-2r) + (k_1-k)\cdot(k_1+k-2r)\, n\cdot(k_1+k)) g_{\mu\nu} \\
& \quad + n\cdot(k_1+k-2r)\,(\,(k_1-k)_\mu (k-2k_1)_\nu + (k_1-2k)_\mu (k_1-k)_\nu\,) \\
& \quad + n\cdot(k_1+k)\,(\,(k_1-k)_\mu (k_1-2k+r)_\nu + (k-2k_1+r)_\mu (k_1-k)_\nu\,) \\
& \quad + n\cdot(k-2k_1)(k_1-k)_\mu (k_1-2k+r)_\nu + n\cdot(k-2k_1+r)(k_1-2k)_\mu (k_1-k)_\nu \\
& \quad + 2n\cdot(k_1-k)(k_1-2k)_\mu (k_1-2k+r)_\nu \\
& \quad + (\,(k_1-k)\cdot(k_1+k) + (k_1-k)\cdot(k-2k_1)\,)\, n_\mu (k_1-2k+r)_\nu \\
& \quad + (k_1-k)\cdot(k_1+k-2r) n_\mu (k-2k_1)_\nu + (k-2k_1)\cdot(k-2k_1+r) n_\mu (k_1-k)_\nu \\
& \quad + (\,(k_1-k)\cdot(k_1+k-2r) + (k_1-k)\cdot(k-2k_1+r)\,)\,(k_1-2k)_\mu n_\nu \\
& \quad + (k-2k_1)\cdot(k-2k_1+r)(k_1-k)_\mu n_\nu + (k_1-k)\cdot(k_1+k)(k-2k_1+r)_\mu n_\nu\,] \\
& - n\cdot(k-r)\,[\,n\cdot(k-2k_1)\; k\cdot(k-2k_1+r) g_{\mu\nu} \\
& \quad + n\cdot(k-2k_1)(k_\mu (k_1-2k+r)_\nu + (k_1+k-2r)_\mu k_\nu) \\
& \quad + n\cdot(k_1+k-2r)(k_1-2k)_\mu k_\nu + n\cdot k (k_1-2k)_\mu (k_1-2k+r)_\nu \\
& \quad + (k_1+k)\cdot(k_1+k-2r)\, n_\mu k_\nu + k\cdot(k-2k_1+r) n_\mu (k_1+k)_\nu + k\cdot(k_1+k) n_\mu (k_1-2k+r)_\nu \\
& \quad + k\cdot(k-2k_1+r)(k_1-2k)_\mu n_\nu\,]
\end{aligned}
$$

$$
\begin{aligned}
&+\frac{n\cdot(k-r)}{n\cdot(k_1-k)}\,[\,n\cdot(k_1+k-2r)\,(\,n\cdot(k-2k_1)(k_1-k)_\mu k_\nu+n\cdot(k_1-k)(k_1-2k)_\mu k_\nu\,)\\
&\quad+\,(\,n\cdot(k-2k_1)(k_1-k)\cdot(k_1+k-2r)+n\cdot(k_1+k-2r)(k_1-k)\cdot(k_1+k)\\
&\qquad+n\cdot(k_1+k)(k_1-k)\cdot(k_1+k-2r)\,)\,n_\mu k_\nu\\
&\quad+k\cdot(k-2k_1+r)\,(\,(k_1-k)\cdot(k_1+k)\,n_\mu n_\nu+(\,n\cdot(k_1+k)+n\cdot(k-2k_1)\,)n_\mu(k_1-k)_\nu\\
&\qquad+n\cdot(k-2k_1)(k_1-k)_\mu n_\nu+n\cdot(k_1-k)(k_1-2k)_\mu n_\nu\,)\\
&\quad+(\,(\,n\cdot(k-2k_1)\,k\cdot(k_1-k)+n\cdot k\,(k_1-k)\cdot(k_1+k)+n\cdot(k_1+k)\,k\cdot(k_1-k)\,)\,n_\mu\\
&\qquad+n\cdot k\,(\,n\cdot(k-2k_1)(k_1-k)_\mu+n\cdot(k_1-k)(k_1-2k)_\mu\,)\,)\,(k_1-2k+r)_\nu\,]\\
&-n\cdot k\,[\,n\cdot(k-2k_1+r)\,(k-2k_1)\cdot(k-r)\,g_{\mu\nu}+(k-2k_1)\cdot(k-r)\,n_\mu(k_1-2k+r)_\nu\\
&\quad+n\cdot(k-2k_1+r)\,(\,(k_1-2k)_\mu(k-r)_\nu+(k-r)_\mu(k_1+k)_\nu\,)\\
&\quad+(\,n\cdot(k_1+k)\,(k-r)_\mu+n\cdot(k-r)\,(k_1-2k)_\mu\,)\,(k_1-2k+r)_\nu\\
&\quad+(\,(k-2k_1)\cdot(k-r)\,(k_1+k-2r)_\mu+(k_1+k)\cdot(k_1+k-2r)\,(k-r)_\mu\\
&\qquad+(k-r)\cdot(k_1+k-2r)\,(k_1-2k)_\mu\,)\,n_\nu\,]\\
&+\frac{n\cdot k}{n\cdot(k_1-k)}\,[\,(\,(k-2k_1)\cdot(k-r)\,(\,n\cdot(k_1+k-2r)\,(k_1-k)_\mu+(k_1-k)\cdot(k_1+k-2r)\,n_\mu\,)\\
&\qquad+(\,n\cdot(k_1+k-2r)\,(k_1-k)\cdot(k_1+k)+n\cdot(k_1+k)\,(k_1-k)\cdot(k_1+k-2r)\,)\,(k-r)_\mu\\
&\qquad+(\,n\cdot(k_1+k-2r)\,(k_1-k)\cdot(k-r)+n\cdot(k-r)\,(k_1-k)\cdot(k_1+k-2r)\,)\,(k_1-2k)_\mu\,)\,n_\nu\\
&\quad+n\cdot(k-2k_1+r)\,(\,(\,n\cdot(k_1+k)\,(k-r)_\mu+n\cdot(k-r)\,(k_1-2k)_\mu\\
&\qquad\qquad\qquad\qquad+(k-2k_1)\cdot(k-r)\,n_\mu\,)\,(k_1-k)_\nu\\
&\qquad+(\,(k-2k_1)\cdot(k-r)\,(k_1-k)_\mu+(k_1-k)\cdot(k_1+k)\,(k-r)_\mu\\
&\qquad\qquad\qquad\qquad+(k_1-k)\cdot(k-r)\,(k_1-2k)_\mu\,)\,n_\nu\,)\\
&\quad+n\cdot(k_1-k)\,(\,n\cdot(k_1+k)\,(k-r)_\mu+n\cdot(k-r)\,(k_1-2k)_\mu\\
&\qquad+(k-2k_1)\cdot(k-r)\,n_\mu\,)\,(k_1-2k+r)_\nu\,]\\
&+k\cdot(k-r)\,[\,n\cdot(k-2k_1)\,n\cdot(k-2k_1+r)\,g_{\mu\nu}+(k_1+k)\cdot(k_1+k-2r)\,n_\mu n_\nu
\end{aligned}
$$

$$
\begin{aligned}
&\quad + n\cdot(k-2k_1+r)\, n_\mu(k_1+k)_\nu + (\, n\cdot(k-2k_1) + n\cdot(k_1+k)\,)\, n_\mu(k_1-2k+r)_\nu \\
&\quad + n\cdot(k-2k_1)\, (k_1+k-2r)_\mu n_\nu + (\, n\cdot(k_1+k-2r) + n\cdot(k-2k_1+r)\,)\, (k_1-2k)_\mu n_\nu \,] \\
&- \frac{k\cdot(k-r)}{n\cdot(k_1-k)}\, [\, (\, n\cdot(k-2k_1) + n\cdot(k_1+k)\,)\, n\cdot(k-2k_1+r)\, n_\mu(k_1-k)_\nu \\
&\quad + (\, n\cdot(k-2k_1) + n\cdot(k_1+k)\,)\, n\cdot(k_1-k)\; n_\mu(k_1-2k+r)_\nu \\
&\quad + (n\cdot(k_1+k-2r)+n\cdot(k-2k_1+r))\, (n\cdot(k-2k_1)\, (k_1-k)_\mu + n\cdot(k_1-k)\, (k_1-2k)_\mu)\, n_\nu \\
&\quad + (\, (\, n\cdot(k-2k_1) + n\cdot(k_1+k)\,)\, (k_1-k)\cdot(k_1+k-2r) \\
&\qquad + (\, n\cdot(k_1+k-2r) + n\cdot(k-2k_1+r)\,)\, (k_1-k)\cdot(k_1+k)\;)\; n_\mu n_\nu \,]\,, \qquad \text{(B.14)}
\end{aligned}
$$

where we have used $n\cdot n = n^2 = 0$.

Similar to the calculation of γ_{qq} we perform the k_- integral in (B.3) first. The discussion of different regions of k_+ and the corresponding pole values goes almost identical. We have

$$
\begin{aligned}
\gamma_{gg}^{+,1} &= -ig^2 C_A \int \frac{d^4k}{(2\pi)^4} \frac{2(2k-r)_+^{J-n-2} r_+^n A_{\mu\nu}^g(k,r)}{(k^2+i\epsilon)((k_1-k)^2+i\epsilon)((k-r)^2+i\epsilon)} \\
&= -i\frac{\alpha_s C_A}{(2\pi)^2} \int \frac{d\underline{k}^2}{\underline{k}^4} \left(\frac{-2\pi i}{r_+ k_{1+}} \int_0^{r_+} dk_+ k_+ A_{\mu\nu}^{g,(1)} (2k-r)_+^{J-n-2} r_+^n \right. \\
&\qquad \left. + \frac{2\pi i}{-k_{1+}(k_1-r)_+} \int_{r_+}^{k_{1+}} dk_+ (k_1-k)_+ A_{\mu\nu}^{g,(2)} (2k-r)_+^{J-n-2} r_+^n \right) \\
&= -\frac{\alpha_s C_A}{2\pi} \int \frac{d\underline{k}^2}{\underline{k}^4} \left(\int_0^{r_+} \frac{dk_+ k_+}{r_+ k_{1+}} (2k-r)_+^{J-n-2} r_+^n A_{\mu\nu}^{g,(1)} \right. \\
&\qquad \left. + \int_{r_+}^{k_{1+}} \frac{dk_+ (k_1-k)_+}{-k_{1+}(k_1-r)_+} (2k-r)_+^{J-n-2} r_+^n A_{\mu\nu}^{g,(2)} \right) \\
&\equiv \gamma_{gg}^{1,(1)} + \gamma_{gg}^{1,(2)}\,, \qquad \text{(B.15)}
\end{aligned}
$$

where $A^{g,(1)}$ and $A^{g,(2)}$ are $A_{\mu\nu}^g(k,r)$ evaluated at the poles $k_-^{(1)}$ and $k_-^{(2)}$, respectively, and again we have

$$\begin{aligned} k_-^{(1)} = \frac{\underline{k}^2}{2k_+} = \frac{\underline{k}^2}{2k_{1+}\omega} \qquad & \& \quad k^2 = 2k_+k_- - \underline{k}^2 = 0, \\ k_-^{(2)} = \frac{\underline{k}^2}{2(k_+ - k_{1+})} = \frac{-\underline{k}^2}{2k_{1+}(1-\omega)} \quad & \& \quad k^2 = -\frac{\underline{k}^2}{(1-\omega)}. \end{aligned} \tag{B.16}$$

To extract the gluon-gluon anomalous dimension, we do not need to calculate further the full expression B.14. Rather, the projection of $A^g_{\mu\nu}$ onto the transverse directions will suffice. That is, by effectively treating the gluon vertex to be

$$V^g_{ij} = g_{ij} n\cdot k\, n\cdot(k-r)\ . \tag{B.17}$$

we only need to keep in equation B.14 those terms proportional to g_{ij} and $k_i k_j \leftrightarrow -\frac{1}{2} g_{ij} \underline{k}^2$. Thus we define

$$A^g_{ij} \equiv g_{ij} A_I + k_i k_j A_{II} \tag{B.18}$$

and calculate A_I and A_{II} separately. Note as stated in the main text, we may choose k_1 and r such that only their plus components are non-vanishing, that is,

$$k_1 = (k_+, 0, \underline{0})\,, \qquad r = (r_+, 0, \underline{0})\,, \tag{B.19}$$

which also leads to

$$k_1 \cdot r = k_1^2 = r^2 = 0\ . \tag{B.20}$$

Reading off (and simplifying) from the full expression of $A^g_{\mu\nu}$ we arrive at

$$\begin{aligned} A_I \;=\; & n\cdot k n\cdot(k-r)(2k^2 - 2k_1\cdot k - k\cdot r) \\ & + \frac{n\cdot k n\cdot(k-r)}{n\cdot(k_1-k)}(2k^2 n\cdot(k_1+k) - 2k^2 n\cdot r - 2n\cdot(k_1+k)k\cdot r) \\ & - n\cdot(k-r)n\cdot(k-2k_1)(k^2 - 2k_1\cdot k + k\cdot r) - n\cdot k n\cdot(k-2k_1)(k^2 - 2k_1\cdot k) \\ & - n\cdot k(n\cdot r(k^2 - 2k_1\cdot k - k\cdot r) - n\cdot(k-2k_1)k\cdot r) \\ & + (k^2 - k\cdot r)((n\cdot(k-2k_1))^2 + n\cdot r(k-2k_1))\ . \end{aligned} \tag{B.21}$$

Same as the discussion of the quark-quark anomalous dimension, we are only interested in the leading powers of $\underline{k}$, which is also quadratic, and, from the pole values, can only come from terms proportional to either k^2 or k_-. Keeping only such terms we obtain

$$A_I = k^2 A_I^a + (k^2 - 2k_1{\cdot}k) A_I^b + r_+ k_- A_I^c \,, \tag{B.22}$$

where

$$\begin{aligned} A_I^a &= 2\left(k_+^2 - 2k_{1+}k_+ + 2k_{1+}^2 - k_{1+}r_+ + \frac{k_+^2 k_{1+} + k_+^3 - k_{1+}k_+r_+ - 2k_+^2 r_+ + k_+ r_+^2}{(k_1-k)_+}\right) \\ &= 2k_{1+}^2\left(\omega^2 - 2\omega + 2 - \nu + \frac{\omega^2 + \omega^3 - \omega\nu - 2\omega^2\nu + \omega\nu^2}{1-\omega}\right) \\ A_I^b &= 4k_{1+}k_+ - k_+^2 - 2k_{1+}r_+ - k_+r_+ \;=\; k_{1+}^2(4\omega - \omega^2 - 2\nu - \omega\nu) \\ A_I^c &= -2\left(k_+^2 - k_+r_+ - 2k_{1+}k_+ + 2k_{1+}^2 + \frac{k_+^3 + k_+^2 k_{1+} - k_+^2 r_+ - k_{1+}k_+r_+}{(k_1-k)_+}\right) \\ &= -2k_{1+}^2(\omega^2 - \omega\nu - 2\omega + 2 + \frac{\omega^3 + \omega^2 - \omega^2\nu - \omega\nu}{1-\omega}) \,. \end{aligned} \tag{B.23}$$

At the $k_-^{(1)}$ pole, from (B.16) we have

$$k^2 = 0\,, \quad 2k_1{\cdot}k = \frac{\underline{k}^2}{\omega}\,, \tag{B.24}$$

thus

$$\begin{aligned} A_I^{(1)} &= -\frac{\underline{k}^2}{\omega}A_I^b + \frac{\nu\underline{k}^2}{2\omega}A_I^c \\ &= -\frac{\underline{k}^2}{\omega}k_{1+}^2(4\omega - \omega^2 - 2\nu - \omega\nu) \\ &\quad -\frac{\nu\underline{k}^2}{\omega}k_{1+}^2(\omega^2 - \omega\nu - 2\omega + 2 + \frac{\omega^3 + \omega^2 - \omega^2\nu - \omega\nu}{1-\omega}) \\ &= -\frac{\underline{k}^2}{\omega}k_{1+}^2(4\omega - \omega^2 + \nu(\omega^2 - 3\omega - \omega\nu + \frac{\omega^3 + \omega^2 - \nu(\omega^2 + \omega)}{1-\omega})) \,. \end{aligned} \tag{B.25}$$

At the $k_-^{(2)}$ pole, on the other hand, from (B.16) we have

$$k^2 = 2k_1{\cdot}k = -\frac{\underline{k}^2}{1-\omega}\,, \tag{B.26}$$

and thus

$$
\begin{aligned}
A_I^{(2)} &= -\frac{\underline{k}^2}{1-\omega}A_I^a - \frac{\nu\underline{k}^2}{2(1-\omega)}A_I^c \\
&= -\frac{\underline{k}^2}{1-\omega}k_{1+}^2(2(\omega^2-2\omega+2-\nu+\frac{\omega^2+\omega^3-\omega\nu-2\omega^2\nu+\omega\nu^2}{1-\omega}) \\
&\quad -\nu(\omega^2-\omega\nu-2\omega+2+\frac{\omega^3+\omega^2-\omega^2\nu-\omega\nu}{1-\omega})) \\
&= -\frac{\underline{k}^2}{1-\omega}2k_{1+}^2(\omega^2-2\omega+2-2\nu+\frac{\omega^3+\omega^2-2\omega\nu(2\omega-\nu)}{1-\omega}) \ . \qquad \text{(B.27)}
\end{aligned}
$$

On the other hand, from the full expression of $A^g_{\mu\nu}$ (B.14) we have

$$
\begin{aligned}
A_{II} &= 10n{\cdot}kn{\cdot}(k-r) - 2n{\cdot}kn{\cdot}(k-r) - n{\cdot}(k-r)n{\cdot}(k+4r) - n{\cdot}kn{\cdot}(k-5r) \\
&\quad + \frac{n{\cdot}kn{\cdot}(k-r)}{n{\cdot}(k_1-k)}(n{\cdot}(k_1-k-2r) + n{\cdot}(k_1-k+2r)) \\
&= 8k_+^2 - 8k_+r_+ + 4r_+^2 \\
&= 4k_{1+}^2(2\omega^2-2\omega\nu+\nu^2) \ . \qquad \text{(B.28)}
\end{aligned}
$$

Note that A_{II} has the same value at both poles.

So now we have

$$
\begin{aligned}
A^g_{ij} &\equiv g_{ij}A_I + k_ik_jA_{II} \\
&= g_{ij}(A_I - \frac{1}{2}\underline{k}^2A_{II}) \\
&= g_{ij}(A_I - \underline{k}^2k_{1+}^2 2(2\omega^2-2\omega\nu+\nu^2)) \ . \qquad \text{(B.29)}
\end{aligned}
$$

Substitute the above result with the values of A_I at different poles into (B.15), we have

$$
\begin{aligned}
\gamma_{gg,ij}^{1,(1)} &= -\frac{\alpha_s C_A}{2\pi}\int\frac{d\underline{k}^2}{\underline{k}^4}\int_0^{r_+}\frac{dk_+k_+}{r_+k_{1+}}(2k-r)_+^{J-n-2}r_+^nA_{ij}^{g,(1)} \\
&= -\frac{\alpha_s C_A}{2\pi}\int\frac{d\underline{k}^2}{\underline{k}^4}\int_0^{\nu}\frac{d\omega\omega}{\nu}k_{1+}^{J-2}(2\omega-\nu)^{J-n-2}\nu^n g_{ij}(A_I^{(1)} - \frac{1}{2}\underline{k}^2A_{II})
\end{aligned}
$$

$$\begin{aligned}
&= -\frac{\alpha_s C_A}{2\pi}\int \frac{d\underline{k}^2}{\underline{k}^2} 2 g_{ij} k_{1+}^J \int_0^\nu d\omega (2\omega-\nu)^{J-n-2}\nu^n \frac{\omega}{2\nu} \\
&\quad \cdot(-\frac{1}{\omega}(4\omega-\omega^2+\nu(\omega^2-3\omega-\omega\nu+\frac{\omega^3+\omega^2-\nu(\omega^2+\omega)}{1-\omega})) \\
&\qquad -2(2\omega^2-2\omega\nu+\nu^2)) \\
&= \frac{\alpha_s C_A}{2\pi}\int \frac{d\underline{k}^2}{\underline{k}^2} 2 g_{ij} k_{1+}^J \int_0^\nu d\omega (2\omega-\nu)^{J-n-2}\nu^n \\
&\quad \cdot\frac{1}{2}\left(\frac{1}{\nu}(4\omega^3-\omega^2+4\omega)+\frac{\omega^3+\omega^2-2\omega^2\nu}{1-\omega}-3(\omega^2+\omega)\right) \qquad \text{(B.30)}
\end{aligned}$$

and

$$\begin{aligned}
\gamma_{gg,ij}^{1,(2)} &= -\frac{\alpha_s C_A}{2\pi}\int \frac{d\underline{k}^2}{\underline{k}^4}\int_{r_+}^{k_1 1+} \frac{dk_+(k_1-k)_+}{k_{1+}(k_1-r)_+}(2k-r)_+^{J-n-2} r_+^n A_{ij}^{g,(2)} \\
&= -\frac{\alpha_s C_A}{2\pi}\int \frac{d\underline{k}^2}{\underline{k}^4}\int_\nu^1 \frac{d\omega(1-\omega)}{1-\nu} k_{1+}^{J-2}(2\omega-\nu)^{J-n-2}\nu^n g_{ij}(A_I^{(2)}-\frac{1}{2}\underline{k}^2 A_{II}) \\
&= -\frac{\alpha_s C_A}{2\pi}\int \frac{d\underline{k}^2}{\underline{k}^2} 2 g_{ij} k_{1+}^J \int_\nu^1 d\omega (2\omega-\nu)^{J-n-2}\nu^n \frac{1-\omega}{2(1-\nu)} \\
&\quad \cdot(-\frac{2}{1-\omega}(\omega^2-2\omega+2-2\nu+\frac{\omega^3+\omega^2-2\omega\nu(2\omega-\nu)}{1-\omega}) \\
&\qquad -2(2\omega^2-2\omega\nu+\nu^2)) \\
&= \frac{\alpha_s C_A}{2\pi}\int \frac{d\underline{k}^2}{\underline{k}^2} 2 g_{ij} k_{1+}^J \int_\nu^1 d\omega (2\omega-\nu)^{J-n-2}\nu^n \\
&\quad \cdot\frac{1}{1-\nu}\left(\frac{2(1-\omega+\omega^2)^2+\nu(\nu(1+\omega^2)-2(1+\omega^3))}{1-\omega}\right)\ . \qquad \text{(B.31)}
\end{aligned}$$

Same as in the case of the quark-quark anomalous dimension, the ω integration is divergent at the end point $\omega = 1$. This divergence is again cancelled by self-energy graphs. However, in this case for each gluon propagator connecting to the vertex, there are two of them, a gluon loop and a quark loop (figure 22). Their values in the LC gauge can again be taken readily from [47] as

$$\begin{aligned}
Z_g(x) &= 1+\frac{\alpha_s}{2\pi}2C_A\frac{2}{\epsilon}(-I_0-\log|x|+\frac{11}{12}) \\
Z_q(x) &= 1+\frac{\alpha_s}{2\pi}2C_A\frac{2}{\epsilon}(-\frac{n_f}{18})\ . \qquad \text{(B.32)}
\end{aligned}$$

Again the self-energies depend on the longitudinal momentum fraction x, with I_0 the LC colinear divergence (see equation 4.51). Using the same argument as that was used for the quark-quark graph, we need add to the gluon-gluon anomalous dimension the following contribution from self-energy diagrams:

$$\frac{1}{2}(Z_g(1)+Z_q(1)+Z_g(1-\nu)+Z_q(1-\nu)) = \frac{\alpha_s}{2\pi}2C_A\frac{2}{\epsilon}(-I_0-\frac{1}{2}\log(1-\nu)+\frac{11}{12}-\frac{n_f}{18})\ . \tag{B.33}$$

Identifying $\frac{2}{\epsilon}$ as the logarithmic divergence in transverse momentum and that

$$\frac{11}{12}-\frac{n_f}{18} = \frac{1}{4N_c}(11-\frac{2}{3}n_f) \equiv \frac{1}{4N_c}b_0 \tag{B.34}$$

with b_0 the leading coefficient of the QCD $\beta-$function, we can see that we have shown that the gluon-gluon anomalous dimension is indeed given by the same expression of γ_{gg} as in (4.53), that is,

$$\begin{aligned}\gamma_{gg}^{ij} \ =\ & \frac{\alpha_s C_A}{2\pi}2g^{ij}k_{1+}^J\nu^n\int_0^1 d\omega(2\omega-\nu)^{J-n-2}\Big(\Theta(\nu-\omega)\frac{1}{2}(\frac{1}{\nu}(4\omega^3-\omega^2+4\omega) \\ & +(\frac{\omega^3+\omega^2-2\omega^2\nu}{1-\omega}-3(\omega^2+\omega)))+\frac{\Theta(\omega-\nu)}{1-\nu}(\frac{2(1-\omega+\omega^2)^2}{1-\omega}+\nu\frac{\nu(1+\omega^2)-2(1+\omega^3)}{1-\omega}) \\ & +(1-\nu)\delta(1-\omega)(-2I_0-\log(1-\nu)+\frac{b_0}{2C_A})\Big)\ .\end{aligned} \tag{B.35}$$

Appendix C

Gluon-Quark Anomalous Dimension

In this appendix we show the details of computing the gluon-quark anomalous dimension. The gluon-quark transition comes from the contribution of the second triangle diagram, as shown in detail in figure 19. By the same Feynman rules and in the LC gauge, we can write the value of the gluon-quark diagram as

$$\begin{aligned}
\gamma_{gq}^{+,1} &= (-ig)^2 \sum_{i,b} (T^i_{ab} T^i_{ba'}) \int \frac{d^4k}{(2\pi)^4} \gamma_\mu \frac{i}{\not{k}_1 - \not{k}} \gamma_\nu \frac{-iD_{\nu\beta}(k-r)}{(k-r)^2 + i\epsilon} \\
&\qquad \cdot O^{(J,n)(+)}_{g,\beta\alpha}(k, k-r) \frac{-iD_{\alpha\mu}(k)}{k^2 + i\epsilon} \\
&= ig^2 C_F \delta_{aa'} \int \frac{d^4k}{(2\pi)^4} \frac{2(2k-r)_+^{J-n-2} r_+^n B^g(k,r)}{(k^2 + i\epsilon)((k_1 - k)^2 + i\epsilon)((k-r)^2 + i\epsilon)} \,. \qquad \text{(C.1)}
\end{aligned}$$

Similar to the calculation of the quark-quark diagram (see section 4.2.1) we have set the color label of the two quark lines to be equal ($a = a'$), and defined

$$B^g(k,r) = \gamma_\mu (\not{k}_1 - \not{k}) \gamma_\nu D_{\nu\beta}(k-r) V^g_{\beta\alpha} D_{\alpha\mu}(k) \,. \qquad \text{(C.2)}$$

The LC gluon projectors are

$$\begin{aligned} D_{\nu\beta}(k-r) &= g_{\nu\beta} - \frac{n_\nu (k-r)_\beta + (k-r)_\nu n_\beta}{n\cdot(k-r)}\,, \\ D_{\alpha\mu}(k) &= g_{\alpha\mu} - \frac{n_\alpha k_\mu + k_\alpha n_\mu}{n\cdot k}\,. \end{aligned} \tag{C.3}$$

From our convention of the gluon vertex (4.23), by exactly the same argument as that for the gluon-gluon diagram (see the derivation of equation B.7 in appendix B) we have

$$\begin{aligned} &D_{\nu\beta}(k-r)\, V^g_{\beta\alpha}(k,k-r)\, D_{\alpha\mu}(k) \equiv V^g_{\nu\mu}(k,k-r) \\ &\quad = g_{\nu\mu} n\cdot k n\cdot(k-r) - n\cdot(k-r)k_\nu n_\mu - n\cdot n_\nu (k-r)_\mu + k\cdot(k-r) n_\nu n_\mu\,. \end{aligned} \tag{C.4}$$

By the same steps we followed before in computing the k_- integral and picking up different poles at different k_+ regions, we have

$$\begin{aligned} \gamma^{+,1}_{gq} &= \frac{\alpha_s C_F}{2\pi} \int \frac{d\underline{k}^2}{\underline{k}^4} \Bigg(\int_0^{r_+} \frac{dk_+ k_+}{r_+ k_{1+}} (2k-r)_+^{J-n-2} r_+^n B^{g,(1)} \\ &\qquad + \int_{r_+}^{k_{1+}} \frac{dk_+ (k_1-k)_+}{k_{1+}(k_1-r)_+} (2k-r)_+^{J-n-2} r_+^n B^{g,(2)} \Bigg) \\ &\equiv \gamma^{1,(1)}_{gq} + \gamma^{1,(2)}_{gq} \end{aligned} \tag{C.5}$$

where $B^{g,(1)}$ and $B^{g,(2)}$ are $B^g(k,r)$ evaluated at the poles $k_-^{(1)}$ and $k_-^{(2)}$, respectively, with

$$k_-^{(1)} = \frac{\underline{k}^2}{2k_+} = \frac{\underline{k}^2}{2k_{1+}\omega} \qquad \& \quad k^2 = 2k_+k_- - \underline{k}^2 = 0,$$

$$k_-^{(2)} = \frac{\underline{k}^2}{2(k_+ - k_{1+})} = \frac{-\underline{k}^2}{2k_{1+}(1-\omega)} \qquad \& \quad k^2 = -\frac{\underline{k}^2}{(1-\omega)}.$$

For the evaluation of $B^g(k,r)$ we have

$$\begin{aligned}
B^g(k,r) &= \gamma_\mu(\not{k}_1 - \not{k})\gamma_\nu D_{\nu\beta}(k-r)V^g_{\beta\alpha}D_{\alpha\mu}(k) \\
&= \gamma_\mu(\not{k}_1 - \not{k})\gamma_\nu V^g_{\nu\mu}(k,k-r) = \gamma_\mu(\not{k}_1 - \not{k})\gamma_\nu(g_{\nu\mu}n\cdot kn\cdot(k-r) \\
&\quad -n\cdot(k-r)k_\nu n_\mu - n\cdot n_\nu(k-r)_\mu + k\cdot(k-r)n_\nu n_\mu) \\
&= n\cdot kn\cdot(k-r)\gamma_\mu(\not{k}_1 - \not{k})\gamma_\mu - n\cdot k(\not{k} - \not{r})(\not{k}_1 - \not{k})\not{n} \\
&\quad -n\cdot(k-r)\not{n}(\not{k}_1 - \not{k})\not{k} + k\cdot(k-r)\not{n}(\not{k}_1 - \not{k})\not{n} \\
&= -2n\cdot kn\cdot(k-r)(\not{k}_1 - \not{k}) - n\cdot k(\not{k}\not{k}_1 - \not{r}\not{k}_1 + \not{r}\not{k} - k^2)\not{n} \\
&\quad -n\cdot(k-r)\not{n}(\not{k}_1\not{k} - k^2) + k\cdot(k-r)\not{n}(\not{k}_1 - \not{k})\not{n} \\
&= -2k_+(k-r)_+(\not{k}_1 - \not{k}) - k_+(\not{k}\not{k}_1 - \not{r}\not{k}_1 + \not{r}\not{k} - k^2)\gamma_+ \\
&\quad -(k-r)_+\gamma_+(\not{k}_1\not{k} - k^2) + (k^2 - k\cdot r)\gamma_+(\not{k}_1 - \not{k})\gamma_+\,.
\end{aligned} \tag{C.6}$$

As before we are interested in picking up the leading logarithmic divergence terms which are again quadratic and come from terms proportional to k_- (and k^2 for the pole $k_-^{(2)}$). Since γ_{gq} is related to the gluon-quark transition in evolution, the contributing terms will also have to have the structure of a quark vertex, that is, be proportional to γ_+ as well. We obtain

$$\begin{aligned}
B^g(k,r) &= -2k_+(k-r)_+(-k_-\gamma_+) - k_+(\gamma_+\gamma_-\gamma_+(k_-k_{1+}) - 0 - 0 - k^2\gamma_+) \\
&\quad -(k-r)_+(\gamma_+\gamma_-\gamma_+k_{1+}k_- - k^2\gamma_+) + (k^2 - k_-r_+)\gamma_+\gamma_-\gamma_+(k_1-k)_+ \\
&= 2k_+(k-r)_+k_-\gamma_+ - 2\gamma_+k_+k_-k_{1+} + \gamma_+k_+k^2 - 2(k-r)_+\gamma_+k_{1+}k_- \\
&\quad +(k-r)_+k^2\gamma_+ + 2k^2\gamma_+(k_1-k)_+ - 2(k_1-k)_+k_-r_+\gamma_+\ .
\end{aligned} \tag{C.7}$$

Thus, for the $k_-^{(1)}$ pole we have

$$\begin{aligned}
B^{g,(1)} &= \gamma_+ \Bigg(\frac{2k_+(k-r)_+}{2k_+} - \frac{2k_+k_{1+}}{2k_+} + 0 - \frac{2(k-r)_+k_{1+}}{2k_+} \\
&\quad - \frac{2(k_1-k)_+r_+}{2k_+} + 0 + 0 \Bigg) \underline{k}^2 \\
&= \gamma_+\underline{k}^2((k-r)_+ - k_{1+} - (k-r)_+\frac{k_{1+}}{k_+} - r_+\frac{(k_1-k)_+}{k_+} \\
&= \gamma_+\underline{k}^2k_{1+}(\omega-\nu-1-\frac{\omega-\nu}{\omega} - \nu\frac{1-\omega}{\omega}) \\
&= \frac{\underline{k}^2}{\omega}\gamma_+k_{1+}(\omega^2-2\omega)\ , \qquad \text{(C.8)}
\end{aligned}$$

while for the $k_-^{(2)}$ pole we obtain

$$\begin{aligned}
B^{g,(2)} &= \gamma_+ \Bigg(-\frac{2k_+(k-r)_+}{2(1-\omega)k_{1+}} + \frac{2k_+k_{1+}}{2(1-\omega)k_{1+}} - \frac{k_+}{1-\omega} + \frac{2(k-r)_+k_{1+}}{2(1-\omega)k_{1+}} \\
&\quad - \frac{(k-r)_+}{1-\omega} - \frac{2(k_1-k)_+}{1-\omega} + \frac{2(k_1-k)_+r_+}{2(1-\omega)k_{1+}} \Bigg) \underline{k}^2 \\
&= \frac{\gamma_+\underline{k}^2}{1-\omega}(-\omega(k-r)_+ + k_+ - k_+ + (k-r)_+ - (k-r)_+ - 2k_{1+}(1-\omega) + r_+(1-\omega)) \\
&= -\frac{\gamma_+\underline{k}^2k_{1+}}{1-\omega}(\omega^2-2\omega+2-\nu) \\
&= -\frac{\underline{k}^2}{1-\omega}k_{1+}\gamma_+(1+(1-\omega)^2-\nu)\ . \qquad \text{(C.9)}
\end{aligned}$$

Subsitute these back into (C.5) we have

$$\begin{aligned}
\gamma_{gq}^{+,1} &= \frac{\alpha_sC_F}{2\pi}\int\frac{d\underline{k}^2}{\underline{k}^4}\Bigg(\int_0^{r_+}\frac{dk_+k_+}{2r_+k_{1+}}(2k-r)_+^{J-n-2}r_+^n\frac{\underline{k}^2}{\omega}\gamma_+k_{1+}(\omega^2-2\omega) \\
&\qquad + \int_{r_+}^{k_{1+}}\frac{dk_+(k_1-k)_+}{2k_{1+}(k_1-r)_+}(2k-r)_+^{J-n-2}r_+^n(-\frac{\underline{k}^2}{1-\omega}k_{1+}\gamma_+(1+(1-\omega)^2-\nu))\Bigg) \\
&= \frac{\alpha_sC_F}{2\pi}\int\frac{d\underline{k}^2}{\underline{k}^2}\Bigg(\int_0^{\nu}\frac{d\omega\omega}{\nu}\frac{\omega^2-2\omega}{\omega}k_{1+}k_{1+}^{J-2}(2\omega-\nu)^{J-n-2}\nu_+^n\gamma_+ \\
&\quad - \int_{\nu}^{1}\frac{d\omega(1-\omega)}{1-\nu}\frac{1+(1-\omega)^2-\nu}{1-\omega}k_{1+}k_{1+}^{J-2}(2\omega-\nu)^{J-n-2}\nu_+^n\gamma_+\Bigg)\ , \qquad \text{(C.10)}
\end{aligned}$$

upon which we have recovered the result listed in (4.53), that is, after factoring out the logarithmic divergence in the transverse momentum, we obtain

$$\begin{aligned}\gamma_{gq}^{(+)} &= \frac{\alpha_s}{2\pi}C_F\,\gamma_+k_{1+}^{J-1}\nu^n\int_0^1 d\omega(2\omega-\nu)^{J-n-2}\left(\Theta(\nu-\omega)\frac{1}{\nu}(\omega^2-2\omega)\right.\\ &\qquad\left.-\Theta(\omega-\nu)\frac{1}{1-\nu}(1+(1-\omega)^2-\nu)\right)\ . \qquad\text{(C.11)}\end{aligned}$$

Figures



Figure 1: Diffractive Vector Meson Production

This is the diagrammatic representation of diffractive vector meson production (Part (a)). Part (b) is the two-gluon exchange diagram that has the dominate contribution when the double leading logarithmic approximation is taken.



Figure 2: Deeply Inelastic Scattering

This is the diagrammatic representation of the Deeply Inelastic Scattering process. This is a neutral current process via one photon exchange. The initial proton is $|p, \sigma\rangle$, the initial and final electron are k, λ and k', λ', respectively, and n labels collectively the different final states of the proton.



Figure 3: Forward Amplitude and Structure Function

This is the graphic representation of the DIS forward amplitude (a) and the DIS structure function (b). The difference lies in the cut (the dotted line) in the middle of (b) which puts all the intermediate particles it passes through on the mass shell.



Figure 4: Analyticity of the Forward Amplitude

This figure shows the analytic properties of the forward amplitude on the complex ω plane. The crosses are poles and the bold regions on the real axis are the branching cuts. The apmlitude has a convergence circle of unit radius. c is the original contour used in the dispersion integral. It lies completely within the unit circle and thus encloses no singularity.



Figure 5: Dispersion Integral Contour

This figure shows the contour c' of the dispersion integral of the forward amplitude. It is a continuous distortion from the c contour of figure 4. c' does not cross any cut nor does it enclose any singularity on the complex ω plane.



Figure 6: Integration Region and Change of Variables

This figure illustrates the change of variables and integration limits in section 1.4.2, where we show the equivalence between the DGLAP equation and the RGE from OPE analysis. The shaded area is the integration region.



Figure 7: Forward Operator Product Expansion

The operator product of the forward amplitude to the lowest order. The upper part of the righ hand side is the Wilson coefficient. The lower part is the reduced matrix element. We have only shown the leading quark-quark transition.


$\gamma_+ (\frac{k}{p})^{n-1}$
k
k
p
p
$\cdot \frac{(2\pi)\, 2E_p}{2\, p_+} = xP^f(x)$

Figure 8: The Lowest Order Quark Distribution Function

This is the quark distribution function $P^f(x)$ where we have put in explicitly the lowest order quark vertex in the light cone gauge. The dotted line is the cut that puts all the intermediate lines it passes through on the mass shell.



Figure 9: The Lowest Order Quark-Quark Diagram

This is the quark-quark diagram of the first order radiative correction to the quark distribution function. a, b are color indices and i labels the on-shell gluon.



Figure 10: Quark Self Energy Diagram

These are the two quark self energy diagrams that are needed to be added to figure 9 to cancel infrared divergence at the end points of integration.



Figure 11: The Quark-Gluon Mixing Graph

This is the graph that generates the anomalous dimension for quark to gluon mixing in forward DIS.



Figure 12: Non-forward Scattering Amplitude

This is the unequal mass double virtual Compton scattering diagram. Both q and q' are virtual photons and they are of different invariant masses. There is a non-zero four-momentum transfer r from the incoming virtual photon to the target proton.



Figure 13: Symmetry of the Non-forward Scattering Amplitude

Because the two diagrams shown are indeed identical after the momentum labeling change $\mu \leftrightarrow \nu,\ q \leftrightarrow -q'$, the non-forward amplitude is invariant under the same transformation.



Figure 14: Handbag Diagrams in QCD-Parton Picture

These are the so-called "handbag" diagrams contributing to the non-forward amplitude in a QCD-Parton picture. We have explicitly included the cross diagram.



Figure 15: Triangle Graphs for the Anomalous Dimensions

These are the four lowest order "triangle" diagrams contributing to the anomalous dimensions. The momentum labels of the four are the same (only the quark-quark diagram shows them explicitly).



Figure 16: The Quark-Quark Diagram of the Anomalous Dimensions

This is the detailed version of the first triangle graph, the quark-quark diagram. α and β are Lorentz indices; a, a', b and b' are color indices for the quarks while i labels the qluon line. The difference between this diagram and Figure 9 in the forward case, besides the non-forwardness, is that we are not putting the gluon on-shell.



Figure 17: The Poles and Contours of k_- Integration

The figures show, on the k_- complex plane, the poles and different contours of k_- integration in the four regions of its value (see section 4.2.1). The dots are the pole positions with their distances to the real k_- axis exaggerated.



Figure 18: The Tree Level Born Diagrams of the Non-forward Amplitude

These are the tree level (Born) terms contributing to the non-forward amplitude from which we can calculate the lowest order Wilson coefficients. We have explicitly included the cross term.



Figure 19: The Gluon-Quark Diagram of the Anomalous Dimensions

This is the detailed version of the second triangle graph, the gluon-quark diagram. α, β, μ and ν are Lorentz indices; a, a', and b are color indices for the quarks while i labels the gluon lines.



Figure 20: The Quark-Gluon Diagram of the Anomalous Dimensions

This is the detailed version of the third triangle graph, the quark-gluon diagram. α, β are Lorentz indices; a, b are color indices for the quarks while i and i' label the gluon lines.



Figure 21: The Gluon-Gluon Diagram of the Anomalous Dimensions

This is the detailed version of the last triangle graph–the gluon-gluon diagram. α, α', β, β', γ, δ, μ and ν are Lorentz indices; i, j, k and k' label the gluon lines.



Figure 22: The Gluon Self-Energy Diagram

These are the gluon self-energy diagrams that are needed to cancel the end-point divergences in the LC calculation. We need to include both the gluon loop and the quark loop contributions.